



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07H 21/04, C07K 14/705, C12N 15/09, 15/63, C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/11015</b> <b>(43) International Publication Date:</b> 2 March 2000 (02.03.00)																								
<b>(21) International Application Number:</b> PCT/US99/19351 <b>(22) International Filing Date:</b> 24 August 1999 (24.08.99)  <b>(30) Priority Data:</b> <table border="0"> <tr> <td>60/097,638</td> <td>24 August 1998 (24.08.98)</td> <td>US</td> </tr> <tr> <td>60/097,659</td> <td>24 August 1998 (24.08.98)</td> <td>US</td> </tr> <tr> <td>60/099,618</td> <td>9 September 1998 (09.09.98)</td> <td>US</td> </tr> <tr> <td>60/102,092</td> <td>28 September 1998 (28.09.98)</td> <td>US</td> </tr> <tr> <td>60/109,978</td> <td>25 November 1998 (25.11.98)</td> <td>US</td> </tr> <tr> <td>60/113,645</td> <td>23 December 1998 (23.12.98)</td> <td>US</td> </tr> <tr> <td>60/113,646</td> <td>23 December 1998 (23.12.98)</td> <td>US</td> </tr> <tr> <td>09/379,246</td> <td>23 August 1999 (23.08.99)</td> <td>US</td> </tr> </table> <b>(71) Applicant:</b> ALPHAGENE, INC. [US/US]; 260 West Cummings Park, Woburn, MA 01801 (US). <b>(72) Inventors:</b> VALENZUELA, Dario; 260 West Cummings Park, Woburn, MA 01801 (US). YUAN, Olive; 260 West Cummings Park, Woburn, MA 01801 (US). HOFFMAN, Heidi; 260 West Cummings Park, Woburn, MA 01801 (US). HALL, Jeff; 260 West Cummings Park, Woburn, MA 01801 (US). RAPIEJKO, Peter; 260 West Cummings Park, Woburn, MA 01801 (US).		60/097,638	24 August 1998 (24.08.98)	US	60/097,659	24 August 1998 (24.08.98)	US	60/099,618	9 September 1998 (09.09.98)	US	60/102,092	28 September 1998 (28.09.98)	US	60/109,978	25 November 1998 (25.11.98)	US	60/113,645	23 December 1998 (23.12.98)	US	60/113,646	23 December 1998 (23.12.98)	US	09/379,246	23 August 1999 (23.08.99)	US	<b>(74) Agent:</b> SPRUNGER, Suzanne, A.; American Home Products Corporation, Patent & Trademark Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
60/097,638	24 August 1998 (24.08.98)	US																								
60/097,659	24 August 1998 (24.08.98)	US																								
60/099,618	9 September 1998 (09.09.98)	US																								
60/102,092	28 September 1998 (28.09.98)	US																								
60/109,978	25 November 1998 (25.11.98)	US																								
60/113,645	23 December 1998 (23.12.98)	US																								
60/113,646	23 December 1998 (23.12.98)	US																								
09/379,246	23 August 1999 (23.08.99)	US																								
<b>(54) Title:</b> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM  <b>(57) Abstract</b>  Novel polynucleotides and the proteins encoded thereby are disclosed.																										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

5

This application is a continuation-in-part of the following applications:

- (1) provisional application Ser. No. 60/097,638 (GI 6908), filed August 24, 1998;
  - (2) provisional application Ser. No. 60/097,659 (GI 6909), filed August 24, 1998;
  - (3) provisional application Ser. No. 60/099,618 (GI 6910), filed September 9, 1998;
  - 10 (4) provisional application Ser. No. 60/102,092 (GI 6912), filed September 28, 1998;
  - (5) provisional application Ser. No. 60/109,978 (GI 6914), filed November 25, 1998;
  - (6) provisional application Ser. No. 60/113,645 (GI 6916), filed December 23, 1998; and
  - (7) provisional application Ser. No. 60/113,646 (GI 6917), filed December 23, 1998;
- all of which are incorporated by reference herein.

15

FIELD OF THE INVENTION

The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins.

20

BACKGROUND OF THE INVENTION

Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques  
25 clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader  
30 sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these proteins and the  
35 polynucleotides encoding them that the present invention is directed.

### SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb11\_1 deposited with the ATCC under  
10 accession number 98846;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;
- (e) a polynucleotide comprising the nucleotide sequence of a mature  
15 protein coding sequence of clone vb11\_1 deposited with the ATCC under accession number 98846;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;
- (g) a polynucleotide encoding a protein comprising the amino acid  
20 sequence of SEQ ID NO:2;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of  
25 (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:1.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934; the nucleotide sequence of the full-length



protein coding sequence of clone vb11\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vb11\_1 deposited with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846.

In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:2.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:1.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and

(ab) the nucleotide sequence of the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and

(bb) the nucleotide sequence of the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:1 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the  
15 cDNA sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:2;

25 (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:2. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:2.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5                   (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID  
10               NO:3 from nucleotide 201 to nucleotide 482;
- (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (e)     a polynucleotide encoding the full-length protein encoded by the  
15               cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;
- 20               (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i)     a polynucleotide encoding a protein comprising a fragment of the  
25               amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k)     a polynucleotide which encodes a species homologue of the protein  
30               of (h) or (i) above ;
- (l)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:3.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482; the nucleotide sequence of SEQ ID NO:3 from nucleotide 201 to nucleotide 482; the nucleotide sequence of the full-length protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:4.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:3.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(bb) the nucleotide sequence of the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:3 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 201 to nucleotide 482, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 201 to nucleotide 482, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 201 to nucleotide 482.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

- (b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:4. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:4, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:4.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 1195 to nucleotide 1527;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:5 from nucleotide 1468 to nucleotide 1527;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:5.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:5 from nucleotide 1195 to nucleotide 1527; the nucleotide sequence of SEQ ID NO:5 from nucleotide 1468 to nucleotide 1527; the nucleotide sequence of the full-length protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846.

In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising the amino acid sequence from amino acid 50 to amino acid 59 of SEQ ID NO:6.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:5.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (aa) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(ab) the nucleotide sequence of the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

10 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

15 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

20 (bb) the nucleotide sequence of the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:5 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5. Also preferably the polynucleotide isolated  
30 according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 1195 to nucleotide 1527, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from nucleotide 1195 to nucleotide 1527, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 1195 to



nucleotide 1527. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 1468 to nucleotide 1527, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from  
5 nucleotide 1468 to nucleotide 1527, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 1468 to nucleotide 1527.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:6;
- (b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
vb14\_1 deposited with the ATCC under accession number 98846;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:6. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:6, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising the amino acid sequence from amino acid 50 to amino acid 59 of SEQ ID NO:6.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:7 from nucleotide 109 to nucleotide 294;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:8;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:7.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294; the nucleotide sequence of SEQ ID NO:7 from  
25 nucleotide 109 to nucleotide 294; the nucleotide sequence of the full-length protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
30 of clone ve11\_1 deposited with the ATCC under accession number 98846. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 30 to amino acid 39 of SEQ ID NO:8.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:7.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and
    - (ab) the nucleotide sequence of the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and
    - (bb) the nucleotide sequence of the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7, and extending

contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:7 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the

5 cDNA sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294. Also preferably the polynucleotide isolated according to the above

10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 109 to nucleotide 294, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 109 to nucleotide 294, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 109 to nucleotide 294.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the
- 20 fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8. In further preferred

25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from

30 amino acid 30 to amino acid 39 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 22 to nucleotide 468;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 118 to nucleotide 468;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:9.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:9 from nucleotide 22 to nucleotide 468; the nucleotide sequence of SEQ ID NO:9 from nucleotide 118 to nucleotide 468; the nucleotide sequence of the full-length protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vf2\_1 deposited with

the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 69 to amino acid 78 of SEQ ID NO:10.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:9.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 30 (ba) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

- (bb) the nucleotide sequence of the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:9 to  
10 a nucleotide sequence corresponding to the 3' end of SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 22 to nucleotide 468, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of  
15 SEQ ID NO:9 from nucleotide 22 to nucleotide 468, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 22 to nucleotide 468. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 118 to nucleotide 468, and extending contiguously from a  
20 nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 118 to nucleotide 468, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 118 to nucleotide 468.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
25 consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
- (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vf2\_1  
30 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:10. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 69 to amino acid 78 of SEQ ID NO:10.

5           In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11;
- 10       (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641;
- (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vg2\_1 deposited with the ATCC under  
15       accession number 98846;
- (e)     a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (f)     a polynucleotide comprising the nucleotide sequence of a mature  
20       protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (h)     a polynucleotide encoding a protein comprising the amino acid  
25       sequence of SEQ ID NO:12;
- (i)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of  
30       (a)-(g) above;
- (k)     a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and



(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:11.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641; the nucleotide sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641; the nucleotide sequence of the full-length protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence from amino acid 248 to amino acid 257 of SEQ ID NO:12.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:11.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(bb) the nucleotide sequence of the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:11 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:12;

- (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:12. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:12, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence from amino acid 248 to amino acid 257 of SEQ ID NO:12.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:13 from nucleotide 416 to nucleotide 892;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:14;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:13.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892; the nucleotide sequence of SEQ ID NO:13 from nucleotide 416 to nucleotide 892; the nucleotide sequence of the full-length protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 80 to amino acid 89 of SEQ ID NO:14.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:13.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:13; and

5 (ab) the nucleotide sequence of the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

10 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15

(ba) SEQ ID NO:13; and

(bb) the nucleotide sequence of the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

20 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13, and  
25 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:13 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:13. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892, and extending contiguously from a nucleotide sequence  
30 corresponding to the 5' end of said sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 416 to nucleotide 892, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from nucleotide 416 to nucleotide 892, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 416 to nucleotide 892.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
- (b) a fragment of the amino acid sequence of SEQ ID NO:14, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:14. In further preferred  
15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence  
20 from amino acid 80 to amino acid 89 of SEQ ID NO:14.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
25 NO:15 from nucleotide 62 to nucleotide 1057;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057;
- (d) a polynucleotide comprising the nucleotide sequence of the full-  
30 length protein coding sequence of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vl1\_1 deposited with the ATCC under accession number 98846;
- 5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- 10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 15 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:15.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:15 from nucleotide 62 to nucleotide 1057; the nucleotide sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057; the nucleotide sequence of the full-length protein coding sequence of clone vl1\_1 deposited with the ATCC under accession number 98846; or the nucleotide sequence of a mature protein coding sequence of clone vl1\_1 deposited
- 25 with the ATCC under accession number 98846. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having
- 30 biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 161 to amino acid 170 of SEQ ID NO:16.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:15.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:15, but excluding the poly(A) tail at the  
10 3' end of SEQ ID NO:15; and
- (ab) the nucleotide sequence of the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 15 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
20 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and
- (bb) the nucleotide sequence of the cDNA insert of clone  
25 vl1\_1 deposited with the ATCC under accession number 98846;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

- 30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:15 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15. Also preferably the



polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 62 to nucleotide 1057, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 62 to nucleotide 1057, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 62 to nucleotide 1057. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:16. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 161 to amino acid 170 of SEQ ID NO:16.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 74 to nucleotide 529;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:17.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:17 from nucleotide 74 to nucleotide 529; the nucleotide sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529; the nucleotide sequence of the full-length protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838; or the nucleotide sequence of a mature protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert

30

of clone vk2\_1 deposited with the ATCC under accession number 98838. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, 5 most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 71 to amino acid 80 of SEQ ID NO:18.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ 10 ID NO:17.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize 15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and
    - (ab) the nucleotide sequence of the cDNA insert of clone 20 vk2\_1 deposited with the ATCC under accession number 98838;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 25 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 30 (ba) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:17 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17. Also preferably the
- 10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 74 to nucleotide 529, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 74 to nucleotide 529, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide
- 15 74 to nucleotide 529. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529, to a nucleotide sequence corresponding to the 3' end of
- 20 said sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:18;
- 25 (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;

the protein being substantially free from other mammalian proteins. Preferably such

30 protein comprises the amino acid sequence of SEQ ID NO:18. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 71 to amino acid 80 of SEQ ID NO:18.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19;
- (b)    a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170;
- (c)    a polynucleotide comprising the nucleotide sequence of SEQ ID  
10       NO:19 from nucleotide 1098 to nucleotide 3170;
- (d)    a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (e)    a polynucleotide encoding the full-length protein encoded by the  
15       cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (f)    a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862;
- 20       (g)    a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (h)    a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (i)    a polynucleotide encoding a protein comprising a fragment of the  
25       amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (j)    a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k)    a polynucleotide which encodes a species homologue of the protein  
30       of (h) or (i) above ;
- (l)    a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:19.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170; the nucleotide sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170; the nucleotide sequence of the full-length protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862. In other preferred  
5      NO:19 from nucleotide 174 to nucleotide 3170; the nucleotide sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170; the nucleotide sequence of the full-length protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862. In other preferred  
10    embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862.

In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably  
15    twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 494 to amino acid 503 of SEQ ID NO:20.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:19.  
20    ID NO:19.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
25      in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
30      vb21\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and

(bb) the nucleotide sequence of the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:19 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:20;

- (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:20. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:20, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 494 to amino acid 503 of SEQ ID NO:20.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 74 to nucleotide 1453;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:21 from nucleotide 224 to nucleotide 1453;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;



- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- 5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:21.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:21 from nucleotide 74 to nucleotide 1453; the nucleotide sequence of SEQ ID NO:21 from nucleotide 224 to nucleotide 1453; the nucleotide sequence of the full-length protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 225 to amino acid 234 of SEQ ID NO:22.

15  
20  
25

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:21.

30 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 5 (aa) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and
- (ab) the nucleotide sequence of the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 10 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 15 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 20 vc35\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:21 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21. Also preferably the
- 30 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 74 to nucleotide 1453, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 74 to nucleotide 1453, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide

74 to nucleotide 1453. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 224 to nucleotide 1453, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from  
5 nucleotide 224 to nucleotide 1453, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 224 to nucleotide 1453.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:22;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:22. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:22, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 225 to amino acid 234 of SEQ ID NO:22.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:23 from nucleotide 243 to nucleotide 368;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:23.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368; the nucleotide sequence of SEQ ID NO:23  
25 from nucleotide 243 to nucleotide 368; the nucleotide sequence of the full-length protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
30 of clone vc36\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:24, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:24.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:23.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:23 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368. Also preferably the polynucleotide isolated according to the above  
10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 243 to nucleotide 368, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 243 to nucleotide 368, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 243 to nucleotide 368.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:24;
- (b) a fragment of the amino acid sequence of SEQ ID NO:24, the  
20 fragment comprising eight contiguous amino acids of SEQ ID NO:24; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:24. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:24, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence  
30 from amino acid 34 to amino acid 43 of SEQ ID NO:24.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25 from nucleotide 370 to nucleotide 1662;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc38\_1 deposited with the ATCC under accession number 98862;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc38\_1 deposited with the ATCC under accession number 98862;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:26;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:25.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:25 from nucleotide 370 to nucleotide 1662; the nucleotide sequence of the full-length protein coding sequence of clone vc38\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc38\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862.

In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:26, or a  
5 polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 210 to amino acid 219 of SEQ ID NO:26.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:25.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
15 consisting of:

(aa) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and

(ab) the nucleotide sequence of the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and

(bb) the nucleotide sequence of the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;



- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:25 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25 from nucleotide  
10 1662, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:25 from nucleotide 370 to nucleotide 1662, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:25 from nucleotide 370 to nucleotide 1662.

In other embodiments, the present invention provides a composition comprising  
15 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:26;
- (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
- 20 (c) the amino acid sequence encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:26. In further preferred embodiments, the present invention provides a protein comprising a fragment of the  
25 amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:26, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 210 to amino acid 219 of SEQ ID NO:26.

30 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:27.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365; the nucleotide sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365; the nucleotide sequence of the full-length protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc39\_1 deposited

with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:28.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:27.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 30 (ba) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

- (bb) the nucleotide sequence of the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:27 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:28;
- (b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:28. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:28.

5           In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29;
- 10           (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc40\_1 deposited with the ATCC under  
15           accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
20           protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25           sequence of SEQ ID NO:30;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30           (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:29.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066; the nucleotide sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066; the nucleotide sequence of the full-length protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 167 to amino acid 176 of SEQ ID NO:30.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:29.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and

(bb) the nucleotide sequence of the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:29 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:30;

- (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:30. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:30, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 167 to amino acid 176 of SEQ ID NO:30.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31 from nucleotide 38 to nucleotide 553;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:31 from nucleotide 104 to nucleotide 553;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:32;



- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:31.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:31 from nucleotide 38 to nucleotide 553; the nucleotide sequence of SEQ ID NO:31 from nucleotide 104 to nucleotide 553; the nucleotide sequence of the full-length protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:32, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 81 to amino acid 90 of SEQ ID NO:32.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:31.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 5 (aa) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and
- (ab) the nucleotide sequence of the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 10 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 15 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 20 vc46\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:31 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31. Also preferably the
- 30 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 38 to nucleotide 553, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from nucleotide 38 to nucleotide 553, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide

38 to nucleotide 553. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 104 to nucleotide 553, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from  
5 nucleotide 104 to nucleotide 553, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide 104 to nucleotide 553.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:32;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:32. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:32, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 81 to amino acid 90 of SEQ ID NO:32.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:33 from nucleotide 242 to nucleotide 2548;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:33.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548; the nucleotide sequence of SEQ ID NO:33 from nucleotide 242 to nucleotide 2548; the nucleotide sequence of the full-length protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:34, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 392 to amino acid 401 of SEQ ID NO:34.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:33.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vc49\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said primer(s) to human genomic DNA in  
30 conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:33 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548. Also preferably the polynucleotide isolated according to the above  
10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33 from nucleotide 242 to nucleotide 2548, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 242 to nucleotide 2548, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:33 from nucleotide 242 to nucleotide 2548.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
- (b) a fragment of the amino acid sequence of SEQ ID NO:34, the  
20 fragment comprising eight contiguous amino acids of SEQ ID NO:34; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:34. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:34, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence  
30 from amino acid 392 to amino acid 401 of SEQ ID NO:34.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:35.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776; the nucleotide sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776; the nucleotide sequence of the full-length protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc50\_1 deposited

with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:36, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:36.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:35.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and



- (bb) the nucleotide sequence of the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:35 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:36;
- (b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:36. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:36, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:36.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:37 from nucleotide 139 to nucleotide 1308;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc51\_1 deposited with the ATCC under  
15 accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
20 protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25 sequence of SEQ ID NO:38;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30 (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:37.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:37 from nucleotide 139 to nucleotide 1308; the nucleotide sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308; the nucleotide sequence of the full-length protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:38, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 190 to amino acid 199 of SEQ ID NO:38.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:37.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37; and

(bb) the nucleotide sequence of the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:37 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37 from nucleotide 139 to nucleotide 1308, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from nucleotide 139 to nucleotide 1308, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide 139 to nucleotide 1308. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:38;

- (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:38. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:38, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 190 to amino acid 199 of SEQ ID NO:38.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 21 to nucleotide 1142;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:39 from nucleotide 114 to nucleotide 1142;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:40;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:39.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:39 from nucleotide 21 to nucleotide 1142; the nucleotide sequence of SEQ ID NO:39 from nucleotide 114 to nucleotide 1142; the nucleotide sequence of the full-length protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862; or the nucleotide sequence of a mature protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 182 to amino acid 191 of SEQ ID NO:40.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:39.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 5 (aa) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and
- (ab) the nucleotide sequence of the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 10 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 15 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 20 vc52\_1 deposited with the ATCC under accession number 98862;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:39 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39. Also preferably the
- 30 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 21 to nucleotide 1142, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from nucleotide 21 to nucleotide 1142, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide

21 to nucleotide 1142. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 114 to nucleotide 1142, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from  
5 nucleotide 114 to nucleotide 1142, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide 114 to nucleotide 1142.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:40;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:40. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:40, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 182 to amino acid 191 of SEQ ID NO:40.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:41 from nucleotide 346 to nucleotide 1416;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886;



- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:41.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416; the nucleotide sequence of SEQ ID NO:41  
25 from nucleotide 346 to nucleotide 1416; the nucleotide sequence of the full-length protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
30 of clone vc33\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence from amino acid 229 to amino acid 238 of SEQ ID NO:42.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:41.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vc33\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 20 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (ba) SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said primer(s) to human genomic DNA in  
30 conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:41 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416. Also preferably the polynucleotide isolated according to the above  
10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 346 to nucleotide 1416, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 346 to nucleotide 1416, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 346 to nucleotide 1416.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
- (b) a fragment of the amino acid sequence of SEQ ID NO:42, the  
20 fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:42. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence  
30 from amino acid 229 to amino acid 238 of SEQ ID NO:42.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:44;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:43.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461; the nucleotide sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461; the nucleotide sequence of the full-length protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vc34\_1 deposited

with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence from amino acid 200 to amino acid 209 of SEQ ID NO:44.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:43.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

- (bb) the nucleotide sequence of the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ  
10 ID NO:43 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
15 of said sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461, and extending contiguously from a  
20 nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
25 consisting of:

- (a) the amino acid sequence of SEQ ID NO:44;
- (b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
30 vc34\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:44. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence from amino acid 200 to amino acid 209 of SEQ ID NO:44.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:45 from nucleotide 1922 to nucleotide 2350;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc47\_1 deposited with the ATCC under  
15 accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
20 protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25 sequence of SEQ ID NO:46;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30 (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:45.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:45 from nucleotide 1922 to nucleotide 2350; the nucleotide sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350; the nucleotide sequence of the full-length protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886. In other preferred  
5 NO:45 from nucleotide 1922 to nucleotide 2350; the nucleotide sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350; the nucleotide sequence of the full-length protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886. In other preferred  
10 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886.

In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably  
15 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:46, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 66 to amino acid 75 of SEQ ID NO:46.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:45.  
20 ID NO:45.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
30 vc47\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);



and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

(bb) the nucleotide sequence of the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:45 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45 from nucleotide 1922 to nucleotide 2350, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:45 from nucleotide 1922 to nucleotide 2350, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:45 from nucleotide 1922 to nucleotide 2350. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:46;

(b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;

- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:46. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:46, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 66 to amino acid 75 of SEQ ID NO:46.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 111 to nucleotide 1337;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:47 from nucleotide 246 to nucleotide 1337;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:48;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:47.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:47 from nucleotide 111 to nucleotide 1337; the nucleotide sequence of SEQ ID NO:47 from nucleotide 246 to nucleotide 1337; the nucleotide sequence of the full-length protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 199 to amino acid 208 of SEQ ID NO:48.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:47.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 5 (aa) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and
- (ab) the nucleotide sequence of the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 10 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 15 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 20 vc54\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:47 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47. Also preferably the
- 30 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 111 to nucleotide 1337, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 111 to nucleotide 1337, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide

111 to nucleotide 1337. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 246 to nucleotide 1337, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from  
5 nucleotide 246 to nucleotide 1337, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 246 to nucleotide 1337.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:48;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:48. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:48, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 199 to amino acid 208 of SEQ ID NO:48.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:49 from nucleotide 270 to nucleotide 1637;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:49.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637; the nucleotide sequence of SEQ ID NO:49  
25 from nucleotide 270 to nucleotide 1637; the nucleotide sequence of the full-length protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
30 of clone vc57\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:50, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence from amino acid 236 to amino acid 245 of SEQ ID NO:50.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:49.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:49 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637. Also preferably the polynucleotide isolated according to the above  
10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 270 to nucleotide 1637, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:49 from nucleotide 270 to nucleotide 1637, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide 270 to nucleotide 1637.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
- (b) a fragment of the amino acid sequence of SEQ ID NO:50, the  
20 fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:50. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:50, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence  
30 from amino acid 236 to amino acid 245 of SEQ ID NO:50.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51;



- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:51.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934; the nucleotide sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934; the nucleotide sequence of the full-length protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone

ve13\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886.

In further preferred embodiments, the present invention provides a polynucleotide  
5 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of  
10 SEQ ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 315 to amino acid 324 of SEQ ID NO:52.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:51.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 20 (aa) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and
- (ab) the nucleotide sequence of the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and

- (bb) the nucleotide sequence of the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:51 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 25 (a) the amino acid sequence of SEQ ID NO:52;
- (b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- 30

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:52. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 315 to amino acid 324 of SEQ ID NO:52.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:53 from nucleotide 240 to nucleotide 503;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ve16\_1 deposited with the ATCC under  
15 accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
20 protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25 sequence of SEQ ID NO:54;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30 (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:53.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:53 from nucleotide 240 to nucleotide 503; the nucleotide sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503; the nucleotide sequence of the full-length protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:54.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:53.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and
    - (ab) the nucleotide sequence of the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and

(bb) the nucleotide sequence of the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:53 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 240 to nucleotide 503, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 240 to nucleotide 503, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 240 to nucleotide 503. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

- (b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:54. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:54, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:54.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55 from nucleotide 11 to nucleotide 1063;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:55 from nucleotide 71 to nucleotide 1063;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:56;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:55.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:55 from nucleotide 11 to nucleotide 1063; the nucleotide sequence of SEQ ID NO:55 from nucleotide 71 to nucleotide 1063; the nucleotide sequence of the full-length protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 170 to amino acid 179 of SEQ ID NO:56.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:55.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:



- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and
  - (ab) the nucleotide sequence of the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:55 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 11 to nucleotide 1063, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from nucleotide 11 to nucleotide 1063, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide

11 to nucleotide 1063. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 71 to nucleotide 1063, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from  
5 nucleotide 71 to nucleotide 1063, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 71 to nucleotide 1063.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:56;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:56. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:56, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 170 to amino acid 179 of SEQ ID NO:56.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:57 from nucleotide 755 to nucleotide 886;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:57.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886; the nucleotide sequence of SEQ ID NO:57  
25 from nucleotide 755 to nucleotide 886; the nucleotide sequence of the full-length protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
30 of clone vj2\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:58, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:58.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:57.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:57 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886. Also preferably the polynucleotide isolated according to the above  
10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57 from nucleotide 755 to nucleotide 886, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:57 from nucleotide 755 to nucleotide 886, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:57 from nucleotide 755 to nucleotide 886.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:58;
- (b) a fragment of the amino acid sequence of SEQ ID NO:58, the  
20 fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:58. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:58, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence  
30 from amino acid 52 to amino acid 61 of SEQ ID NO:58.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:59.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344; the nucleotide sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344; the nucleotide sequence of the full-length protein coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vp7\_1 deposited

with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:60.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:59.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and

- (bb) the nucleotide sequence of the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:59 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 25 (a) the amino acid sequence of SEQ ID NO:60;
- (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- 30

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:60. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably



comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:60.

5           In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61;
- 10           (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp8\_1 deposited with the ATCC under  
15           accession number 98886;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
20           protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25           sequence of SEQ ID NO:62;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30           (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:61.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757; the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757; the nucleotide sequence of the full-length protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886; or the nucleotide sequence of a mature protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:62, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:62.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:61.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

(bb) the nucleotide sequence of the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:61 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:62;

- (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:62. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:62, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:62.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 1048 to nucleotide 3726;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb22\_1 deposited with the ATCC under
- 20 accession number 98933;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- 25 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb22\_1 deposited with the ATCC under accession number 98933;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- 30 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

5 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:63.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:63 from nucleotide 1048 to nucleotide 3726; the nucleotide sequence of the full-length protein coding sequence of clone vb22\_1 deposited with the ATCC under accession number 98933; or the nucleotide sequence of a mature protein coding sequence of clone vb22\_1 deposited with the ATCC under accession number 98933. In other preferred  
15 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933.

In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably  
20 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:64, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 441 to amino acid 450 of SEQ ID NO:64.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
25 ID NO:63.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63; and

- (ab) the nucleotide sequence of the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 vb22\_1 deposited with the ATCC under accession number 98933;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:63 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63 from nucleotide 1048 to nucleotide 3726, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from nucleotide 1048 to nucleotide 3726, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide
- 30 1048 to nucleotide 3726.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:64;

- (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:64. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:64, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 441 to amino acid 450 of SEQ ID NO:64.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 134 to nucleotide 667;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:65 from nucleotide 191 to nucleotide 667;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA
- 30 insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:66;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- 5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:65.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:65 from nucleotide 134 to nucleotide 667; the nucleotide sequence of SEQ ID NO:65 from nucleotide 191 to nucleotide 667; the nucleotide sequence of the full-length protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933; or the nucleotide sequence of a mature protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:66, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 84 to amino acid 93 of SEQ ID NO:66.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:65.

30 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:



- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 5 (aa) SEQ ID NO:65, but excluding the poly(A) tail at the 3' end of SEQ ID NO:65; and
- (ab) the nucleotide sequence of the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 10 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 15 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:65, but excluding the poly(A) tail at the 3' end of SEQ ID NO:65; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 20 vc48\_1 deposited with the ATCC under accession number 98933;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:65 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:65, but excluding the poly(A) tail at the 3' end of SEQ ID NO:65. Also preferably the
- 30 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 134 to nucleotide 667, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from nucleotide 134 to nucleotide 667, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide

134 to nucleotide 667. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 191 to nucleotide 667, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from  
5 nucleotide 191 to nucleotide 667, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide 191 to nucleotide 667.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:66;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:66. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:66, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 84 to amino acid 93 of SEQ ID NO:66.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:67 from nucleotide 158 to nucleotide 457;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:67.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457; the nucleotide sequence of SEQ ID NO:67 from nucleotide 158 to nucleotide 457; the nucleotide sequence of the full-length protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933; or the nucleotide sequence of a mature protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising the amino acid sequence from amino acid 60 to amino acid 69 of SEQ ID NO:68.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:67.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and
    - (bb) the nucleotide sequence of the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:67 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457. Also preferably the polynucleotide isolated according to the above  
10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 158 to nucleotide 457, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from nucleotide 158 to nucleotide 457, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 158 to nucleotide 457.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
- (b) a fragment of the amino acid sequence of SEQ ID NO:68, the  
20 fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:68. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising the amino acid sequence  
30 from amino acid 60 to amino acid 69 of SEQ ID NO:68.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 10 207012;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA 15 insert of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:70;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment 20 comprising eight contiguous amino acids of SEQ ID NO:70;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:69.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387; the nucleotide sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387; the nucleotide sequence of the full-length protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012; or the nucleotide sequence of a mature protein coding sequence of clone vc61\_1

deposited with the ATCC under accession number 207012. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising the amino acid sequence from amino acid 221 to amino acid 230 of SEQ ID NO:70.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:69.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and

- (bb) the nucleotide sequence of the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:69 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:70;
- (b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:70. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably



comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising the amino acid sequence from amino acid 221 to amino acid 230 of SEQ ID NO:70.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:71 from nucleotide 44 to nucleotide 1513;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458;
- (e) a polynucleotide comprising the nucleotide sequence of the full-  
15 length protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number  
20 207012;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA  
25 insert of clone vp15\_1 deposited with the ATCC under accession number 207012;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:72;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment  
30 comprising eight contiguous amino acids of SEQ ID NO:72;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

(n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:71.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:71 from nucleotide 44 to nucleotide 1513; the nucleotide sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513; the nucleotide sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458; the nucleotide sequence of the full-length protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012; or the nucleotide sequence of a mature protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:72 from amino acid 1 to amino acid 139. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 240 to amino acid 249 of SEQ ID NO:72.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:71.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and

- (ab) the nucleotide sequence of the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 vp15\_1 deposited with the ATCC under accession number 207012;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:71 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 44 to nucleotide 1513, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from nucleotide 44 to nucleotide 1513, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:71 from nucleotide
- 30 44 to nucleotide 1513. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513, to a nucleotide sequence corresponding to the 3' end of

said sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:72;
- (b) the amino acid sequence of SEQ ID NO:72 from amino acid 1 to amino acid 139;
- (c) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and
- (d) the amino acid sequence encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:72 or the amino acid sequence of SEQ ID NO:72 from amino acid 1 to amino acid 139. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 240 to amino acid 249 of SEQ ID NO:72.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:73.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743; the nucleotide sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743; the nucleotide sequence of the full-length protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012; or the nucleotide sequence of a mature protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012. In further preferred embodiments, the present invention provides a

polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:74.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:73.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

- (ab) the nucleotide sequence of the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;

- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

- (bb) the nucleotide sequence of the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:73 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
- (b) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:74. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:74.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp19\_1 deposited with the ATCC under accession number 207012;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp19\_1 deposited with the ATCC under  
15 accession number 207012;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- 20 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 25 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:75.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461; the nucleotide sequence of the full-length protein coding sequence of clone vp19\_1 deposited with the ATCC under accession



number 207012; or the nucleotide sequence of a mature protein coding sequence of clone vp19\_1 deposited with the ATCC under accession number 207012. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number  
5 207012. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence  
10 of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:76.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:75.

Further embodiments of the invention provide isolated polynucleotides produced  
15 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (aa) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and

(ab) the nucleotide sequence of the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;

25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and

- (bb) the nucleotide sequence of the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:75 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
- (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:76. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:76.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide 368;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 141 to nucleotide 368;
- 10 (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 51 to nucleotide 332;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012;
- 15 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012;
- 20 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- 25 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- 30 (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

(n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:77.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide 368; the nucleotide sequence of SEQ ID NO:77 from nucleotide 141 to nucleotide 368; the nucleotide sequence of SEQ ID NO:77 from nucleotide 51 to nucleotide 332; the nucleotide sequence of the full-length protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012; or the nucleotide sequence of a mature protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 93. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:78.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:77.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and

(bb) the nucleotide sequence of the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77, and  
20 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:77 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide  
25 368, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide 368, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide 368. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
30 NO:77 from nucleotide 141 to nucleotide 368, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 141 to nucleotide 368, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide 141 to nucleotide 368. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide

sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 51 to nucleotide 332, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 51 to nucleotide 332, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from  
5 nucleotide 51 to nucleotide 332.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
- 10 (b) the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 93;
- (c) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
- (d) the amino acid sequence encoded by the cDNA insert of clone  
15 vq1\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:78 or the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 93. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid  
20 sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:78.

25 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:79 from nucleotide 2 to nucleotide 1018;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 20 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least
- 25 25% of the length of SEQ ID NO:79.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:79 from nucleotide 2 to nucleotide 1018; the nucleotide sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018; the nucleotide sequence of the full-length protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number

30 207011; or the nucleotide sequence of a mature protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011. In further preferred embodiments, the present invention provides a

polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 164 to amino acid 173 of SEQ ID NO:80.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:79.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79; and

- (ab) the nucleotide sequence of the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79; and

- (bb) the nucleotide sequence of the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and



(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:79 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 2 to nucleotide 1018, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 2 to nucleotide 1018, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 2 to nucleotide 1018. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:80;
- (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:80. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 164 to amino acid 173 of SEQ ID NO:80.

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial, yeast, insect and mammalian cells, transformed with such polynucleotide compositions. Also provided by the present invention are organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein.

Processes are also provided for producing a protein, which comprise:

- (a) growing a culture of the host cell transformed with such polynucleotide compositions in a suitable culture medium; and
- (b) purifying the protein from the culture.

The protein produced according to such methods is also provided by the present invention.

Protein compositions of the present invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody which specifically reacts with such protein are also provided by the present invention.

Methods are also provided for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition comprising a protein of the present invention and a pharmaceutically acceptable carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B are schematic representations of the pED6 and pNOTs vectors, respectively, used for deposit of clones disclosed herein.

#### DETAILED DESCRIPTION

##### ISOLATED PROTEINS AND POLYNUCLEOTIDES

Nucleotide and amino acid sequences, as presently determined, are reported below for each clone and protein disclosed in the present application. The nucleotide sequence of each clone can readily be determined by sequencing of the deposited clone in accordance with known methods. The predicted amino acid sequence (both full-length and mature forms) can then be determined from such nucleotide sequence. The amino acid sequence of the protein encoded by a particular clone can also be determined by expression of the clone in a suitable host cell, collecting the protein and determining its sequence. For each disclosed protein applicants have identified what they have

determined to be the reading frame best identifiable with sequence information available at the time of filing.

As used herein a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum.

10        Clone "vb11\_1"

A polynucleotide of the present invention has been identified as clone "vb11\_1". vb11\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb11\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb11\_1 protein").

The nucleotide sequence of vb11\_1 as presently determined is reported in SEQ ID NO:1, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb11\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:2. Another potential vb11\_1 reading frame and predicted amino acid sequence that could be encoded by basepairs 84 to 236 of SEQ ID NO:1 is reported in SEQ ID NO:121. Amino acids 13 to 25 of SEQ ID NO:121 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26 of SEQ ID NO:121. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:121.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb11\_1 should be approximately 1751 bp.

The nucleotide sequence disclosed herein for vb11\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb11\_1 demonstrated at least some similarity with sequences identified as N94870 (yy63b05.r1 Homo sapiens cDNA clone 278193 5'). Based upon sequence similarity, vb11\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane

domain within the vb11\_1 protein sequence centered around amino acid 27 of SEQ ID NO:2.

Clone "vb12\_1"

5 A polynucleotide of the present invention has been identified as clone "vb12\_1". vb12\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb12\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb12\_1 protein").

10 The nucleotide sequence of vb12\_1 as presently determined is reported in SEQ ID NO:3, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb12\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:4. Amino acids 34 to 46 of SEQ ID NO:4 are a predicted leader/signal sequence, with the predicted  
15 mature amino acid sequence beginning at amino acid 47. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb12\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 vb12\_1 should be approximately 2289 bp.

The nucleotide sequence disclosed herein for vb12\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb12\_1 demonstrated at least some similarity with sequences identified as AA426009 (zw49e11.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA  
25 clone 773420 3', mRNA sequence). Based upon sequence similarity, vb12\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the vb12\_1 protein sequence, centered around amino acids 11, 60, and 104 of SEQ ID NO:4, respectively. The nucleotide sequence of vb12\_1 indicates that it may contain a THE1B  
30 repeat sequence.

vb12\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 17 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "vb14\_1"

A polynucleotide of the present invention has been identified as clone "vb14\_1". vb14\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb14\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb14\_1 protein").

The nucleotide sequence of vb14\_1 as presently determined is reported in SEQ ID NO:5, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb14\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:6. Amino acids 79 to 91 of SEQ ID NO:6 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 92. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb14\_1 protein. Another potential vb14\_1 reading frame and predicted amino acid sequence that could be encoded by basepairs 182 to 484 of SEQ ID NO:5 is reported in SEQ ID NO:122.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb14\_1 should be approximately 2377 bp.

The nucleotide sequence disclosed herein for vb14\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb14\_1 demonstrated at least some similarity with sequences identified as AF007149 (Homo sapiens clone 23568, 23621, 23795, 23873 and 23874 mRNA sequences), AF070612 (Homo sapiens clone 24771 mRNA sequence), T23635 (Human gene signature HUMGS05495; standard; cDNA to mRNA), and W02197 (za57e04.r1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone 296670 5', mRNA sequence). Based upon sequence similarity, vb14\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "ve11\_1"

A polynucleotide of the present invention has been identified as clone "ve11\_1". ve11\_1 was isolated from a human adult brain (Alzheimer's hippocampus level 7) cDNA library and was identified as encoding a secreted or transmembrane protein on the basis

of computer analysis of the amino acid sequence of the encoded protein. ve11\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ve11\_1 protein").

The nucleotide sequence of ve11\_1 as presently determined is reported in SEQ ID NO:7, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ve11\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:8. Amino acids 1 to 9 of SEQ ID NO:8 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 10. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ve11\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ve11\_1 should be approximately 984 bp.

The nucleotide sequence disclosed herein for ve11\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ve11\_1 demonstrated at least some similarity with sequences identified as F22745 (H.sapiens EST sequence (LL45/C09) from skeletal muscle, mRNA sequence) and Q60824 (Human brain Expressed Sequence Tag EST00928; standard; DNA). Based upon sequence similarity, ve11\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the ve11\_1 protein sequence centered around amino acid 35 of SEQ ID NO:8.

#### Clone "vf2\_1"

A polynucleotide of the present invention has been identified as clone "vf2\_1". vf2\_1 was isolated from a human adult heart cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vf2\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vf2\_1 protein").

The nucleotide sequence of vf2\_1 as presently determined is reported in SEQ ID NO:9, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vf2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:10. Amino acids 20 to 32

of SEQ ID NO:10 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 33. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vf2\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vf2\_1 should be approximately 1162 bp.

The nucleotide sequence disclosed herein for vf2\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vf2\_1 demonstrated at least some similarity with sequences identified as AA605037 (no68h10.s1 NCI\_CGAP\_AA1 Homo sapiens cDNA clone IMAGE:1112035 similar to contains Alu repetitive element;contains element THR repetitive element; mRNA sequence). Based upon sequence similarity, vf2\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the vf2\_1 protein sequence, one centered around amino acid 30 and another around amino acid 70 of SEQ ID NO:10. The nucleotide sequence of vf2\_1 indicates that it may contain an Alu repetitive element.

#### Clone "vg2\_1"

A polynucleotide of the present invention has been identified as clone "vg2\_1". vg2\_1 was isolated from a human adult brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vg2\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vg2\_1 protein").

The nucleotide sequence of vg2\_1 as presently determined is reported in SEQ ID NO:11, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vg2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:12. Amino acids 34 to 46 of SEQ ID NO:12 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 47. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should

the predicted leader/signal sequence not be separated from the remainder of the vg2\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vg2\_1 should be approximately 1993 bp.

- 5       The nucleotide sequence disclosed herein for vg2\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vg2\_1 demonstrated at least some similarity with sequences identified as AA830272 (oc45g11.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 1352708 3' similar to TR Q92853 Q92853 HU-K4; mRNA sequence) and D31740
- 10 (Homo sapiens DNA, CpG island). The predicted amino acid sequence disclosed herein for vg2\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vg2\_1 protein demonstrated at least some similarity to sequences identified as AF026124 (schwannoma-associated protein [Mus musculus]) and U60644 (HU-K4 [Homo sapiens]). Based upon sequence similarity,
- 15 vg2\_1 proteins and each similar protein or peptide may share at least some activity. Profile hidden markov model analysis (Eddy, S. R., 1996, *Curr. Opin. Struct. Biol.* 6(3): 361-365; incorporated by reference herein) of the predicted vg2\_1 protein revealed two phospholipase D active sites (amino acid residues 209 to 236 and 423 to 449 of SEQ ID NO:12). Phospholipase D (PLD) genes are members of a superfamily that is defined by
- 20 several highly conserved motifs. In mammals, it has been proposed that phospholipase D plays a role in membrane vesicular trafficking and in signal transduction. Using site-directed mutagenesis, twenty-five point mutants have been made in human PLD1 (hPLD1) and then characterized (Sung *et al.*, 1997, *EMBO J.* 16(15): 4519-4530; which is incorporated by reference herein). Sung *et al.* found that a motif (HxKxxxxD; see for
- 25 example amino acids 214-221 of SEQ ID NO:12) and a serine/threonine conserved in all members of the PLD superfamily are critical for PLD biochemical activity, suggesting a possible catalytic mechanism. The vg2\_1 clone appears to encode a membrane protein that may be a phospholipase related to the phospholipase D family. The TopPredII computer program predicts four potential transmembrane domains within the vg2\_1
- 30 protein sequence, centered around amino acids 40, 305, 330, and 455 of SEQ ID NO:12, respectively.



Clone "vj1\_1"

A polynucleotide of the present invention has been identified as clone "vj1\_1". vj1\_1 was isolated from a human fetal brain cDNA library (enriched for G-protein-coupled receptors) and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vj1\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vj1\_1 protein").

The nucleotide sequence of vj1\_1 as presently determined is reported in SEQ ID NO:13. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vj1\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:14. Amino acids 1 to 12 of SEQ ID NO:14 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 13. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vj1\_1 protein. Another potential vj1\_1 reading frame and predicted amino acid sequence that could be encoded by basepairs 1795 to 2064 of SEQ ID NO:13 is reported in SEQ ID NO:123.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vj1\_1 should be approximately 2895 bp.

The nucleotide sequence disclosed herein for vj1\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vj1\_1 demonstrated at least some similarity with sequences identified as AA410352 (zv11f01.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 753337 5', mRNA sequence). Based upon sequence similarity, vj1\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vj1\_1 protein sequence centered around amino acid 70 of SEQ ID NO:14. The nucleotide sequence of vj1\_1 indicates that it may contain repetitive elements.

Clone "vl1\_1"

A polynucleotide of the present invention has been identified as clone "vl1\_1". vl1\_1 was isolated from a human fetal cartilage cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the

amino acid sequence of the encoded protein. vl1\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vl1\_1 protein").

The nucleotide sequence of vl1\_1 as presently determined is reported in SEQ ID NO:15, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the vl1\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:16. Amino acids 187 to 199 of SEQ ID NO:16 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 200. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
10 the predicted leader/signal sequence not be separated from the remainder of the vl1\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vl1\_1 should be approximately 1936 bp.

The nucleotide sequence disclosed herein for vl1\_1 was searched against the  
15 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vl1\_1 demonstrated at least some similarity with sequences identified as AA464362 (zx81b12.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 810143 5', mRNA sequence), M90089 (Mouse inositol 1,4,5-triphosphate receptor mRNA sequence), and T21689 (Human gene signature HUMGS03131; standard; cDNA  
20 to mRNA). The predicted amino acid sequence disclosed herein for vl1\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vl1\_1 protein demonstrated at least some similarity to the sequence identified as U80846 (partial CDS [Caenorhabditis elegans]). Based upon sequence similarity, vl1\_1 proteins and each similar protein or peptide may share at least  
25 some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the vl1\_1 protein sequence, one centered around amino acid 192 and another around amino acid 234 of SEQ ID NO:16.

vl1\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 37 kDa was detected in membrane fractions using SDS  
30 polyacrylamide gel electrophoresis.

Clone "vk2\_1"

A polynucleotide of the present invention has been identified as clone "vk2\_1". vk2\_1 was isolated from a human adult brain cDNA library (enriched for G-protein-coupled receptors) and was identified as encoding a secreted or transmembrane protein  
5 on the basis of computer analysis of the amino acid sequence of the encoded protein. vk2\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vk2\_1 protein").

The nucleotide sequence of vk2\_1 as presently determined is reported in SEQ ID NO:17, and includes a poly(A) tail. What applicants presently believe to be the proper  
10 reading frame and the predicted amino acid sequence of the vk2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:18. Amino acids 10 to 22 of SEQ ID NO:18 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
15 the predicted leader/signal sequence not be separated from the remainder of the vk2\_1 protein. Basepairs 416 to 418 of SEQ ID NO:17 may represent the site of an alternatively spliced exon that is not present in clone vk2\_1.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vk2\_1 should be approximately 1284 bp.

20 The nucleotide sequence disclosed herein for vk2\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vk2\_1 demonstrated at least some similarity with sequences identified as AA152101 (zl49f09.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 505289 3', mRNA sequence) and Q78696 (Sequence encoding therapeutic  
25 polypeptide from glioblastoma cell line; standard; cDNA to mRNA). The predicted amino acid sequence disclosed herein for vk2\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vk2\_1 protein demonstrated at least some similarity to the sequence identified as R66278 (Therapeutic polypeptide from glioblastoma cell line). Based upon sequence similarity,  
30 vk2\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the vk2\_1 protein sequence, one centered around amino acid 61 and another around amino acid 97 of SEQ ID NO:18.

Clone "vb21\_1"

A polynucleotide of the present invention has been identified as clone "vb21\_1". vb21\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb21\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb21\_1 protein").

The nucleotide sequence of vb21\_1 as presently determined is reported in SEQ ID NO:19, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb21\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:20. Amino acids 296 to 308 of SEQ ID NO:20 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 309. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb21\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb21\_1 should be approximately 4159 bp.

The nucleotide sequence disclosed herein for vb21\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb21\_1 demonstrated at least some similarity with sequences identified as AA026150 (zj99c10.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 469170 3', mRNA sequence), T72108 (Human semaphorin Z gene; standard; cDNA to mRNA), U52840 (Human semaphorin F homolog), X97817 (M. musculus mRNA for semaphorin F), and X97818 (M. musculus mRNA for semaphorin G). The predicted amino acid sequence disclosed herein for vb21\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vb21\_1 protein demonstrated at least some similarity to sequences identified as W19857 (Human semaphorin Z) and X97818 (semaphorin G [Mus musculus]). Semaphorins are important membrane proteins involved in axonal guidance in the embryonic stage, and may also have a role in nerve regeneration after injury. Based upon sequence similarity, vb21\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four additional potential transmembrane domains within the vb21\_1 protein sequence, centered around amino acids 237, 523, 769, and 895 of SEQ ID NO:20, respectively.

Clone "vc35\_1"

A polynucleotide of the present invention has been identified as clone "vc35\_1". vc35\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc35\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc35\_1 protein").

The nucleotide sequence of vc35\_1 as presently determined is reported in SEQ ID NO:21, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc35\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:22. Amino acids 38 to 50 of SEQ ID NO:22 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 51. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc35\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc35\_1 should be approximately 3042 bp.

The nucleotide sequence disclosed herein for vc35\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc35\_1 demonstrated at least some similarity with sequences identified as AA532364 (nj12a08.s1 NCI\_CGAP\_Pr22 Homo sapiens cDNA clone IMAGE:986102, mRNA sequence), AF029343 (human protocadherin 68), and T22263 (Human gene signature HUMGS03835; standard; cDNA to mRNA). The predicted amino acid sequence disclosed herein for vc35\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc35\_1 protein demonstrated at least some similarity to sequences identified as Y08715 (protocadherin-4 [Mus musculus]). Based upon sequence similarity, vc35\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vc36\_1"

A polynucleotide of the present invention has been identified as clone "vc36\_1". vc36\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the

amino acid sequence of the encoded protein. vc36\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc36\_1 protein").

The nucleotide sequence of vc36\_1 as presently determined is reported in SEQ ID NO:23, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the vc36\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:24. Amino acids 24 to 36 of SEQ ID NO:24 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 37. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
10 the predicted leader/signal sequence not be separated from the remainder of the vc36\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc36\_1 should be approximately 1395 bp.

The nucleotide sequence disclosed herein for vc36\_1 was searched against the  
15 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc36\_1 demonstrated at least some similarity with sequences identified as AA259070 (zs33c04.r1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 686982 5', mRNA sequence) and W67508 (zd40f11.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 343149 3', mRNA sequence). Based upon sequence similarity, vc36\_1  
20 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vc36\_1 indicates that it may contain repetitive elements.

#### Clone "vc38\_1"

A polynucleotide of the present invention has been identified as clone "vc38\_1".  
25 vc38\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc38\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc38\_1 protein").

The nucleotide sequence of vc38\_1 as presently determined is reported in SEQ ID  
30 NO:25, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc38\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:26.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc38\_1 should be approximately 2468 bp.

The nucleotide sequence disclosed herein for vc38\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc38\_1 demonstrated at least some similarity with sequences identified as AF037400 (neuropeptide Y/peptide YY receptor Ya [Danio rerio]). Motifs analysis and profile hidden markov model analysis of the predicted vc38\_1 protein both reveal the presence of the G-protein-coupled receptor signature. G-protein-coupled receptors (also called R7G) are an extensive group of hormones, neurotransmitters, odorants, and light receptors which transduce extracellular signals by interaction with guanine nucleotide-binding (G) proteins. Most G-protein-coupled receptors lack a signal peptide, as does the predicted vc38\_1 protein. Based upon sequence similarity, vc38\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts seven potential transmembrane domains within the vc38\_1 protein sequence, centered around amino acids 60, 90, 130, 170, 225, 280, and 318 of SEQ ID NO:26, respectively.

vc38\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 71 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

#### Clone "vc39\_1"

A polynucleotide of the present invention has been identified as clone "vc39\_1". vc39\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc39\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc39\_1 protein").

The nucleotide sequence of vc39\_1 as presently determined is reported in SEQ ID NO:27, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc39\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:28. Amino acids 2 to 14 of SEQ ID NO:28 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc39\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc39\_1 should be approximately 2048 bp.

The nucleotide sequence disclosed herein for vc39\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc39\_1 demonstrated at least some similarity with sequences identified as AA631722 (np79d04.s1 NCI\_CGAP\_Pr2 Homo sapiens cDNA clone IMAGE:1132519 similar to gb:M21121 T-CELL SPECIFIC RANTES PROTEIN PRECURSOR (HUMAN); contains Alu repetitive element; mRNA sequence). Based upon sequence similarity, vc39\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the vc39\_1 protein sequence centered around amino acid 40 of SEQ ID NO:28. The nucleotide sequence of vc39\_1 indicates that it may contain an Alu/SVA repetitive element.

15        Clone "vc40\_1"

A polynucleotide of the present invention has been identified as clone "vc40\_1". vc40\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc40\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc40\_1 protein").

The nucleotide sequence of vc40\_1 as presently determined is reported in SEQ ID NO:29, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc40\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:30. Amino acids 19 to 31 of SEQ ID NO:30 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 32. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc40\_1 protein.

30        The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc40\_1 should be approximately 2297 bp.

The nucleotide sequence disclosed herein for vc40\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc40\_1 demonstrated at least some similarity with sequences



identified as AA143014 (zl48g04.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 505206 5', mRNA sequence) and T20006 (Human gene signature HUMGS01143; standard; cDNA to mRNA). Based upon sequence similarity, vc40\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the vc40\_1 protein sequence, centered around amino acids 101, 136, and 182 of SEQ ID NO:30, respectively.

#### Clone "vc46\_1"

A polynucleotide of the present invention has been identified as clone "vc46\_1". vc46\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc46\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc46\_1 protein").

The nucleotide sequence of vc46\_1 as presently determined is reported in SEQ ID NO:31, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc46\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:32. Amino acids 10 to 22 of SEQ ID NO:32 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc46\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc46\_1 should be approximately 2938 bp.

The nucleotide sequence disclosed herein for vc46\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc46\_1 demonstrated at least some similarity with sequences identified as AA029404 (ze94e06.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 366658 5', mRNA sequence) and AQ071029 (human genomic fragment). Based upon sequence similarity, vc46\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the vc46\_1 protein sequence, one centered around amino acid 70 and another around amino acid 130 of SEQ ID NO:32.

vc46\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 19 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

5        Clone "vc49\_1"

A polynucleotide of the present invention has been identified as clone "vc49\_1". vc49\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc49\_1 is a full-length clone, including the  
10    entire coding sequence of a secreted protein (also referred to herein as "vc49\_1 protein").

The nucleotide sequence of vc49\_1 as presently determined is reported in SEQ ID NO:33, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc49\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:34. Amino acids 14 to 26  
15    of SEQ ID NO:34 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc49\_1 protein.

20        The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc49\_1 should be approximately 3471 bp.

The nucleotide sequence disclosed herein for vc49\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc49\_1 demonstrated at least some similarity with sequences  
25    identified as AI075929 (ov46h11.x1 Soares\_testis\_NHT Homo sapiens cDNA clone IMAGE 1640421 3' similar to TR Q63418 Q63418 PROTOCADHERIN-3; mRNA sequence), I79964 (Sequence 109 from patent US 5708143), and T03572 (Human protocadherin pc3 coding sequence; standard; cDNA). The predicted amino acid sequence disclosed herein for vc49\_1 was searched against the GenPept and GeneSeq amino acid sequence databases  
30    using the BLASTX search protocol. The predicted vc49\_1 protein demonstrated at least some similarity to sequences identified as L43592 (protocadherin-3 [Rattus norvegicus]) and R86865 (Human protocadherin pc3). Based upon sequence similarity, vc49\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the vc49\_1

protein sequence, one definite transmembrane domain centered around amino acid 700 and another possible transmembrane domain centered around amino acid 260 of SEQ ID NO:34. Profile hidden markov model and motifs analyses of the predicted vc49\_1 protein sequence have revealed it to contain five cadherin extracellular repeated domain signatures at amino acids 142 to 242, 251 to 347, 356 to 451, 460 to 561, and 576 to 671 of SEQ ID NO:34. Cadherins are a family of animal glyco-proteins responsible for calcium-dependent cell-cell adhesion. Cadherins preferentially interact with themselves in a homophilic manner in connecting cells; thus acting as both receptor and ligand. Structurally, cadherins are built of the following domains: a signal sequence, followed by a propeptide of about 130 residues, then an extracellular domain of around 600 residues, then a transmembrane region, and finally a C-terminal cytoplasmic domain of about 150 residues. The predicted vc49\_1 protein sequence almost exactly follows this structure (its cytoplasmic domain being approximately 100 amino acids). Clearly, vc49\_1 protein appears to represent a novel member of the cadherin superfamily.

#### Clone "vc50\_1"

A polynucleotide of the present invention has been identified as clone "vc50\_1". vc50\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc50\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc50\_1 protein").

The nucleotide sequence of vc50\_1 as presently determined is reported in SEQ ID NO:35, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc50\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:36. Amino acids 20 to 32 of SEQ ID NO:36 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 33. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc50\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc50\_1 should be approximately 3819 bp.

The nucleotide sequence disclosed herein for vc50\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and

FASTA search protocols. vc50\_1 demonstrated at least some similarity with sequences identified as AA193122 (zr39d05.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 665769 5', mRNA sequence), T26031 (Human gene signature HUMGS08267; standard; cDNA to mRNA), Z31718 (H.sapiens gene for myelin protein zero), and Z99943 (Human DNA sequence from PAC 313L4 on chromosome 1q24). The predicted amino acid sequence disclosed herein for vc50\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc50\_1 protein demonstrated at least some similarity to the sequence identified as K03242 (rat P0 myelin prepeptide), L24893 (myelin protein zero [Homo sapiens]), and M62860 (mouse peripheral myelin protein). Based upon sequence similarity, vc50\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the vc50\_1 protein sequence centered around amino acid 181 of SEQ ID NO:36.

vc50\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 26 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "vc51\_1"

A polynucleotide of the present invention has been identified as clone "vc51\_1". vc51\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc51\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc51\_1 protein").

The nucleotide sequence of vc51\_1 as presently determined is reported in SEQ ID NO:37, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc51\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:38. Amino acids 12 to 24 of SEQ ID NO:38 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 25. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc51\_1 protein. If the "G" residue at position 388 of SEQ ID NO:37 were deleted, two alternative potential vc51\_1 reading frames and predicted amino acid sequences that could be

encoded by basepairs 333 to 1310 of SEQ ID NO:37 and by basepairs 139 to 522 of SEQ ID NO:37 are reported in SEQ ID NO:124 and SEQ ID NO:125, respectively.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc51\_1 should be approximately 1992 bp.

5       The nucleotide sequence disclosed herein for vc51\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc51\_1 demonstrated at least some similarity with sequences identified as T21514 (Human gene signature HUMGS02887; standard; cDNA to mRNA) and W52782 (zd13h06.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 340571  
10   5', mRNA sequence). The predicted amino acid sequence disclosed herein for vc51\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc51\_1 protein demonstrated at least some similarity to sequences identified as U90716 (human cell surface protein HCAR), Y07593 (coxsackie and adenovirus receptor protein [Homo sapiens]), Y10320 (mouse coxsackie  
15   and adenovirus receptor homolog), and W14146 (Human A33 antigen). Based upon sequence similarity, vc51\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four additional potential transmembrane domains within the vc51\_1 protein sequence centered around amino acids 17, 216, 260, and 373 of SEQ ID NO:38, respectively.

20

#### Clone "vc52\_1"

A polynucleotide of the present invention has been identified as clone "vc52\_1". vc52\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the  
25   amino acid sequence of the encoded protein. vc52\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc52\_1 protein").

The nucleotide sequence of vc52\_1 as presently determined is reported in SEQ ID NO:39, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc52\_1 protein corresponding  
30   to the foregoing nucleotide sequence is reported in SEQ ID NO:40. Amino acids 19 to 31 of SEQ ID NO:40 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 32. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should

the predicted leader/signal sequence not be separated from the remainder of the vc52\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc52\_1 should be approximately 2018 bp.

5        The nucleotide sequence disclosed herein for vc52\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc52\_1 demonstrated at least some similarity with sequences identified as AA075627 (zm89a01.s1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone 545064 3', mRNA sequence) and T24879 (Human gene signature  
10 HUMGS06985; standard; cDNA to mRNA). The predicted amino acid sequence disclosed herein for vc52\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc52\_1 protein demonstrated at least some similarity to sequences identified as AL021890 (putative protein [Arabidopsis thaliana]), L47993 (ORF YJR072c [Saccharomyces cerevisiae]), and U10402  
15 (undefined protein [Caenorhabditis elegans]). Based upon sequence similarity, vc52\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vc52\_1 protein sequence centered around amino acid 145 of SEQ ID NO:40.

vc52\_1 protein was expressed in a COS cell expression system, and an expressed  
20 protein band of approximately 44 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "vc33\_1"

A polynucleotide of the present invention has been identified as clone "vc33\_1".  
25 vc33\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc33\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc33\_1 protein").

The nucleotide sequence of vc33\_1 as presently determined is reported in SEQ ID  
30 NO:41, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc33\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:42. Amino acids 99 to 111 of SEQ ID NO:42 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 112. Due to the hydrophobic nature of the

predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc33\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
5 vc33\_1 should be approximately 2877 bp.

The nucleotide sequence disclosed herein for vc33\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc33\_1 demonstrated at least some similarity with sequences identified as AA846599 (aj97g02.s1 Soares parathyroid tumor NbHPA Homo sapiens  
10 cDNA clone IMAGE:1404434 3' similar to gb:M95549 SODIUM/GLUCOSE COTRANSPORTER-LIKE (HUMAN); mRNA sequence), M95549 (Homo sapiens sodium/glucose cotransporter-like protein mRNA, complete cds), and Q89779 (Cotransporter protein SNST1 cDNA; standard; cDNA). The predicted amino acid  
15 amino acid sequence disclosed herein for vc33\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc33\_1 protein demonstrated at least some similarity to sequences identified as M95549 (sodium/glucose cotransporter-like protein [Homo sapiens]) and R73593 (Cotransporter protein SNST1). Based upon sequence similarity, vc33\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts  
20 three additional potential transmembrane domains within the vc33\_1 protein sequence, centered around amino acids 186, 260, and 324 of SEQ ID NO:42, respectively.

vc33\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 45 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

25

#### Clone "vc34\_1"

A polynucleotide of the present invention has been identified as clone "vc34\_1". vc34\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the  
30 amino acid sequence of the encoded protein. vc34\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc34\_1 protein").

The nucleotide sequence of vc34\_1 as presently determined is reported in SEQ ID NO:43, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc34\_1 protein corresponding

to the foregoing nucleotide sequence is reported in SEQ ID NO:44. Amino acids 4 to 16 of SEQ ID NO:44 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
5 the predicted leader/signal sequence not be separated from the remainder of the vc34\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc34\_1 should be approximately 3062 bp.

The nucleotide sequence disclosed herein for vc34\_1 was searched against the  
10 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc34\_1 demonstrated at least some similarity with sequences identified as AA927558 (om71e04.s1 NCI\_CGAP\_GC4 Homo sapiens cDNA clone IMAGE 1552638 3', mRNA sequence) and U79281 (Human clone 23588 mRNA sequence). Based upon sequence similarity, vc34\_1 proteins and each similar protein or peptide may share  
15 at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the vc34\_1 protein sequence, one centered around amino acid 251 and another around amino acid 283 of SEQ ID NO:44.

vc34\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 72 kDa was detected in conditioned medium using SDS  
20 polyacrylamide gel electrophoresis.

#### Clone "vc47\_1"

A polynucleotide of the present invention has been identified as clone "vc47\_1". vc47\_1 was isolated from a human fetal brain cDNA library and was identified as  
25 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc47\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc47\_1 protein").

The nucleotide sequence of vc47\_1 as presently determined is reported in SEQ ID NO:45, and includes a poly(A) tail. What applicants presently believe to be the proper  
30 reading frame and the predicted amino acid sequence of the vc47\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:46. Amino acids 93 to 105 of SEQ ID NO:46 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 106. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should



the predicted leader/signal sequence not be separated from the remainder of the vc47\_1 protein.

Another potential vc47\_1 reading frame and predicted amino acid sequence that could be encoded by basepairs 1047 to 1322 of SEQ ID NO:45 is reported in SEQ ID NO:126. Amino acids 11 to 23 of SEQ ID NO:126 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of this predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:126.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc47\_1 should be approximately 3676 bp.

The nucleotide sequence disclosed herein for vc47\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc47\_1 demonstrated at least some similarity with sequences identified as AA339320 (EST44392 Fetal brain I Homo sapiens cDNA 5' end, mRNA sequence) and R02462 (ye82h04.r1 Homo sapiens cDNA clone 124279 5'). Based upon sequence similarity, vc47\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vc47\_1 indicates that it may contain one or more of the following repetitive elements: Alu, L1MB7.

#### Clone "vc54\_1"

A polynucleotide of the present invention has been identified as clone "vc54\_1". vc54\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc54\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc54\_1 protein").

The nucleotide sequence of vc54\_1 as presently determined is reported in SEQ ID NO:47, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc54\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:48. Amino acids 33 to 45 of SEQ ID NO:48 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 46. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should

the predicted leader/signal sequence not be separated from the remainder of the vc54\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc54\_1 should be approximately 2083 bp.

5       The nucleotide sequence disclosed herein for vc54\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc54\_1 demonstrated at least some similarity with sequences identified as AF007152 (Homo sapiens clone 23649 and 23755 unknown mRNA, partial cds), Q76901 (Human genome fragment (Preferred); standard; DNA), and T46905 (EST014  
10 BL29 Burkitt's lymphoma, Pascalis Sideras Homo sapiens cDNA clone BL29-14 5', mRNA sequence). The predicted amino acid sequence disclosed herein for vc54\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc54\_1 protein demonstrated at least some similarity to the sequence identified as AF007152 (unknown [Homo sapiens]). Based upon sequence  
15 similarity, vc54\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the vc54\_1 protein sequence, one centered around amino acid 220 and another around amino acid 247 of SEQ ID NO:48.

vc54\_1 protein was expressed in a COS cell expression system, and an expressed  
20 protein band of approximately 44 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "vc57\_1"

A polynucleotide of the present invention has been identified as clone "vc57\_1".  
25 vc57\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc57\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc57\_1 protein").

The nucleotide sequence of vc57\_1 as presently determined is reported in SEQ ID  
30 NO:49, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc57\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:50. Amino acids 15 to 27 of SEQ ID NO:50 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 28. Due to the hydrophobic nature of the

predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc57\_1 protein.

5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc57\_1 should be approximately 2564 bp.

10 The nucleotide sequence disclosed herein for vc57\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc57\_1 demonstrated at least some similarity with sequences identified as AA156231 (zl50a11.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 505340 3', mRNA sequence). The predicted amino acid sequence disclosed herein for vc57\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc57\_1 protein demonstrated at least some similarity to the sequence identified as U41635 (OS-9 precursor [Homo sapiens]). Based upon sequence similarity, vc57\_1 proteins and each similar protein or peptide may share at least some activity.

15 vc57\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 51 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

20 Clone "ve13\_1"

A polynucleotide of the present invention has been identified as clone "ve13\_1". ve13\_1 was isolated from a human adult brain (Alzheimer's hippocampus level 7) cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ve13\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ve13\_1 protein").

25 The nucleotide sequence of ve13\_1 as presently determined is reported in SEQ ID NO:51, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ve13\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:52. Amino acids 551 to 563 of SEQ ID NO:52 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 564. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should

the predicted leader/signal sequence not be separated from the remainder of the ve13\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ve13\_1 should be approximately 3046 bp.

5 The nucleotide sequence disclosed herein for ve13\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ve13\_1 demonstrated at least some similarity with sequences identified as AA587395 (nn82h06.s1 NCI\_CGAP\_Co9 Homo sapiens cDNA clone IMAGE:1090427 similar to contains element THR repetitive element; mRNA sequence)  
10 and Q76778 (Human genome fragment (Preferred); standard; DNA). The predicted amino acid sequence disclosed herein for ve13\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ve13\_1 protein demonstrated at least some similarity to the sequence identified as U50828 (sel-1 gene product [Caenorhabditis elegans]). Based upon sequence similarity, ve13\_1 proteins  
15 and each similar protein or peptide may share at least some activity.

#### Clone "ve16\_1"

A polynucleotide of the present invention has been identified as clone "ve16\_1". ve16\_1 was isolated from a human adult brain (Alzheimer's hippocampus level 7) cDNA  
20 library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ve16\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ve16\_1 protein").

The nucleotide sequence of ve16\_1 as presently determined is reported in SEQ ID  
25 NO:53, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ve16\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:54. Amino acids 14 to 26 of SEQ ID NO:54 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the  
30 predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ve16\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ve16\_1 should be approximately 2033 bp.

The nucleotide sequence disclosed herein for ve16\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the databases. The nucleotide sequence of ve16\_1 indicates that it may contain one or more of the following repetitive elements:

5 Alu, MER.

Clone "vf3\_1"

A polynucleotide of the present invention has been identified as clone "vf3\_1". vf3\_1 was isolated from a human adult heart cDNA library and was identified as  
10 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vf3\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vf3\_1 protein").

The nucleotide sequence of vf3\_1 as presently determined is reported in SEQ ID NO:55, and includes a poly(A) tail. What applicants presently believe to be the proper  
15 reading frame and the predicted amino acid sequence of the vf3\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:56. Amino acids 8 to 20 of SEQ ID NO:56 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
20 the predicted leader/signal sequence not be separated from the remainder of the vf3\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vf3\_1 should be approximately 2987 bp.

The nucleotide sequence disclosed herein for vf3\_1 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vf3\_1 demonstrated at least some similarity with sequences identified as Z78394 (*H.sapiens* mRNA, expressed sequence tag ICRFp507K11187 (5'), mRNA sequence). The predicted amino acid sequence disclosed herein for vf3\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the  
30 BLASTX search protocol. The predicted vf3\_1 protein demonstrated at least some similarity to the sequence identified as U41558 (K02B2.3 gene product [*Caenorhabditis elegans*]). Based upon sequence similarity, vf3\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two

additional potential transmembrane domains within the vf3\_1 protein sequence, one centered around amino acid 242 and another around amino acid 275 of SEQ ID NO:56.

vf3\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 39 kDa was detected in membrane fractions using SDS  
5 polyacrylamide gel electrophoresis.

#### Clone "vj2\_1"

A polynucleotide of the present invention has been identified as clone "vj2\_1". vj2\_1 was isolated from a human fetal brain (whole brain, enriched for G-protein-coupled  
10 receptors) cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vj2\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vj2\_1 protein").

The nucleotide sequence of vj2\_1 as presently determined is reported in SEQ ID  
15 NO:57, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vj2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:58. Amino acids 59 to 71 of SEQ ID NO:58 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 72. Due to the hydrophobic nature of the  
20 predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vj2\_1 protein.

Another potential vj2\_1 reading frame and predicted amino acid sequence that could be encoded by basepairs 146 to 400 of SEQ ID NO:57 is reported in SEQ ID NO:127.  
25 The TopPredII computer program predicts two potential transmembrane domains within the amino acid sequence of SEQ ID NO:127.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vj2\_1 should be approximately 1762 bp.

The nucleotide sequence disclosed herein for vj2\_1 was searched against the  
30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vj2\_1 demonstrated at least some similarity with sequences identified as N36445 (yx83c04.r1 Homo sapiens cDNA clone 268326 5'). Based upon sequence similarity, vj2\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three potential transmembrane

domains within the vj2\_1 protein sequence, centered around amino acids 30, 67, and 90 of SEQ ID NO:58, respectively. The nucleotide sequence of vj2\_1 indicates that it may contain one or more repetitive elements.

5        Clone "vp7\_1"

A polynucleotide of the present invention has been identified as clone "vp7\_1". vp7\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp7\_1 is a full-length clone, including the  
10    entire coding sequence of a secreted protein (also referred to herein as "vp7\_1 protein").

The nucleotide sequence of vp7\_1 as presently determined is reported in SEQ ID NO:59, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp7\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:60. Amino acids 6 to 18  
15    of SEQ ID NO:60 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 19. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp7\_1 protein. Another potential vp7\_1 reading frame and predicted amino acid sequence that  
20    could be encoded by basepairs 2071 to 2430 of SEQ ID NO:59 is reported in SEQ ID NO:128.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp7\_1 should be approximately 2638 bp.

The nucleotide sequence disclosed herein for vp7\_1 was searched against the  
25    GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp7\_1 demonstrated at least some similarity with sequences identified as N49433 (yv21e12.r1 Homo sapiens cDNA clone 243406 5') and Q63862 (AP2 sequence obtained by PCR for tumour specific DNA; standard; cDNA). Based upon sequence similarity, vp7\_1 proteins and each similar protein or peptide may share at least  
30    some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the vp7\_1 protein sequence centered around amino acid 75 of SEQ ID NO:60. The nucleotide sequence of vp7\_1 indicates that it may contain one or more Alu repeat sequences.

Clone "vp8\_1"

A polynucleotide of the present invention has been identified as clone "vp8\_1". vp8\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp8\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp8\_1 protein").

The nucleotide sequence of vp8\_1 as presently determined is reported in SEQ ID NO:61, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp8\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:62. Amino acids 20 to 32 of SEQ ID NO:62 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 33. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp8\_1 protein. If two insertions of "C" residues were made in the nucleotide sequence of SEQ ID NO:61, one after the "A" at position 380 and another after the "G" at position 382, the resulting nucleotide sequence would be predicted to encode the amino acid sequence reported in SEQ ID NO:129.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp8\_1 should be approximately 1513 bp.

The nucleotide sequence disclosed herein for vp8\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp8\_1 demonstrated at least some similarity with sequences identified as AA284421 (zs59c10.r1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 701778 5' similar to contains Alu repetitive element; mRNA sequence) and AC002086 (Human PAC clone DJ525N14 from Xq23, complete sequence). Based upon sequence similarity, vp8\_1 proteins and each similar protein or peptide may share at least some activity. Profile hidden markov model analysis reveals the presence of an SH2 domain in the predicted vp8\_1 protein (SEQ ID NO:62). SH2 domains function as regulatory modulators of intra-cellular signalling cascades by interacting with high affinity to phosphotyrosine-containing target peptides in a sequence-specific and strictly phosphorylation-dependent manner. The nucleotide sequence of vp8\_1 indicates that it may contain one or more Alu repeat sequences.



vp8\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 34 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5        Clone "vb22\_1"

A polynucleotide of the present invention has been identified as clone "vb22\_1". vb22\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb22\_1 is a full-length clone, including the  
10    entire coding sequence of a secreted protein (also referred to herein as "vb22\_1 protein").

The nucleotide sequence of vb22\_1 as presently determined is reported in SEQ ID NO:63, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb22\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:64.  
15    Another potential vb22\_1 reading frame and predicted amino acid sequence is encoded by basepairs 152 to 1006 of SEQ ID NO:63 and is reported in SEQ ID NO:130.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb22\_1 should be approximately 4176 bp.

The nucleotide sequence disclosed herein for vb22\_1 was searched against the  
20    GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb22\_1 demonstrated at least some similarity with sequences identified as L10335 (Homo sapiens neuroendocrine-specific protein C (NSP) mRNA, complete cds), N21304 (yx53f07.s1 Homo sapiens cDNA clone 265477 3' similar to SP:A60021 A60021 TROPOMYOSIN-RELATED PROTEIN, NEURONAL), and V23695  
25    (Human NSPLP protein A coding sequence; standard; cDNA). The predicted amino acid sequence disclosed herein for vb22\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vb22\_1 protein demonstrated at least some similarity to sequences identified as L10333 (nueroendocrine-specific protein A [Homo sapiens]) and W53947 (Human NSPLP protein  
30    A). Based upon sequence similarity, vb22\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the vb22\_1 protein sequence, one centered around amino acid 730 and another around amino acid 846 of SEQ ID NO:64. The nucleotide sequence of vb22\_1 appears to contain a short simple nucleotide repeat ("GGA") region.

Clone "vc48\_1"

A polynucleotide of the present invention has been identified as clone "vc48\_1". vc48\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc48\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc48\_1 protein").

The nucleotide sequence of vc48\_1 as presently determined is reported in SEQ ID NO:65, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc48\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:66. Amino acids 7 to 19 of SEQ ID NO:66 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc48\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc48\_1 should be approximately 3096 bp.

The nucleotide sequence disclosed herein for vc48\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc48\_1 demonstrated at least some similarity with sequences identified as AA292779 (zt56c06.s1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 726346 3', mRNA sequence). The predicted amino acid sequence disclosed herein for vc48\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc48\_1 protein demonstrated at least some similarity to sequences identified as AL031765 (Drosophila genomic product 22E5.z) and Z81058 (F11E6.e [Caenorhabditis elegans]). Based upon sequence similarity, vc48\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four potential transmembrane domains within the vc48\_1 protein sequence, one centered around amino acid 39 and others around amino acids 69, 107 and 134 of SEQ ID NO:66, respectively. The nucleotide sequence of vc48\_1 appears to contain a simple nucleotide repeat ("AC") and one or more of the following repetitive elements: Alu and MIR.

Clone "vp3\_1"

A polynucleotide of the present invention has been identified as clone "vp3\_1". vp3\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp3\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp3\_1 protein").

The nucleotide sequence of vp3\_1 as presently determined is reported in SEQ ID NO:67, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp3\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:68. Amino acids 19 to 31 of SEQ ID NO:68 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 32. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp3\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp3\_1 should be approximately 552 bp.

The nucleotide sequence disclosed herein for vp3\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp3\_1 demonstrated at least some similarity with sequences identified as AA225045 (nc34c06.r1 NCI\_CGAP\_Pr2 Homo sapiens cDNA clone IMAGE 1010026, mRNA sequence), M18157 (Human glandular kallikrein gene, complete cds), and T35868 (Prostate-specific antigen gene partial sequence; standard; DNA). Based upon sequence similarity, vp3\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vc61\_1"

A polynucleotide of the present invention has been identified as clone "vc61\_1". vc61\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc61\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc61\_1 protein").

The nucleotide sequence of vc61\_1 as presently determined is reported in SEQ ID NO:69, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the vc61\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:70. Amino acids 16 to 28 of SEQ ID NO:70 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 29. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc61\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc61\_1 should be approximately 3199 bp.

10 The nucleotide sequence disclosed herein for vc61\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc61\_1 demonstrated at least some similarity with sequences identified as AI028115 (ow51d09.x1 Soares\_parathyroid\_tumor\_NbHPA Homo sapiens cDNA clone IMAGE 1650353 3' similar to gb S67859 TRANSCRIPTION INITIATION  
15 FACTOR IIE-ALPHA CHAIN (HUMAN); mRNA), V20913 (Human induced tumour protein cDNA), and Z99129 (Human DNA sequence from clone 425C14 on chromosome 6q22 Contains the HSF2 gene for Heat Shock Factor 2 (Heat Shock Transcription Factor 2, HSTF 2) and an unknown gene similar to the placental protein DIFF33 gene; Contains ESTs, STSs and GSSs, complete sequence). The predicted amino acid sequence disclosed  
20 herein for vc61\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc61\_1 protein demonstrated at least some similarity to sequences identified as W52812 (Human induced tumour protein) and Z99129 (dJ425C14.2 (Placental protein DIFF33 LIKE) [Homo sapiens]). The deduced vc61\_1 protein has amino acid similarity to human and mouse diff33 protein.  
25 Diff33 is a transmembrane protein which is overexpressed in testicular tumors from polyomavirus large T-antigen transgenic mice. Based upon sequence similarity, vc61\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts nine additional potential transmembrane domains within the vc61\_1 protein sequence, centered around amino acids 50, 100, 150, 210, 240,  
30 270, 320, 390, and 430 of SEQ ID NO:70, respectively. The nucleotide sequence of vc61\_1 indicates that it may contain an Alu repetitive element.

Clone "vp15\_1"

A polynucleotide of the present invention has been identified as clone "vp15\_1". vp15\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp15\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp15\_1 protein").

The nucleotide sequence of vp15\_1 as presently determined is reported in SEQ ID NO:71, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp15\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:72. Amino acids 4 to 16 of SEQ ID NO:72 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp15\_1 protein. If a "C" residue were inserted between nucleotides 458 and 459 of SEQ ID NO:71, nucleotides 44 to 568 of the resulting nucleotide sequence would encode a protein having an amino acid sequence reported as SEQ ID NO:131.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp15\_1 should be approximately 2033 bp.

The nucleotide sequence disclosed herein for vp15\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp15\_1 demonstrated at least some similarity with sequences identified as AI033082 (ow97g04.s1 Soares\_fetal\_liver\_spleen\_1NFLS\_S1 Homo sapiens cDNA clone IMAGE 1654806 3', mRNA sequence) and T21877 (Human gene signature HUMGS03418). The predicted amino acid sequence disclosed herein for vp15\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vp15\_1 protein demonstrated at least some similarity to sequences identified as R45335 (Thrombomodulin analogue Q336N, Q365E) and U94333 (C1qR(p) [Homo sapiens]). The predicted vp15\_1 protein shows some amino acid similarity to multiple thrombomodulin analogues (such as GeneSeq accession number R45335), and shows some end-to-end similarity to GenPept accession number U94333, which is described as a "... human C1q/MBL/SPA receptor that mediates enhanced phagocytosis *in vitro*" (Nepomuceno *et al.*, 1997, *Immunity* 6(2): 119-129, which

is incorporated by reference herein). Based upon sequence similarity, vp15\_1 proteins and each similar protein or peptide may share at least some activity.

vp15\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 24 kDa was detected in conditioned medium and  
5 membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "vp17\_1"

A polynucleotide of the present invention has been identified as clone "vp17\_1". vp17\_1 was isolated from a human adult prostate cDNA library and was identified as  
10 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp17\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp17\_1 protein").

The nucleotide sequence of vp17\_1 as presently determined is reported in SEQ ID NO:73, and includes a poly(A) tail. What applicants presently believe to be the proper  
15 reading frame and the predicted amino acid sequence of the vp17\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:74. Amino acids 10 to 22 of SEQ ID NO:74 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain  
20 should the predicted leader/signal sequence not be separated from the remainder of the vp17\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp17\_1 should be approximately 3150 bp.

The nucleotide sequence disclosed herein for vp17\_1 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp17\_1 demonstrated at least some similarity with sequences identified as AI056890 (oz03g07.x1 Soares\_fetal\_liver\_spleen\_1NFLS\_S1 Homo sapiens cDNA clone IMAGE 1674300 3', mRNA sequence) and T64815 (Tumour suppressor activated pathway gene TSAP6). Based upon sequence similarity, vp17\_1 proteins and  
30 each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the vp17\_1 protein sequence, one centered around amino acid 50 and another around amino acid 80 of SEQ ID NO:74.

Clone "vp19\_1"

A polynucleotide of the present invention has been identified as clone "vp19\_1". vp19\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp19\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp19\_1 protein").

The nucleotide sequence of vp19\_1 as presently determined is reported in SEQ ID NO:75, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp19\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:76.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp19\_1 should be approximately 971 bp.

The nucleotide sequence disclosed herein for vp19\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp19\_1 demonstrated at least some similarity with sequences identified as AA716408 (zg64b02.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 398091 3', mRNA sequence) and T20711 (Human gene signature HUMGS01928). Based upon sequence similarity, vp19\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vp19\_1 protein sequence centered around amino acid 23 of SEQ ID NO:76; due to its hydrophobic nature, this region (amino acids 20 to 32) could also be a leader/signal sequence, with the mature protein beginning at amino acid 33 of SEQ ID NO:76.

Clone "vq1\_1"

A polynucleotide of the present invention has been identified as clone "vq1\_1". vq1\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq1\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq1\_1 protein").

The nucleotide sequence of vq1\_1 as presently determined is reported in SEQ ID NO:77, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq1\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:78. Amino acids 17 to 29

of SEQ ID NO:78 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq1\_1 protein. If a "T" residue were inserted between nucleotides 332 and 333 of SEQ ID NO:77, nucleotides 54 to 496 of the resulting nucleotide sequence would encode a protein having an amino acid sequence reported as SEQ ID NO:132.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq1\_1 should be approximately 873 bp.

The nucleotide sequence disclosed herein for vq1\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq1\_1 demonstrated at least some similarity with sequences identified as No hits were found in the databases. The TopPredII computer program predicts an additional potential transmembrane domain within the vq1\_1 protein sequence, extending from about amino acid 36 to about amino acid 76 of SEQ ID NO:78. The nucleotide sequence of vq1\_1 indicates that it may contain an Alu repetitive element.

#### Clone "vp14\_1"

A polynucleotide of the present invention has been identified as clone "vp14\_1". vp14\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp14\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp14\_1 protein").

The nucleotide sequence of vp14\_1 as presently determined is reported in SEQ ID NO:79, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp14\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:80. Amino acids 5 to 17 of SEQ ID NO:80 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp14\_1 protein.



The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp14\_1 should be approximately 1355 bp.

The nucleotide sequence disclosed herein for vp14\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp14\_1 demonstrated at least some similarity with sequences identified as AI052724 (oz27a12.x1 Soares\_total\_fetus\_Nb2HF8\_9w Homo sapiens cDNA clone IMAGE:1676542 3' similar to SW:YQJQ\_BACSU P54554 HYPOTHETICAL OXIDOREDUCTASE IN GLNQ-ANSR INTERGENIC REGION; mRNA sequence) and T20001 (Human gene signature HUMGS01138). The predicted amino acid sequence disclosed herein for vp14\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vp14\_1 protein demonstrated at least some similarity to sequences identified as R61477 (Clavulanic acid dehydrogenase sequence) and Z99116 (similar to ketoacyl reductase [Bacillus subtilis]). The predicted vp14\_1 protein shows some amino acid similarity to various dehydrogenases due to the presence of a short-chain alcohol dehydrogenase family signature at amino acids 51 to 240 of SEQ ID NO:80, as detected by motifs and hidden markov model analysis. Based upon sequence similarity, vp14\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four additional potential transmembrane domains within the vp14\_1 protein sequence, centered around amino acids 55, 195, 230, and 300 of SEQ ID NO:80, respectively.

#### Deposit of Clones

Clones vb11\_1, vb12\_1, vb14\_1, ve11\_1, vf2\_1, vg2\_1, vj1\_1, and vl1\_1 were deposited on August 20, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98846, from which each clone comprising a particular polynucleotide is obtainable.

Clone vk2\_1 was deposited on August 20, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number 98838, from which the vk2\_1 clone comprising a particular polynucleotide is obtainable.

Clones vb21\_1, vc35\_1, vc36\_1, vc38\_1, vc39\_1, vc40\_1, vc46\_1, vc49\_1, vc50\_1, vc51\_1, and vc52\_1 were deposited on September 2, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98862, from which each clone comprising a particular polynucleotide is obtainable.

Clones vc33\_1, vc34\_1, vc47\_1, vc54\_1, vc57\_1, ve13\_1, ve16\_1, vf3\_1, vj2\_1, vp7\_1, and vp8\_1 were deposited on September 22, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98886, from which each clone comprising a particular polynucleotide is obtainable.

Clones vb22\_1, vc48\_1, and vp3\_1 were deposited on October 16, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98933, from which each clone comprising a particular polynucleotide is obtainable.

Clones vc61\_1, vp15\_1, vp17\_1, vp19\_1, and vq1\_1 were deposited on December 23, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 207012, from which each clone comprising a particular polynucleotide is obtainable.

Clone vp14\_1 was deposited on December 23, 1998 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number 207011, from which the vp14\_1 clone comprising a particular polynucleotide is obtainable.

All restrictions on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent, except for the requirements specified in 37 C.F.R. § 1.808(b), and the term of the deposit will comply with 37 C.F.R. § 1.806.

Each clone has been transfected into separate bacterial cells (*E. coli*) in this composite deposit. Each clone can be removed from the vector in which it was deposited by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI) to produce the appropriate fragment for such clone. Each clone was deposited in either the pED6 or pNOTs vector depicted in Figures 1A and 1B, respectively. The pED6dpc2 vector

("pED6") was derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning (Kaufman *et al.*, 1991, *Nucleic Acids Res.* 19: 4485-4490); the pNOTs vector was derived from pMT2 (Kaufman *et al.*, 1989, *Mol. Cell. Biol.* 9: 946-958) by deletion of the DHFR sequences, insertion of a new polylinker, and insertion of the M13 origin of replication in the ClaI site. In some instances, the deposited clone can become "flipped" (i.e., in the reverse orientation) in the deposited isolate. In such instances, the cDNA insert can still be isolated by digestion with EcoRI and NotI. However, NotI will then produce the 5' site and EcoRI will produce the 3' site for placement of the cDNA in proper orientation for expression in a suitable vector. The cDNA may also be expressed from the vectors in which they were deposited.

Bacterial cells containing a particular clone can be obtained from the composite deposit as follows:

An oligonucleotide probe or probes should be designed to the sequence that is known for that particular clone. This sequence can be derived from the sequences provided herein, or from a combination of those sequences. The sequence of an oligonucleotide probe that was used to isolate or to sequence each full-length clone is identified below, and should be most reliable in isolating the clone of interest.

	<u>Clone</u>	<u>Probe Sequence</u>
20	vb11_1	SEQ ID NO:81
	vb12_1	SEQ ID NO:82
	vb14_1	SEQ ID NO:83
	ve11_1	SEQ ID NO:84
	vf2_1	SEQ ID NO:85
25	vg2_1	SEQ ID NO:86
	vj1_1	SEQ ID NO:87
	vl1_1	SEQ ID NO:88
	vk2_1	SEQ ID NO:89
	vb21_1	SEQ ID NO:90
30	vc35_1	SEQ ID NO:91
	vc36_1	SEQ ID NO:92
	vc38_1	SEQ ID NO:93
	vc39_1	SEQ ID NO:94
	vc40_1	SEQ ID NO:95

	vc46_1	SEQ ID NO:96
	vc49_1	SEQ ID NO:97
	vc50_1	SEQ ID NO:98
	vc51_1	SEQ ID NO:99
5	vc52_1	SEQ ID NO:100
	vc33_1	SEQ ID NO:101
	vc34_1	SEQ ID NO:102
	vc47_1	SEQ ID NO:103
	vc54_1	SEQ ID NO:104
10	vc57_1	SEQ ID NO:105
	ve13_1	SEQ ID NO:106
	ve16_1	SEQ ID NO:107
	vf3_1	SEQ ID NO:108
	vj2_1	SEQ ID NO:109
15	vp7_1	SEQ ID NO:110
	vp8_1	SEQ ID NO:111
	vb22_1	SEQ ID NO:112
	vc48_1	SEQ ID NO:113
	vp3_1	SEQ ID NO:114
20	vc61_1	SEQ ID NO:115
	vp15_1	SEQ ID NO:116
	vp17_1	SEQ ID NO:117
	vp19_1	SEQ ID NO:118
	vq1_1	SEQ ID NO:119
25	vp14_1	SEQ ID NO:120

In the sequences listed above which include an N at position 2, that position is occupied in preferred probes/primers by a biotinylated phosphoramidite residue rather than a nucleotide (such as, for example, that produced by use of biotin phosphoramidite (1-dimethoxytrityloxy-2-(N-biotinyl-4-aminobutyl)-propyl-3-O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidite) (Glen Research, cat. no. 10-1953)).

The design of the oligonucleotide probe should preferably follow these parameters:

- (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any;
  - (b) It should be designed to have a  $T_m$  of approx. 80 ° C (assuming 2° for each A or T and 4 degrees for each G or C).
- 5 The oligonucleotide should preferably be labeled with  $\gamma$ -<sup>32</sup>P ATP (specific activity 6000 Ci/mmol) and T4 polynucleotide kinase using commonly employed techniques for labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography or other established methods. The amount of radioactivity incorporated into the probe should be quantitated
- 10 by measurement in a scintillation counter. Preferably, specific activity of the resulting probe should be approximately 4e+6 dpm/pmol.

The bacterial culture containing the pool of full-length clones should preferably be thawed and 100 µl of the stock used to inoculate a sterile culture flask containing 25 ml of sterile L-broth containing ampicillin at 100 µg/ml. The culture should preferably be

15 grown to saturation at 37°C, and the saturated culture should preferably be diluted in fresh L-broth. Aliquots of these dilutions should preferably be plated to determine the dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing L-broth containing ampicillin at 100 µg/ml and agar at 1.5% in a 150 mm petri dish when grown overnight at 37°C. Other

20 known methods of obtaining distinct, well-separated colonies can also be employed.

Standard colony hybridization procedures should then be used to transfer the colonies to nitrocellulose filters and lyse, denature and bake them.

The filter is then preferably incubated at 65°C for 1 hour with gentle agitation in 6X SSC (20X stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with

25 NaOH) containing 0.5% SDS, 100 µg/ml of yeast RNA, and 10 mM EDTA (approximately 10 mL per 150 mm filter). Preferably, the probe is then added to the hybridization mix at a concentration greater than or equal to 1e+6 dpm/mL. The filter is then preferably incubated at 65°C with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2X SSC/0.5% SDS at room temperature without agitation, preferably followed

30 by 500 mL of 2X SSC/0.1% SDS at room temperature with gentle shaking for 15 minutes. A third wash with 0.1X SSC/0.5% SDS at 65°C for 30 minutes to 1 hour is optional. The filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed.

The positive colonies are picked, grown in culture, and plasmid DNA isolated using standard procedures. The clones can then be verified by restriction analysis, hybridization analysis, or DNA sequencing.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length form of the such proteins is identified in the sequence listing by translation of the nucleotide sequence of each disclosed clone. The mature form(s) of such protein may be obtained by expression of the disclosed full-length polynucleotide (preferably those deposited with the ATCC) in a suitable mammalian cell or other host cell. The sequence(s) of the mature form(s) of the protein may also be determinable from the amino acid sequence of the full-length form.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that

has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.

The chromosomal location corresponding to the polynucleotide sequences disclosed herein may also be determined, for example by hybridizing appropriately  
5 labeled polynucleotides of the present invention to chromosomes *in situ*. It may also be possible to determine the corresponding chromosomal location for a disclosed polynucleotide by identifying significantly similar nucleotide sequences in public  
10 databases, such as expressed sequence tags (ESTs), that have already been mapped to particular chromosomal locations. For at least some of the polynucleotide sequences disclosed herein, public database sequences having at least some similarity to the  
15 polynucleotide of the present invention have been listed by database accession number. Searches using the GenBank accession numbers of these public database sequences can then be performed at an Internet site provided by the National Center for Biotechnology Information having the address <http://www.ncbi.nlm.nih.gov/UniGene/>, in order to  
20 identify "UniGene clusters" of overlapping sequences. Many of the "UniGene clusters" so identified will already have been mapped to particular chromosomal sites.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The desired change in gene expression can be achieved through the use of antisense  
20 polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, *Trends Pharmacol. Sci.* 15(7): 250-254; Lavarosky *et al.*, 1997, *Biochem. Mol. Med.* 62(1): 11-22; and Hampel, 1998, *Prog. Nucleic Acid Res. Mol. Biol.* 58: 1-39; all of which are incorporated by reference herein). The desired change in gene expression can also be achieved through the use of double-stranded ribonucleotide  
25 molecules having some complementarity to the mRNA transcribed from the gene, and which interfere with the transcription, stability, or expression of the mRNA ("RNA interference" or "RNAi"; Fire *et al.*, 1998, *Nature* 391 (6669): 806-811; Montgomery *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* 95 (26): 15502-15507; and Sharp, 1999, *Genes Dev.* 13 (2): 139-141; all of which are incorporated by reference herein). Transgenic animals that have  
30 multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are

also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through  
5 deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, *Bioessays* 14(9): 629-633; Zwaal *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* 90(16): 7431-7435; Clark *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination,  
10 preferably detected by positive/negative genetic selection strategies (Mansour *et al.*, 1988, *Nature* 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models for  
15 the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms, part  
20 or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information. For example, the TopPredII computer program can be used to predict the location of  
25 transmembrane domains in an amino acid sequence, domains which are described by the location of the center of the transmembrane domain, with at least ten transmembrane amino acids on each side of the reported central residue(s).

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most  
30 preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are



proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

5 In particular, sequence identity may be determined using WU-BLAST (Washington University BLAST) version 2.0 software, which builds upon WU-BLAST version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul and Gish, 1996, Local alignment statistics, Doolittle *ed.*, *Methods in Enzymology* 266: 460-480; Altschul *et al.*, 1990, Basic local alignment search tool, *Journal of*  
10 *Molecular Biology* 215: 403-410; Gish and States, 1993, Identification of protein coding regions by database similarity search, *Nature Genetics* 3: 266-272; Karlin and Altschul, 1993, Applications and statistics for multiple high-scoring segments in molecular sequences, *Proc. Natl. Acad. Sci. USA* 90: 5873-5877; all of which are incorporated by reference herein). WU-BLAST version 2.0 executable programs for several UNIX  
15 platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. The complete suite of search programs (BLASTP, BLASTN, BLASTX, TBLASTN, and TBLASTX) is provided at that site, in addition to several support programs. WU-BLAST 2.0 is copyrighted and may not be sold or redistributed in any form or manner without the express written consent of the author; but the posted executables may otherwise be freely  
20 used for commercial, nonprofit, or academic purposes. In all search programs in the suite -- BLASTP, BLASTN, BLASTX, TBLASTN and TBLASTX -- the gapped alignment routines are integral to the database search itself, and thus yield much better sensitivity and selectivity while producing the more easily interpreted output. Gapping can optionally be turned off in all of these programs, if desired. The default penalty (Q) for a gap of length  
25 one is Q=9 for proteins and BLASTP, and Q=10 for BLASTN, but may be changed to any integer value including zero, one through eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer value including zero, one, two, three, four, five, six,  
30 seven, eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. Any combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps.

The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized.

Species homologues of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or  
5 polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with  
10 the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from  
15 the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*,  
20 *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis catus*, *Mustela vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species  
25 (O'Brien and Seuáñez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien *et al.*, 1993, *Nature Genetics* 3:103-112; Johansson *et al.*, 1995, *Genomics* 25: 682-690; Lyons *et al.*, 1997, *Nature Genetics* 15: 47-56; O'Brien *et al.*, 1997, *Trends in Genetics* 13(10): 393-399; Carver and Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed polynucleotides  
30 or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotides which also encode proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90%

identity) with the given polynucleotide, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided herein and  
5 screening a suitable nucleic acid source from individuals of the appropriate species.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides that hybridize under reduced stringency conditions, more preferably stringent conditions, and most preferably highly  
10 stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) <sup>†</sup>	Hybridization Temperature and Buffer <sup>‡</sup>	Wash Temperature and Buffer <sup>‡</sup>
5	A	≥ 50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
	B	<50	T <sub>B</sub> <sup>*</sup> ; 1xSSC	T <sub>B</sub> <sup>*</sup> ; 1xSSC
	C	≥ 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC
	D	<50	T <sub>D</sub> <sup>*</sup> ; 1xSSC	T <sub>D</sub> <sup>*</sup> ; 1xSSC
	E	≥ 50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
	F	<50	T <sub>F</sub> <sup>*</sup> ; 1xSSC	T <sub>F</sub> <sup>*</sup> ; 1xSSC
10	G	≥ 50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
	H	<50	T <sub>H</sub> <sup>*</sup> ; 4xSSC	T <sub>H</sub> <sup>*</sup> ; 4xSSC
	I	≥ 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
	J	<50	T <sub>J</sub> <sup>*</sup> ; 4xSSC	T <sub>J</sub> <sup>*</sup> ; 4xSSC
	K	≥ 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
	L	<50	T <sub>L</sub> <sup>*</sup> ; 2xSSC	T <sub>L</sub> <sup>*</sup> ; 2xSSC
15	M	≥ 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
	N	<50	T <sub>N</sub> <sup>*</sup> ; 6xSSC	T <sub>N</sub> <sup>*</sup> ; 6xSSC
	O	≥ 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
	P	<50	T <sub>P</sub> <sup>*</sup> ; 6xSSC	T <sub>P</sub> <sup>*</sup> ; 6xSSC
	Q	≥ 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
	R	<50	T <sub>R</sub> <sup>*</sup> ; 4xSSC	T <sub>R</sub> <sup>*</sup> ; 4xSSC

†: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

‡: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub>, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

\*T<sub>B</sub> - T<sub>R</sub>: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T<sub>m</sub>) of the hybrid, where T<sub>m</sub> is determined according to the following equations. For hybrids less than 18 base pairs in length, T<sub>m</sub>(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T<sub>m</sub>(°C) = 81.5 + 16.6(log<sub>10</sub>[Na<sup>+</sup>]) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na<sup>+</sup>] is the concentration of sodium ions in the hybridization buffer ([Na<sup>+</sup>] for 1xSSC = 0.165 M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

The isolated polynucleotide encoding the protein of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial

strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *e.g.*, Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLabs (Beverly, MA), Pharmacia (Piscataway, NJ) and Invitrogen Corporation (Carlsbad, CA), respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from the Eastman Kodak Company (New Haven, CT).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art

given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

### USES AND BIOLOGICAL ACTIVITY

5       The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies  
10   or vectors suitable for introduction of DNA).

#### Research Uses and Utilities

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express  
15   recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare  
20   with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for  
25   examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, those  
30   described in Gyuris *et al.*, 1993, *Cell* 75: 791-803 and in Rossi *et al.*, 1997, *Proc. Natl. Acad. Sci. USA* 94: 8405-8410, all of which are incorporated by reference herein) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.



The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine  
5 levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the  
10 protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent  
15 grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to  
20 Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

#### Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein  
25 or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention  
30 can be added to the medium in or on which the microorganism is cultured.

#### Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may

induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

- 10        Assays for T-cell or thymocyte proliferation include without limitation those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., *J. Immunol.* 137:3494-3500, 1986;
- 15        Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J. Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152: 1756-1761, 1994.

- 20        Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon  $\gamma$ , Schreiber, R.D. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

- 25        Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., *J. Exp. Med.* 173:1205-1211, 1991; Moreau et al., *Nature* 336:690-692, 1988; Greenberger et al., *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-2938, 1983;
- 30        Measurement of mouse and human interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991;

Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

- Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

15        Immune Stimulating or Suppressing Activity

- A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, *Leishmania* spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

- 30        Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease.

Such a protein of the present invention may also to be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to regulate immune responses in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), *e.g.*, preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (*e.g.*, B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term

tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

- 5           The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as
- 10 described in Lenschow *et al.*, Science 257:789-792 (1992) and Turka *et al.*, Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.
- 15           Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms.
- 20 Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from
- 25 the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/*lpr/lpr* mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and
- 30 murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune

response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (*e.g.*, a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  chain protein and  $\beta_2$

microglobulin protein or an MHC class II  $\alpha$  chain protein and an MHC class II  $\beta$  chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated  
5 immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated  
10 immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without  
15 limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Herrmann et al., *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann et al., *J. Immunol.* 128:1968-1974, 1982; Handa et al.,  
20 *J. Immunol.* 135:1564-1572, 1985; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Takai et al., *J. Immunol.* 140:508-512, 1988; Herrmann et al., *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann et al., *J. Immunol.* 128:1968-1974, 1982; Handa et al., *J. Immunol.* 135:1564-1572, 1985; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bowman et al., *J. Virology* 61:1992-1998; Takai et al., *J. Immunol.* 140:508-512, 1988; Bertagnolli et al.,  
25 *Cellular Immunology* 133:327-341, 1991; Brown et al., *J. Immunol.* 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, *J. Immunol.* 144:3028-3033, 1990; and Assays for B cell function: *In vitro*  
30 antibody production, Mond, J.J. and Brunswick, M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek,

D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

- 5 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995;
- 10 Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

- Assays for lymphocyte survival/apoptosis (which will identify, among others,
- 15 proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993;
- 20 Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad. Sci. USA 88:7548-7551, 1991.

25

#### Hematopoiesis Regulating Activity

- A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell
- 30 lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid



cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and

Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

5

#### Tissue Growth Activity

A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns,  
10 incisions and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as  
15 well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal  
20 disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue  
25 destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in  
30 circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and

in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation

of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium ).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

#### Activin/Inhibin Activity

A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin  $\alpha$  family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- $\beta$  group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

#### Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al.

APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

#### Hemostatic and Thrombolytic Activity

5 A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting  
10 formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

15 Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

#### Receptor/Ligand Activity

20 A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands,  
25 receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant  
30 receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static  
5 conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

10        Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the  
15 inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic  
20 inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

25

Cadherin/Tumor Invasion Suppressor Activity

Cadherins are calcium-dependent adhesion molecules that appear to play major roles during development, particularly in defining specific cell types. Loss or alteration of normal cadherin expression can lead to changes in cell adhesion properties linked to  
30 tumor growth and metastasis. Cadherin malfunction is also implicated in other human diseases, such as pemphigus vulgaris and pemphigus foliaceus (auto-immune blistering skin diseases), Crohn's disease, and some developmental abnormalities.

The cadherin superfamily includes well over forty members, each with a distinct pattern of expression. All members of the superfamily have in common conserved

extracellular repeats (cadherin domains), but structural differences are found in other parts of the molecule. The cadherin domains bind calcium to form their tertiary structure and thus calcium is required to mediate their adhesion. Only a few amino acids in the first cadherin domain provide the basis for homophilic adhesion; modification of this  
5 recognition site can change the specificity of a cadherin so that instead of recognizing only itself, the mutant molecule can now also bind to a different cadherin. In addition, some cadherins engage in heterophilic adhesion with other cadherins.

E-cadherin, one member of the cadherin superfamily, is expressed in epithelial cell types. Pathologically, if E-cadherin expression is lost in a tumor, the malignant cells  
10 become invasive and the cancer metastasizes. Transfection of cancer cell lines with polynucleotides expressing E-cadherin has reversed cancer-associated changes by returning altered cell shapes to normal, restoring cells' adhesiveness to each other and to their substrate, decreasing the cell growth rate, and drastically reducing anchorage-independent cell growth. Thus, reintroducing E-cadherin expression reverts carcinomas  
15 to a less advanced stage. It is likely that other cadherins have the same invasion suppressor role in carcinomas derived from other tissue types. Therefore, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to treat cancer. Introducing such proteins or polynucleotides into cancer cells can reduce or eliminate the cancerous changes observed  
20 in these cells by providing normal cadherin expression.

Cancer cells have also been shown to express cadherins of a different tissue type than their origin, thus allowing these cells to invade and metastasize in a different tissue in the body. Proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be substituted in these cells for the  
25 inappropriately expressed cadherins, restoring normal cell adhesive properties and reducing or eliminating the tendency of the cells to metastasize.

Additionally, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to generate antibodies recognizing and binding to cadherins. Such antibodies can be used to block  
30 the adhesion of inappropriately expressed tumor-cell cadherins, preventing the cells from forming a tumor elsewhere. Such an anti-cadherin antibody can also be used as a marker for the grade, pathological type, and prognosis of a cancer, i.e. the more progressed the cancer, the less cadherin expression there will be, and this decrease in cadherin expression can be detected by the use of a cadherin-binding antibody.



Fragments of proteins of the present invention with cadherin activity, preferably a polypeptide comprising a decapeptide of the cadherin recognition site, and polynucleotides of the present invention encoding such protein fragments, can also be used to block cadherin function by binding to cadherins and preventing them from binding in ways that produce undesirable effects. Additionally, fragments of proteins of the present invention with cadherin activity, preferably truncated soluble cadherin fragments which have been found to be stable in the circulation of cancer patients, and polynucleotides encoding such protein fragments, can be used to disturb proper cell-cell adhesion.

Assays for cadherin adhesive and invasive suppressor activity include, without limitation, those described in: Hortsch et al. J Biol Chem 270 (32): 18809-18817, 1995; Miyaki et al. Oncogene 11: 2547-2552, 1995; Ozawa et al. Cell 63: 1033-1038, 1990.

#### Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via antibody-dependent cell-mediated cytotoxicity (ADCC)). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth.

#### Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s);

effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic  
5 lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another  
10 material or entity which is cross-reactive with such protein.

#### ADMINISTRATION AND DOSING

A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be used in a  
15 pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the  
20 carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other  
25 agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor,  
30 thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical

compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in  
5 combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other  
10 hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical  
15 composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is  
20 administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention.  
25 When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid  
30 form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present

invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu$ g to about 100 mg (preferably about 0.1mg to about 10 mg, more preferably about 0.1  $\mu$ g to about 1 mg) of protein of the present invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. As used herein, the term "antibody" includes without limitation a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a single-chain antibody, a CDR-grafted antibody, a humanized antibody, or fragments thereof which bind to the indicated protein.

Such term also includes any other species derived from an antibody or antibody sequence which is capable of binding the indicated protein.

Antibodies to a particular protein can be produced by methods well known to those skilled in the art. For example, monoclonal antibodies can be produced by generation of  
5 antibody-producing hybridomas in accordance with known methods (see for example, Goding, 1983, *Monoclonal antibodies: principles and practice*, Academic Press Inc., New York; and Yokoyama, 1992, "Production of Monoclonal Antibodies" in *Current Protocols in Immunology*, Unit 2.5, Greene Publishing Assoc. and John Wiley & Sons). Polyclonal sera and antibodies can be produced by inoculation of a mammalian subject with the  
10 relevant protein or fragments thereof in accordance with known methods. Fragments of antibodies, receptors, or other reactive peptides can be produced from the corresponding antibodies by cleavage of and collection of the desired fragments in accordance with known methods (see for example, Goding, *supra*; and Andrew et al., 1992, "Fragmentation of Immunoglobulins" in *Current Protocols in Immunology*, Unit 2.8, Greene Publishing  
15 Assoc. and John Wiley & Sons). Chimeric antibodies and single chain antibodies can also be produced in accordance with known recombinant methods (see for example, 5,169,939, 5,194,594, and 5,576,184). Humanized antibodies can also be made from corresponding murine antibodies in accordance with well known methods (see for example, U.S. Patent Nos. 5,530,101, 5,585,089, and 5,693,762). Additionally, human antibodies may be  
20 produced in non-human animals such as mice that have been genetically altered to express human antibody molecules (see for example Fishwild *et al.*, 1996, *Nature Biotechnology* 14: 845-851; Mendez *et al.*, 1997, *Nature Genetics* 15: 146-156 (erratum *Nature Genetics* 16: 410); and U.S. Patents 5,877,397 and 5,625,126). Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide  
25 immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J. Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, *FEBS Lett.* 211, 10 (1987).

Monoclonal antibodies binding to the protein of the invention may be useful  
30 diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where

abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

5       For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably  
10 be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the  
15 methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical  
20 applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium  
25 sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxapatite, bioglass, aluminates, or other  
30 ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions  
5 from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of  
10 carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to  
15 provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in  
20 question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to  
25 humans, are desired patients for such treatment with proteins of the present invention.

The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of  
30 a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect



the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such  
5 polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

Cells may also be cultured *ex vivo* in the presence of proteins of the present  
10 invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

Patent and literature references cited herein are incorporated by reference as if fully set forth.

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) the nucleotide sequence of SEQ ID NO:1;
  - (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934;
  - (c) the nucleotide sequence of the full-length protein coding sequence of clone vb11\_1 deposited with the ATCC under accession number 98846;
  - (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;
  - (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
  - (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
  - (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
  - (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:1.
2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.
3. A host cell transformed with the polynucleotide of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:

- (a) growing a culture of a host cell in a suitable culture medium, wherein the host cell has been transformed with the polynucleotide of claim 2; and
  - (b) purifying said protein from the culture.
- 6. A protein produced according to the process of claim 5.
- 7. An isolated polynucleotide encoding the protein of claim 6.
- 8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846.
- 9. A protein comprising an amino acid sequence selected from the group consisting of:
  - (a) the amino acid sequence of SEQ ID NO:2;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;the protein being substantially free from other mammalian proteins.
- 10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.
- 11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.
- 12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) the nucleotide sequence of SEQ ID NO:3;
  - (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482;
  - (c) the nucleotide sequence of SEQ ID NO:3 from nucleotide 201 to nucleotide 482;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
- (b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:5;
- (b) the nucleotide sequence of SEQ ID NO:5 from nucleotide 1195 to nucleotide 1527;
- (c) the nucleotide sequence of SEQ ID NO:5 from nucleotide 1468 to nucleotide 1527;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
- (b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(d) the nucleotide sequence of the full-length protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

(f) the nucleotide sequence of a mature protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

(b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:5;
- (b) the nucleotide sequence of SEQ ID NO:5 from nucleotide 1195 to nucleotide 1527;
- (c) the nucleotide sequence of SEQ ID NO:5 from nucleotide 1468 to nucleotide 1527;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
- (b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846; the protein being substantially free from other mammalian proteins.

16. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:7;
- (b) the nucleotide sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294;
- (c) the nucleotide sequence of SEQ ID NO:7 from nucleotide 109 to nucleotide 294;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:8;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:7.



17. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

18. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:9;
- (b) the nucleotide sequence of SEQ ID NO:9 from nucleotide 22 to nucleotide 468;
- (c) the nucleotide sequence of SEQ ID NO:9 from nucleotide 118 to nucleotide 468;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:9.

19. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
- (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

20. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:11;
- (b) the nucleotide sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641;
- (c) the nucleotide sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:12;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:11.

21. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:12;

(b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and

(c) the amino acid sequence encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:13;

(b) the nucleotide sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892;

(c) the nucleotide sequence of SEQ ID NO:13 from nucleotide 416 to nucleotide 892;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

(f) the nucleotide sequence of a mature protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:14;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:13.

23. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:14;

(b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and

(c) the amino acid sequence encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

24. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:15;

(b) the nucleotide sequence of SEQ ID NO:15 from nucleotide 62 to nucleotide 1057;

(c) the nucleotide sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057;

(d) the nucleotide sequence of the full-length protein coding sequence of clone v11\_1 deposited with the ATCC under accession number 98846;

- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:15.

25. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
- (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

26. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:17;

- (b) the nucleotide sequence of SEQ ID NO:17 from nucleotide 74 to nucleotide 529;
- (c) the nucleotide sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:17.

27. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:18;
- (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;

the protein being substantially free from other mammalian proteins.

28. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:19;
- (b) the nucleotide sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170;
- (c) the nucleotide sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:19.

29. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

30. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 74 to nucleotide 1453;
- (c) the nucleotide sequence of SEQ ID NO:21 from nucleotide 224 to nucleotide 1453;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees



C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:21.

31. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

32. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:23;
- (b) the nucleotide sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368;
- (c) the nucleotide sequence of SEQ ID NO:23 from nucleotide 243 to nucleotide 368;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:23.

33. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:24;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

34. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:25;
- (b) the nucleotide sequence of SEQ ID NO:25 from nucleotide 370 to nucleotide 1662;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vc38\_1 deposited with the ATCC under accession number 98862;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:26;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;

- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:25.

35. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:26;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

36. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:27;
- (b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365;
- (c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:28;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

37. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:28;

(b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

38. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:29;

(b) the nucleotide sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066;

(c) the nucleotide sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;

- (f) the nucleotide sequence of a mature protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:29.

39. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
- (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

40. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:31;
- (b) the nucleotide sequence of SEQ ID NO:31 from nucleotide 38 to nucleotide 553;
- (c) the nucleotide sequence of SEQ ID NO:31 from nucleotide 104 to nucleotide 553;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:32;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:31.

41. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:32;

(b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

42. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:33;
- (b) the nucleotide sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548;
- (c) the nucleotide sequence of SEQ ID NO:33 from nucleotide 242 to nucleotide 2548;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:33.

43. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
- (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862; the protein being substantially free from other mammalian proteins.

44. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:35;
- (b) the nucleotide sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776;
- (c) the nucleotide sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:35.



45. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:36;
- (b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

46. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:37;
- (b) the nucleotide sequence of SEQ ID NO:37 from nucleotide 139 to nucleotide 1308;
- (c) the nucleotide sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:37.

47. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

48. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:39;
- (b) the nucleotide sequence of SEQ ID NO:39 from nucleotide 21 to nucleotide 1142;
- (c) the nucleotide sequence of SEQ ID NO:39 from nucleotide 114 to nucleotide 1142;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:40;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:39.

49. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:40;

(b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

50. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:41;

(b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416;

(c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 346 to nucleotide 1416;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:41.

51. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:42;

(b) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

52. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:43;

(b) the nucleotide sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461;

(c) the nucleotide sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:44;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:43.

53. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

54. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:45;

- (b) the nucleotide sequence of SEQ ID NO:45 from nucleotide 1922 to nucleotide 2350;
- (c) the nucleotide sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:46;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:45.

55. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
- (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

56. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:47;
- (b) the nucleotide sequence of SEQ ID NO:47 from nucleotide 111 to nucleotide 1337;
- (c) the nucleotide sequence of SEQ ID NO:47 from nucleotide 246 to nucleotide 1337;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:47.

57. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- the protein being substantially free from other mammalian proteins.

58. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:49;
- (b) the nucleotide sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637;
- (c) the nucleotide sequence of SEQ ID NO:49 from nucleotide 270 to nucleotide 1637;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees



C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:49.

59. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- the protein being substantially free from other mammalian proteins.

60. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:51;
- (b) the nucleotide sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934;
- (c) the nucleotide sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:51.

61. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:52;
- (b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

62. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:53;
- (b) the nucleotide sequence of SEQ ID NO:53 from nucleotide 240 to nucleotide 503;
- (c) the nucleotide sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:54;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:53.

63. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

(b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and

(c) the amino acid sequence encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

64. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:55;

(b) the nucleotide sequence of SEQ ID NO:55 from nucleotide 11 to nucleotide 1063;

(c) the nucleotide sequence of SEQ ID NO:55 from nucleotide 71 to nucleotide 1063;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:56;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:55.

65. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:56;

(b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and

(c) the amino acid sequence encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

66. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:57;

(b) the nucleotide sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886;

(c) the nucleotide sequence of SEQ ID NO:57 from nucleotide 755 to nucleotide 886;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:57.

67. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:58;
- (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

68. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:59;
- (b) the nucleotide sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344;
- (c) the nucleotide sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:59.

69. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
- (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886; the protein being substantially free from other mammalian proteins.

70. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:61;
- (b) the nucleotide sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757;
- (c) the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:62;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:61.

71. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:62;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- the protein being substantially free from other mammalian proteins.

72. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:63;
- (b) the nucleotide sequence of SEQ ID NO:63 from nucleotide 1048 to nucleotide 3726;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vb22\_1 deposited with the ATCC under accession number 98933;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:63.

73. A protein comprising an amino acid sequence selected from the group consisting of:



- (a) the amino acid sequence of SEQ ID NO:64;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- the protein being substantially free from other mammalian proteins.

74. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:65;
- (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 134 to nucleotide 667;
- (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 191 to nucleotide 667;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.

75. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- the protein being substantially free from other mammalian proteins.

76. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:67;
- (b) the nucleotide sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457;
- (c) the nucleotide sequence of SEQ ID NO:67 from nucleotide 158 to nucleotide 457;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:67.

77. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- the protein being substantially free from other mammalian proteins.

78. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:69;
- (b) the nucleotide sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387;
- (c) the nucleotide sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:70;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:69.

79. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:70;

(b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

80. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:71;

(b) the nucleotide sequence of SEQ ID NO:71 from nucleotide 44 to nucleotide 1513;

(c) the nucleotide sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513;

(d) the nucleotide sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458;

(e) the nucleotide sequence of the full-length protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012;

(f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

(g) the nucleotide sequence of a mature protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012;

(h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

(i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:72;

(j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and

(l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:71.

81. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:72;

(b) the amino acid sequence of SEQ ID NO:72 from amino acid 1 to amino acid 139;

(c) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and

(d) the amino acid sequence encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

82. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:73;
- (b) the nucleotide sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743;
- (c) the nucleotide sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:73.

83. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
- (b) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012; the protein being substantially free from other mammalian proteins.

84. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:75;
- (b) the nucleotide sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:75.

85. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
- (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

86. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:77;
- (b) the nucleotide sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide 368;
- (c) the nucleotide sequence of SEQ ID NO:77 from nucleotide 141 to nucleotide 368;
- (d) the nucleotide sequence of SEQ ID NO:77 from nucleotide 51 to nucleotide 332;
- (e) the nucleotide sequence of the full-length protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (g) the nucleotide sequence of a mature protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
- (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:77.



87. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
- (b) the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 93;
- (c) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
- (d) the amino acid sequence encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

88. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:79;
- (b) the nucleotide sequence of SEQ ID NO:79 from nucleotide 2 to nucleotide 1018;
- (c) the nucleotide sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:79.

89. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:80;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- the protein being substantially free from other mammalian proteins.

Fig. 1A





Fig. 1B



## SEQUENCE LISTING

<110> Valenzuela, Dario  
 Yuan, Olive  
 Hoffman, Heidi  
 Hall, Jeff  
 Rapiejko, Peter

<120> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

<130> GI 6908X

<140>

<141>

<160> 132

<170> PatentIn Ver. 2.0

<210> 1

<211> 1738

<212> DNA

<213> Homo sapiens

<400> 1

```

acttagtgta gcaccagggg gccctagagc tggaggatat cgaatagatt aaatthttgct 60
cgtctcttcc acaagcccta accatgggtc ttaaaaacag cagattctgg gagccttcca 120
tgctctctct ctctctctct ttatctactt ccctcccaaa tgagagagtg acagagaatt 180
gtttttttat aaatcgaagt ttcttaatat tatcagggtt tgatacgtca gtggtctaaa 240
atgctatagt gcaattacta gcagttactg caccggagtg caccgtgcca atagaggact 300
gttggtttta taagggaact cttagcccat ttcttttaaa aaaaatcagt ttaattctta ctgtgtgccc 420
caatgaaata tcatttaaat ttcttttaaa aaaaatcagt ttaattctta ctgtgtgccc 480
aacacgaagg ctttttttga aagaaaaata gaatgttttg cctcaaagta gtccatataa 480
aatgtcttga atagaagaaa aaactaccaa accaaagggt actatthttg aaacatcgtg 540
tgttcattcc agcaaggcag aagactgcac cttctttcca gtgacatgct gtgtcatttt 600
ttttaagtc tcttaatttt tagacacatt ttgtgtttat gttttaacaa tgtatgccta 660
accagtcac ttgtctgcac caatgcgaaag gttcttgaga ggagtattct ctatccctgt 720
ggatataga acactggcat ttcatctatt ttccctttc ctttttaag gatttaactt 780
tggaatcttc caaaggaagt ttggccaatg ccagatcccc aggaatttgg ggggttttct 840
ttcttttcaa ctgaaattgt atctgattcc tactgttcat gttagtgatc atctaatac 900
agagccaaac acttttctcc cctgtgtgga aaagtaggta tgcttttcaa taaaatctgt 960
cttttctggt agaaacctga gccactgaaa ataaaagaga caactagaag cacagtagag 1020
tcccagactg agatctacct ttgagaggct ttgaaagtaa tccctggggg ttggattatt 1080
ttcacaaggg ttatgccgtt ttattcaagt ttgttgctcc gttttgcacc tctgcaataa 1140
aagcaaatg acaaccagta cataaggggt tagcttgaca aagtagactt ccttggtgta 1200
atthtttaagt ttttttttcc ttaactatat ctgtctacag gcagatacag atagttgtat 1260
gaaaatctgc ttgcctgtaa aatttgcatt tataaatgtg ttgccgatgg atcacttggg 1320
cctgtacaca taccaattag cgtgaccact tccatcttaa aaacaaacct aaaaaacaaa 1380
atthattata tatatatata tatatatata aaggactgtg ggttggtatac aaactattgc 1440
aaacacttgt gcaaatctgt cttgatataa aggaaaagca aaatctgtat aacattatta 1500
ctacttgaat gcctctgtga ctgatttttt ttttcatttt aaatataaac ttttttgtga 1560
aaagtatgct caatgttttt tttccctttc cccattccct tgtaaatata ttttgttcta 1620
tgtgacttgg ttggaataa gtttaactgg actgtaattt gcattaaata aaaagtaggt 1680
tagcctggaa atgaaattaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa 1738

```

<210> 2

<211> 84

<212> PRT

<213> Homo sapiens

&lt;400&gt; 2

Met Gln Arg Phe Leu Arg Gly Val Phe Ser Ile Pro Val Asp Met Lys  
 1 5 10 15

Thr Leu Ala Phe His Leu Phe Phe Pro Phe Leu Phe Lys Gly Phe Asn  
 20 25 30

Phe Gly Ile Phe Gln Arg Lys Phe Gly Gln Cys Gln Ile Pro Arg Asn  
 35 40 45

Leu Gly Gly Phe Leu Ser Phe Gln Leu Lys Leu Tyr Leu Ile Pro Thr  
 50 55 60

Val His Val Ser Asp His Leu Ile Thr Glu Pro Asn Thr Phe Leu Pro  
 65 70 75 80

Cys Val Glu Lys

&lt;210&gt; 3

&lt;211&gt; 2198

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

gctctgtgga ttctggccag gccgggttcg gcggttgctg tgagagcggg cttcccaaca 60  
 ccacgcccgc cgccttctct gtcagctctt tccccgtcag catcccagcc gtgctcacgc 120  
 agacgggactg gactgagccc tggctcatgg ggctggccac cttccacgcg ctctgcgtgc 180  
 tcctcacctg cttgtccctc cgaagctaca gactacagat cgggcacttt ctgtgtctag 240  
 tcactttagt ctactgtgct gaatacatca atgagggcgc tgccatgaac tggagattat 300  
 ttctgaaata ccagtatttc gactccaggg ggatgttcat ttctatagta ttttcagccc 360  
 cactgctggt gaatgccatg atcattgtgg ttatgtgggt atggaagact ttgaatgtga 420  
 tgactgacct gaagaatgca caagagagaa gaaaggaaaa gaaaaggaga aggaagaaga 480  
 actgaggggc agcagctgct tggagtttgc gtccctcccg tccaccagct gcagctccca 540  
 gtgctgcagt gtgcgtggcg tgggcatcct tccagctgac tcatggtttg aaaaaccgtt 600  
 gttttattta aatatccaca gtggtaggcg acacactgaa gttggctttt cagccagcac 660  
 tgaatgtatc catcaggaca tgcgtcttca ggtgcctgat cttttagtgc aggcgtgtgg 720  
 aacggtctct gcagagcttc ataactggga atttgatttg aagaagtcca tgtcatatgt 780  
 gtaactagta ctattataa atataaata cacaataata aatatgaaac tcaataataa 840  
 acagtgccac ctgtacatgg gcaccatgcc ctccctctcg tgctgtgttt tctagtgcac 900  
 gccacagttc gcagtagagg gtgttttcac ctccaagac atggggcaaa gtttgagac 960  
 acctggttgt cactggaggg ggtggtgctc ctggctctc ctgtggagcc cggggtgatg 1020  
 cataaaatcc tgtgtgcctg ggtcagccgc atcacagaca atgacttgac atgaaatgac 1080  
 agctgtgctg gggcagagag accttggaag gaagctcttg gaaaatacgt tgtatctcag 1140  
 tttgatgaac caattcaca gaggctaggg cctctctagc aaagttagtg gctgctttac 1200  
 tgaaaacaga atggaagccc tgaagtcaac actccatgga gaagcgtgtc tttcctaagt 1260  
 tcctggtgtt ctgttgattt aggtgcttgg gaacacaatg ctcccagttc tgtaggaca 1320  
 ggcatactgt tactttgcaa tatccacttt ataaaatagc tcctgccagc tggctcttgg 1380  
 ttctctgcaa atgtggacct gtagtttaag aatgacaggt ggtagagac ccagatattt 1440  
 aaaaataggt gttcaataag ggaatactga ttgtgcattg tatctggata gcacgcctaa 1500  
 ttgtgcattt ctgaaagtta ccaattcaaa atgtaattgg aacagttatc tttgattaga 1560  
 caagcctggg aagagaatgt tgaggtgcag agctcaccag ccaagttcat gccctctcgc 1620  
 ggcctttgtg gctgagaagt gggacagaaa gatgattaag gtaatgtgtc ctccctgtag 1680  
 cattgtccag ggcggtgtg tagatatttg acttcaactga cagaaaagaa accagggagt 1740  
 ttgttagagac tgtgcatttt tagtataaca ttttcacat ctgatattgt ttggctttgt 1800  
 gtccccaccc aaattgcatc tcaaatgtga atccccatgt gtcaaggag ggacctgatg 1860  
 ggaggtgatg gatcatggg ggtggtttcc cctatgttgt tatcataata gagagggagt 1920  
 tctcacaaga tctgctcgtt ttaaagacag cagtttcccc tgctgtcact gtctctctcc 1980  
 tgctgccttg tgaagaaggt gcttgtttct ccctctgcca tgattgtaag tttcccagag 2040

```

cctccccggc catgtggaac tgagtcaatt aaacttcttg tttataaagt aaaaaaaaaa 2100
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2160
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa                                     2198

```

<210> 4  
 <211> 140  
 <212> PRT  
 <213> Homo sapiens

```

<400> 4
Met Pro Ser Ala Phe Ser Val Ser Ser Phe Pro Val Ser Ile Pro Ala
  1             5             10             15

Val Leu Thr Gln Thr Asp Trp Thr Glu Pro Trp Leu Met Gly Leu Ala
      20             25             30

Thr Phe His Ala Leu Cys Val Leu Leu Thr Cys Leu Ser Ser Arg Ser
      35             40             45

Tyr Arg Leu Gln Ile Gly His Phe Leu Cys Leu Val Ile Leu Val Tyr
      50             55             60

Cys Ala Glu Tyr Ile Asn Glu Ala Ala Ala Met Asn Trp Arg Leu Phe
      65             70             75             80

Ser Lys Tyr Gln Tyr Phe Asp Ser Arg Gly Met Phe Ile Ser Ile Val
      85             90             95

Phe Ser Ala Pro Leu Leu Val Asn Ala Met Ile Ile Val Val Met Trp
      100            105            110

Val Trp Lys Thr Leu Asn Val Met Thr Asp Leu Lys Asn Ala Gln Glu
      115            120            125

Arg Arg Lys Glu Lys Lys Arg Arg Arg Lys Glu Asp
      130            135            140

```

<210> 5  
 <211> 2229  
 <212> DNA  
 <213> Homo sapiens

```

<400> 5
ggagccgggc aaccagtgga aggagcctgg gagaggccag gcctccccgg actgctagcc 60
tgcttttctt ggggtccctg gagccggagg aagaaccagg atgttgctgc ctgcagaagc 120
tcagctcagg aagacttcca ggaacctgag gaggagctgc cactaacagc catatttccc 180
aatggagact gtgatgacct tggaaggggg tcaaaagcct gtgatggagt cgtacacact 240
cctgctgagc ccaccggaga ctcaagatga aggctggacc cttgcgctgt ccctggctct 300
aacctacaga ctggggcctg gctccgtctt actggccccc aggtctccat ggagactgca 360
gaaacccccg cctgctggag gcctgccaca ctcacagtta ccagctagac agtggggcct 420
actaagacaa gcaggaccta aaacagtgtc tcccctggga acctactccc caccagcat 480
ttgctaagtc tgatcacagg gaggttattt tgtctctctg tctcggtttc tctgagccac 540
tgagacagat ggctgtccgc tttgaggctc tgcagagctg tggcacccca tgggtgtgtc 600
gcagtgttct gggcacatgc atgggcaccc atcggtgaga gtgcagctgg gaagaactct 660
gaaccagaag tcatcagagc tgaggcatgg ccttgaacat gtcactcagt ctctggggct 720
tctgtttcac aaatgcatga gggggccacc agcccagtgg ctttaaacca ggggcagggt 780
gtccctccag gcagcattgg aaatgtgtgt gtgttgaggg ggtcacagtg actgtggggg 840
caccctggc atctagtggg catcccacaa tgtgcagaac agtctctgac agcaaagaat 900
tgggtccattc aatgccaatt gtagtacctt tgagacattc tggctgagcc aatgccttct 960

```

```

ccctgtcaga gtccccaga gcagagaggg tcaggcttcc ctggaccttg gctcccagag 1020
caagccaaaa taaagactac actgttgccct tgggggcttg tcggggccagg gccaagacgg 1080
tctgctgtct gcagggccag gacagaaata gccacacatg ccggtgagaa caaagagcct 1140
ctttctttct catgttgaca tcgactttct gtgccaagtc ctttgggtat aaggatgcta 1200
gggaattcct ataggcacca aacagaagga aagctagggg cttggactac tgggtatagg 1260
acttgctcta gctctcaggt cctagcccaa gctcaatgca aacacagccc ctccgggctc 1320
tctgtttctg tgaggttctg gaatcccttc ctctgtgtcc gtgagtctga cagaatcgat 1380
gatgttcctt tagagctggg aaatccatgt gtttattcac ggagggaact caccattacc 1440
tcccttgctt tctttgcttg ccttggagaa atccagagtc ttcggaatgg caaaggcagc 1500
tcttggaatt ccctggaggg gaggcactag ctgagggaag tagctccctt cattcatgat 1560
gcacagttaa cgcagcagac acacaactgc gcctactatt tgctcggtgc cctgcaaggc 1620
gctgcctaac tttgatttgt tatttcagct ctctccagga tagtgccaaa tgggtgcaatg 1680
ggaaacctgt tttgctgggg ggctctagat cactggctcc agaactcccg gctgccaggg 1740
tagccctac cccagcccc ttgctcctgg acagcagtg gtctcacctt tagcctctgc 1800
ccccagttct ggtctgaccc aacagagggg ctctatgata ttaagaaggg gcccttctctg 1860
ctctgtgctt caacctattc tccataatag ggagtctaatt cctattcctt ccctgctga 1920
tgaggatggt gtgaggatga ggaggacggc atctcatttg gggctttttg gcagtggggc 1980
tcattttaat cctgcagggc tgctgcccag tggatctatc cagctgcttc cttgtagcca 2040
agaatgagtt caatgaattg tgattcactg attttattga ttttgtttta aaacagggag 2100
actggtattt ttgaagctgc tatcattttc tatttcttta ttaatttctt tgtaatcatc 2160
ttattaaagt tttcttattt agtggaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2220
aaaaaaaaa 2229

```

&lt;210&gt; 6

&lt;211&gt; 111

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

```

Met Leu Gly Asn Ser Tyr Arg His Gln Thr Glu Gly Lys Leu Ala
  1              5              10              15

Trp Thr Thr Gly Tyr Arg Thr Cys Ser Ser Ser Gln Val Leu Ala Gln
          20              25              30

Ala Gln Cys Lys His Ser Pro Ser Gly Leu Ser Val Ser Val Arg Phe
          35              40              45

Trp Asn Pro Phe Leu Cys Val Arg Glu Ser Asp Arg Ile Asp Asp Val
          50              55              60

Pro Leu Glu Leu Gly Asn Pro Cys Val Tyr Ser Arg Arg Glu Leu Thr
          65              70              75              80

Ile Thr Ser Leu Val Phe Phe Ala Cys Leu Gly Glu Ile Gln Ser Leu
          85              90              95

Arg Asn Gly Lys Gly Ser Ser Trp Ile Ser Leu Glu Gly Arg His
          100              105              110

```

&lt;210&gt; 7

&lt;211&gt; 915

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

```

gcgtgagcac cctatctcag ctgctattcg ggcattgatc tttgtagagg gtagagtaga 60
cagccccctc ccctactcac catgggtattt ctccctgaat tcctctttct tgttttcttt 120
cctgggtgtg tgaaccagtt gctgctgtca taccctggc agggccaggg gacctctctt 180

```



```

tggctcatttc tgccttttca ctggctgctg ccccaggaag actcctctag gctctccatc 240
tttcccttga gagctggctc cccaccccaa cctgctcagg caccaccgag gatctaggtc 300
tctggctccc catacctgga cccacatggg tgggtgcctg ttgcatgttt aagagagagg 360
ggctgtgagg tgacagggca ctagggcctt cactccttcc tccccttcca tcttttcttt 420
accagtgccca cccatgtccc tagctcccgg gtattggggc tgaggctctg gggcctgtct 480
ccctgccagc gtgagggcaa gaccccagag ccttagctga gcaagcccag aggggcagcg 540
tggcccttcc ctcccctttt cctgccccgt cccatgcctc agcttgcctg ttgtgccagt 600
tgctgttttc gcttcagtgt ttgattctag cacttacatg tgctctcccc accaagccct 660
ctatctcctt ctaatccttc aacccctggc cccctcccca taacagtgc ttttccaggg 720
aggaagaggc agcaggagct gttggccttg gtttgacag agcgggtagg gctgtaggga 780
aagcgggtga gctgttgtgc tgctgggctt cctttggcc ctcgcttccc accctacgat 840
gtatgaaatg tatgtacaga ccagagatgt ttatacagcc gataaagatg gaaaaaaaaa 900
aaaaaaaaaa aaaaaa 915

```

&lt;210&gt; 8

&lt;211&gt; 71

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

```

Met Val Phe Leu Leu Glu Phe Leu Phe Leu Val Phe Phe Pro Gly Cys
  1                      5                      10                     15

```

```

Val Asn Gln Leu Leu Leu Ser Tyr Pro Trp Gln Gly Gln Gly Thr Ser
      20                      25                     30

```

```

Leu Trp Ser Phe Leu Ser Phe His Trp Leu Leu Pro Gln Glu Asp Ser
    35                      40                     45

```

```

Ser Arg Leu Ser Ile Phe Pro Leu Arg Ala Gly Ser Pro Pro Gln Pro
    50                      55                     60

```

```

Ala Gln Ala Pro Pro Arg Ile
    65                      70

```

&lt;210&gt; 9

&lt;211&gt; 1100

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

```

tgacccccaa gacggaagag gatggccgcg gcggtcttga ggagattttg gtcccggcgc 60
cgcgagaggc cggggcgcgcg cgctgtgtat gtgggtctgc tgggcgggcgc ggcggcctgc 120
ttcacgctgg cggccagcga ggggtgccttc gaggaggcgc tgctggaggc gtcggggacc 180
ctcctgctgc tggcgccggc caccgcgaac cgcgggtcct ggacgtgggc ttcgtgggtc 240
gctgggtggg gctggggggc tggatgcgcg actgcgacat caacgacgac gaattcctgc 300
acctgccggc gcatttgcgg gtggtcgggc cccagcagct gcattccgag accaacgagc 360
ggctcttcga tgagaagtac aagcctgtcg tgctcaccga cgatcagggtg gaccaggcgc 420
tgtgggagga gcaggctctg cagaaggaga agaaggacag gctcgccctg agccaggccc 480
actcgctggt gcaggcggag gccccgagat gaaaccctga ggcccccgag tcctggcaaa 540
ctgcttgccct ggggtggtgc agttctgagt gtgcctcacc tgcagaacag ctgagacaga 600
tgatgtgcaa agtgttttct cactggattt gcacaagtgt ggggagcctt tctgcccccc 660
gtctttgttc tttattagct gaagctaatt cagagccacc tgggtccggg agttggggag 720
agcagaacga cttgacacat gttcatcact ggcagagctg gtcattgagcc ttttatataa 780
gcctttttca tcgggcctca gaggccctcc ttaaggaggt accacattgg tcagctgact 840
tgcaaaactct tctaaggcca cttagatttt ctttttcaag tttgggttgt ggctggcat 900
ggtgatgtct gtaatccagc cagtttgcaa actgaggcaa gagaatcgct tgaggccagg 960
agtttgaggc cagctagagt gacatagcaa gactccgatg ctacaaaaaa gaataataa 1020
agtaattaac taaataaaat ttgagctgaa atgttaaaaa aaaaaaaaaa aaaaaaaaaa 1080

```

aaaaaaaaaa aaaaaaaaaa

1100

&lt;210&gt; 10

&lt;211&gt; 149

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

Met Ala Ala Ala Leu Arg Arg Phe Trp Ser Arg Arg Arg Ala Glu  
 1 5 10 15

Ala Gly Arg Ala Ala Val Tyr Val Gly Leu Leu Gly Gly Ala Ala Ala  
 20 25 30

Cys Phe Thr Leu Ala Pro Ser Glu Gly Ala Phe Glu Glu Ala Leu Leu  
 35 40 45

Glu Ala Ser Gly Thr Leu Leu Leu Leu Ala Pro Ala Thr Arg Asn Arg  
 50 55 60

Gly Ser Trp Thr Trp Ala Ser Trp Val Ala Gly Gly Cys Trp Gly Pro  
 65 70 75 80

Gly Cys Ala Thr Ala Thr Ser Thr Thr Thr Asn Ser Cys Thr Cys Arg  
 85 90 95

Arg Ile Cys Gly Trp Ser Gly Pro Ser Ser Cys Ile Pro Arg Pro Thr  
 100 105 110

Ser Gly Ser Ser Met Arg Ser Thr Ser Leu Ser Cys Ser Pro Thr Ile  
 115 120 125

Arg Trp Thr Arg Arg Cys Gly Arg Ser Arg Ser Cys Arg Arg Arg Arg  
 130 135 140

Arg Thr Gly Ser Pro  
 145

&lt;210&gt; 11

&lt;211&gt; 2010

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

tgggtgagctg cagagaagag gaggttggtg tggagcacag gcagcaccga gcctgccccg 60  
 tgagctgagg gcctgcagtc tgcggctgga atcaggatag acaccaaggc aggaccccca 120  
 gagatgctga agcctctttg gaaagcagca gtggcccca catggccatg ctccatgccg 180  
 ccccgccgcc cgtgggacag agaggctggc acgttgacag tcctgggagc gctggctgtg 240  
 ctgtggctgg gctccgtggc tcttatctgc ctctgtggc aagtgccccg tctctccacc 300  
 tggggccagg tgcagcccaa ggacgtggc aggtcctggg agcatggctc cagcccagct 360  
 tgggagcccc tgaagcaga ggccaggcag cagagggact cctgccagct tgccttctg 420  
 gaaagcatcc ccaggacct gccatctgca gccggcagcc cctctgccca gcctctgggc 480  
 caggcctggc tgcagctgct ggacactgcc caggagagcg tccacgtggc ttcatactac 540  
 tggctccctca cagggcctga catcggggtc aacgactcgt cttcccagct gggagaggct 600  
 ctctctgaga agctgcagca gctgctgggc aggaacattt ccctggctgt ggccaccagc 660  
 agccccacac tggccaggac atccaccgac ctgcagggtc tggctgcccc aggtgcccac 720  
 gtacgacagg tgcccctggg gcggctcacc aggggtgttt tgactccaa attctgggtt 780  
 gtggatggac ggcacatata catgggcagt gccaacatgg actggcggtc tctgacgcag 840  
 gtgaaggagc ttggcgctgt catctataac tgcagccacc tggcccaaga cctggagaag 900

```

accttccaga cctactgggt actgggggtg cccaaggctg tcctcccca aacctggcct 960
cagaacttct catctcactt caaccgtttc cagcccttcc acggcctctt tgatgggggtg 1020
cccaccactg cctacttctc agcgctcgcca ccagcactct gtccccaggg ccgcaccccg 1080
gacctggagg cgctgctggc ggtgatgggg agcgcccagg agttcatcta tgctcccggtg 1140
atggagtatt tccccaccac gcgcttcagc cacccccga ggtactggcc ggtgctggac 1200
aacgcgctgc gggcggcagc cttcggaag ggcgtgcgcg tgcgcctgct ggtcggctgc 1260
ggactcaaca cggacccac catgttcccc tacctgcggt ccctgcaggc gctcagcaac 1320
cccgcggcca acgtctctgt ggacgtgaaa gtcttcatcg tgccgggtggg gaaccattcc 1380
aacatcccat tcagcagggt gaaccacagc aagttcatgg tcacggagaa ggcagcctac 1440
ataggcacct ccaactggtc ggaggattac ttcagcagca cggcgggggt gggcttggtg 1500
gtcaccaga gccctggcgc gcagcccgcg ggggccacgg tgcaggagca gctcggcag 1560
ctctttgagc gggactggag ttcgcgctac gccgtcgcc tggaaggaca ggtccgggc 1620
caggactcg tttggcaggg ctgagggggg cctctttttc tctcggcgac cccgccccgc 1680
acgcgcctc cctctgacc cgggcctggg cttcagccgc ttcctcccg aagcagcccg 1740
ggtccgcact gcgccaggag ccgctgcga ccgcccgggc gtcgcaaacc gccgcctgc 1800
tctctgattt ccgagtcag cccccctga gcccacctc ctccaggag ccctccagga 1860
agcccccttc ctgactcctg gccacaggc caggcctaaa aaaaactcgt ggcttaaaaa 1920
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1980
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2010

```

<210> 12

<211> 506

<212> PRT

<213> Homo sapiens

<400> 12

```

Met Leu Lys Pro Leu Trp Lys Ala Ala Val Ala Pro Thr Trp Pro Cys
  1             5             10             15

Ser Met Pro Pro Arg Arg Pro Trp Asp Arg Glu Ala Gly Thr Leu Gln
      20             25             30

Val Leu Gly Ala Leu Ala Val Leu Trp Leu Gly Ser Val Ala Leu Ile
      35             40             45

Cys Leu Leu Trp Gln Val Pro Arg Pro Pro Thr Trp Gly Gln Val Gln
      50             55             60

Pro Lys Asp Val Pro Arg Ser Trp Glu His Gly Ser Ser Pro Ala Trp
      65             70             75             80

Glu Pro Leu Glu Ala Glu Ala Arg Gln Gln Arg Asp Ser Cys Gln Leu
      85             90             95

Val Leu Val Glu Ser Ile Pro Gln Asp Leu Pro Ser Ala Ala Gly Ser
      100            105            110

Pro Ser Ala Gln Pro Leu Gly Gln Ala Trp Leu Gln Leu Leu Asp Thr
      115            120            125

Ala Gln Glu Ser Val His Val Ala Ser Tyr Tyr Trp Ser Leu Thr Gly
      130            135            140

Pro Asp Ile Gly Val Asn Asp Ser Ser Ser Gln Leu Gly Glu Ala Leu
      145            150            155            160

Leu Gln Lys Leu Gln Gln Leu Leu Gly Arg Asn Ile Ser Leu Ala Val
      165            170            175

Ala Thr Ser Ser Pro Thr Leu Ala Arg Thr Ser Thr Asp Leu Gln Val

```

180					185					190						
Leu	Ala	Ala	Arg	Gly	Ala	His	Val	Arg	Gln	Val	Pro	Met	Gly	Arg	Leu	
195					200					205						
Thr	Arg	Gly	Val	Leu	His	Ser	Lys	Phe	Trp	Val	Val	Asp	Gly	Arg	His	
210					215					220						
Ile	Tyr	Met	Gly	Ser	Ala	Asn	Met	Asp	Trp	Arg	Ser	Leu	Thr	Gln	Val	
225					230					235					240	
Lys	Glu	Leu	Gly	Ala	Val	Ile	Tyr	Asn	Cys	Ser	His	Leu	Ala	Gln	Asp	
245					250					255						
Leu	Glu	Lys	Thr	Phe	Gln	Thr	Tyr	Trp	Val	Leu	Gly	Val	Pro	Lys	Ala	
260					265					270						
Val	Leu	Pro	Lys	Thr	Trp	Pro	Gln	Asn	Phe	Ser	Ser	His	Phe	Asn	Arg	
275					280					285						
Phe	Gln	Pro	Phe	His	Gly	Leu	Phe	Asp	Gly	Val	Pro	Thr	Thr	Ala	Tyr	
290					295					300						
Phe	Ser	Ala	Ser	Pro	Pro	Ala	Leu	Cys	Pro	Gln	Gly	Arg	Thr	Arg	Asp	
305					310					315					320	
Leu	Glu	Ala	Leu	Leu	Ala	Val	Met	Gly	Ser	Ala	Gln	Glu	Phe	Ile	Tyr	
325					330					335						
Ala	Ser	Val	Met	Glu	Tyr	Phe	Pro	Thr	Thr	Arg	Phe	Ser	His	Pro	Pro	
340					345					350						
Arg	Tyr	Trp	Pro	Val	Leu	Asp	Asn	Ala	Leu	Arg	Ala	Ala	Ala	Phe	Gly	
355					360					365						
Lys	Gly	Val	Arg	Val	Arg	Leu	Leu	Val	Gly	Cys	Gly	Leu	Asn	Thr	Asp	
370					375					380						
Pro	Thr	Met	Phe	Pro	Tyr	Leu	Arg	Ser	Leu	Gln	Ala	Leu	Ser	Asn	Pro	
385					390					395					400	
Ala	Ala	Asn	Val	Ser	Val	Asp	Val	Lys	Val	Phe	Ile	Val	Pro	Val	Gly	
405					410					415						
Asn	His	Ser	Asn	Ile	Pro	Phe	Ser	Arg	Val	Asn	His	Ser	Lys	Phe	Met	
420					425					430						
Val	Thr	Glu	Lys	Ala	Ala	Tyr	Ile	Gly	Thr	Ser	Asn	Trp	Ser	Glu	Asp	
435					440					445						
Tyr	Phe	Ser	Ser	Thr	Ala	Gly	Val	Gly	Leu	Val	Val	Thr	Gln	Ser	Pro	
450					455					460						
Gly	Ala	Gln	Pro	Ala	Gly	Ala	Thr	Val	Gln	Glu	Gln	Leu	Arg	Gln	Leu	
465					470					475					480	
Phe	Glu	Arg	Asp	Trp	Ser	Ser	Arg	Tyr	Ala	Val	Gly	Leu	Asp	Gly	Gln	
485					490					495						
Ala Pro Gly Gln Asp Cys Val Trp Gln Gly																

500

505

<210> 13  
 <211> 2830  
 <212> DNA  
 <213> Homo sapiens

&lt;400&gt; 13

```

ggcagaggca tgcctcatggt gctccttccc cagcagcaat cctatgctcc ccttccaggt 60
cttctcttcc ttcttttccct cagccagctg gctgctggta agagaggtct ggcgagagtg 120
tgtcccagta gctgctgcta tggatggaac ggagcgggca ggacgtaggc tggaggagtgc 180
tgtcttcccc gcatctttac gagatcgctg ctgcaggacc ttaatcatag catctttctc 240
tatgattttg gcatggagat ttttaatagt gtattccatg tcctgacacc ttctgttggc 300
ctgcaccacc tcctcctcct ctgcccagat gtgggcctcc agcgagctct ctccgtagct 360
gccattccgt gagtggttga ttgtcgtggt gtcctctctca gctgctgcag tggctgcggc 420
attcatggca aagtgtcggg tgggtgctctc ctccaggtagc ttctgctccc actttgtcat 480
gtcggcctcc agggccagga tccgctcctc ctctctccgc acaagttcca ggagggtctg 540
ggcattgtat tccggcatgt tggctggctg gccatttcca tgtttctgct gggttctcag 600
tgcattccagc tctctctcca gccaaagtcg cagtctccgc tccatctgtt ctgccttctc 660
acatgcagac tgcagctggg tcaggccctg ctgcagcttc tcaactttct caacatagc 720
ttgcttctct cgtaactcct ctccagctt gatgacctg gctgggctg tgcctaaagc 780
ctgggtccagg atctcgatgt gtctccgatg gtctcactt gcagtcgca ctgctgctaa 840
ctccatttct aatttctcct ttctcttcaa gaattctttg ttctgggaag cataatgcc 900
ctctgcagct ttgtcttcat gcccttcgta ttccctgctg gatagttgcc tgttagcagt 960
ctctagtcca tctcggaggt ctctgttgaa atcatgaagt ctctaatct cgccttccaa 1020
tttgtttctc atggccttgt ccagcgattc tcgcttgggt gtagacttga ccagacttcc 1080
ataggcttcc gaaattctct gaagtctttt ttcaaacttg tggagcttgt cggcattgtc 1140
gtagtacccc tgaagttcct ggtgaagcac ccggttctcc tctgttaata tctccaccat 1200
ttgctgggct cgctccacaa tcgcaaaggc atctggacca agctgctggc tgggggaggc 1260
ggcagggcgg ggctgtggag caccagggc catcgggagt ggaagcggca gggagacaga 1320
gtgcagtgcc ccgctggcgg aagaggtctg ggaggacatg ggcctgtgct gctgcatggt 1380
tgatgggaac ggaagtggc atgggaggct cgttacccca tactcagggg gtggctggta 1440
ccgcaggacg gccacatctg ttctcacagg ctgaggagca gggtagggct tgaccatctc 1500
gtggagcatc ccgggtgct ggtcaccata aaaaagtcg tgcctcctgg tcttgctgac 1560
tggggacatc atttgcttgg tcttgaagggt gtactcaggt ggaggacccc gagggtcag 1620
cactttacct gcaggctggg caggggaggg ccccccctct actttgaagc cctttccatt 1680
ccccgagccg ggaaggtgtt gcttggcccc attcctctcc agggacagct gcatgattct 1740
ctcgctgagc gagcggacgt ggccctgctt cagttccttc agcgctcctg ccttatgcgc 1800
ctgtccactg ttggcacggt tcacagtggt cctcccctca gttcgggact tctgactggt 1860
gccccctgcc atgtagtaac catggccac cgcctctgct tgcgttggct gctgctgctg 1920
ccccctgaag aactgcgact gtgctttggc ctctcgttaa gtgggcagtt cctcgttgtt 1980
ctgctgaggc tgcgtggacc ggacctgttt ctccatcacc gtattgtcca cctggtgttc 2040
ttgacctgct ggttcttggc gtgctgactg gtagaccatt tgtgggtctt cttgagttag 2100
gttttccgct gaagaaaagt tgtttgtagg atgggctggt cctgcactcc ctgtggcctg 2160
gtgctgaatg gccagcaagt tcatgttctc ggttgggggt ccataccgca gttgttccct 2220
gatcagccgc tgcaatactg ttccagctgc cgcactctcg gaacctctca ttccacctc 2280
tgagatcttc aggaagtgtt gggagtggaa gttacaaaga gtagcttcaa caacttttcc 2340
cctgatgcta ctggttgctt ccaccgtcaa agcagatgtt tcatgcctcc cagaatagag 2400
ctgaaagtct ggggaaaagt aggtggagtc ttctagaacc tggacaggac tactggggct 2460
ataacaagca gaaggggacc ctaactcgtc tgggctgtag gatgcgggct aggcttccac 2520
aaactcactg tctgggggag aagaaaagca gaaaacaact cgaatcgcta ccattcagga 2580
cgaacccgcc aagcaccagc tcaagcgag gtccccggga aaagcggggg ctctctcttc 2640
ccagcgctca gaatccctga gccggaggcc ccgcgggatt cagaccgcca gatccccagg 2700
gagtgcacaa tcgcccagca aacttggggg acaactcggc cctggcaccg cgcggcttcc 2760
aggcgcggtc aggcgcgcgc caactttccc cgcgtgccac cccgcggctc ccccgccgg 2820
gcgctgggccc
2830

```

<210> 14  
 <211> 171

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

Met Ile Val Val Ser Leu Ser Ala Ala Val Ala Ala Phe Met  
 1 5 10 15

Ala Lys Cys Arg Met Val Leu Ser Ser Arg Tyr Phe Cys Ser His Phe  
 20 25 30

Val Met Ser Ala Ser Arg Ala Arg Ile Arg Ser Ser Phe Ser Arg Thr  
 35 40 45

Ser Ser Arg Arg Ala Gly Ala Leu Tyr Ser Gly Met Leu Ala Gly Trp  
 50 55 60

Pro Phe Pro Cys Phe Cys Trp Val Leu Ser Ala Ser Ser Ser Leu Ser  
 65 70 75 80

Ser Gln Val Arg Ser Leu Arg Ser Ile Cys Ser Arg Phe Ser His Ala  
 85 90 95

Asp Cys Ser Trp Val Arg Ala Cys Cys Ser Phe Ser Thr Phe Ser Thr  
 100 105 110

Tyr Ala Cys Phe Ser Arg Asn Ser Ser Ser Ser Leu Met Thr Leu Ala  
 115 120 125

Trp Ala Leu Leu Lys Ala Trp Ser Arg Ile Ser Met Cys Leu Arg Trp  
 130 135 140

Ser Ser Leu Ala Val Arg Thr Ala Ala Asn Ser Ile Ser Asn Phe Ser  
 145 150 155 160

Phe Ser Phe Lys Asn Ser Leu Phe Trp Glu Ala  
 165 170

&lt;210&gt; 15

&lt;211&gt; 2000

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

gcagaagctg tacgggctcc aggctgacat taccatcagc ctggacggcg tgcccttctc 60  
 catgcatgac accaccctgc ggcgcaccac caacgtggag gaggagtcc cggagctggc 120  
 ccgcaggcct gcctccatgc ttaactggac caccctgcag agactcaacg ctggccagtg 180  
 gttcctgaag actgaccctc tctggacagc cagctccctg tcaccctccg accacagaga 240  
 ggcccagaac cagtccatct gcagcctggc agagctcctg gagctggcca agggcaatgc 300  
 cacactgctg ctcaacctgc gtgacccgcc ccgggagcac ccctaccgca gcagttttat 360  
 caacgtgact ctggaggccg tgctgcactc cggttcccc cagcaccagg tcatgtggct 420  
 gcctagcagg cagaggcccc tggcgcgaa ggtggctccc ggcttccaac agacatcagg 480  
 ctccaaggag gcagtcgcca gcctgcggag aggccacatc cagcggctga acctgcgcta 540  
 cactcaggtg tcccgcagg agctcaggga ctacgcgtcc tggaacctga gtgtgaacct 600  
 ctacacagtc aacgcaccgt ggctcttctc cctgctgtgg tgtgcggggg tcccatccgt 660  
 cacctctgac aactcccaca ccctgtccca ggtgccttcc cccctctgga tcatgcccc 720  
 ggacgagtac tgtctcatgt ggtcactgc cgacctgtc tccttcaccc tcatcgtggg 780  
 catcttcgtg ctccagaagt ggcgcctggg tggcatacgg agctacaacc ctgagcagat 840  
 catgctgagt gctgcggtgc gccggaccag ccgggacgtc agcatcatga aggagaagct 900  
 tattttctca gagatcagcg atggtgtaga ggtctccgat gtgctctccg tatgttcaga 960

```

caacagttat gacacatatg ccaacagcac cgccaccctt gtgggcccc gagggggtgg 1020
cagccacacc aagaccctca tagagcggag tgggcgtag ctgaagacat gtctgtccca 1080
cctgtacctg acacagaagc tggggagcct aggagagctg gtggaagtgt gtctgaactc 1140
ggagtgtctt gggagcgggc tccacagcct ccttgtgggc tccagccctt tgcagccgc 1200
agcctctctt gagggggact ccctgtcccc tgaggcccag ctgggccagg actccatcct 1260
ttcagatgcc cctgcaggcc tggggctcct tctgggaagt atggggccta gggcttggtc 1320
cccctcttct gaggcctctt cctgtatccc gacctggaag ctttgatggg tcatgggcca 1380
tgccataccc cctgtggcaa tggagtgtgt ggatgtcac ctgtgccatc tgcctcctg 1440
tctgtgccag gaggcacctg agttctctgc tgttatcctg cccaagggc ctgggccgag 1500
cctctacctg aagcaactct gctcttctg tcagtctcaa agcacaagga ggttcagccc 1560
aggaggaagc cagctgcaat gtggagacac gtctcctcc ccaaccacc tcatgccacc 1620
gccaaccccc tgccccagga gggggcctga gccacgtccc ctaggagcag ctggagatgg 1680
ccaaaagagt gagctcaagg actactggga tccaatgcc agtgtccagc agacctcaag 1740
gcagaagggc cacctaaccc aggagtccac agactgatgt gacctcaggt tcccacatca 1800
gtggccacag ggcaggggcc acctggtaga agtgttctg atatggccag ggtgggtgtg 1860
tggctaagtg ggcctgaaca gagggaaact agggcccttg gcccaatgtg attaaaggct 1920
gccatcttga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1980
aaaaaaaaa aaaaaaaaaa                                2000

```

&lt;210&gt; 16

&lt;211&gt; 332

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

```

Met His Asp Thr Thr Leu Arg Arg Thr Thr Asn Val Glu Glu Glu Phe
  1              5              10              15

Pro Glu Leu Ala Arg Arg Pro Ala Ser Met Leu Asn Trp Thr Thr Leu
      20              25              30

Gln Arg Leu Asn Ala Gly Gln Trp Phe Leu Lys Thr Asp Pro Phe Trp
      35              40              45

Thr Ala Ser Ser Leu Ser Pro Ser Asp His Arg Glu Ala Gln Asn Gln
      50              55              60

Ser Ile Cys Ser Leu Ala Glu Leu Leu Glu Leu Ala Lys Gly Asn Ala
      65              70              75              80

Thr Leu Leu Leu Asn Leu Arg Asp Pro Pro Arg Glu His Pro Tyr Arg
      85              90              95

Ser Ser Phe Ile Asn Val Thr Leu Glu Ala Val Leu His Ser Gly Phe
      100              105              110

Pro Gln His Gln Val Met Trp Leu Pro Ser Arg Gln Arg Pro Leu Val
      115              120              125

Arg Lys Val Ala Pro Gly Phe Gln Gln Thr Ser Gly Ser Lys Glu Ala
      130              135              140

Val Ala Ser Leu Arg Arg Gly His Ile Gln Arg Leu Asn Leu Arg Tyr
      145              150              155              160

Thr Gln Val Ser Arg Gln Glu Leu Arg Asp Tyr Ala Ser Trp Asn Leu
      165              170              175

Ser Val Asn Leu Tyr Thr Val Asn Ala Pro Trp Leu Phe Ser Leu Leu
      180              185              190

```

Trp Cys Ala Gly Val Pro Ser Val Thr Ser Asp Asn Ser His Thr Leu  
 195 200 205  
 Ser Gln Val Pro Ser Pro Leu Trp Ile Met Pro Pro Asp Glu Tyr Cys  
 210 215 220  
 Leu Met Trp Val Thr Ala Asp Leu Val Ser Phe Thr Leu Ile Val Gly  
 225 230 235 240  
 Ile Phe Val Leu Gln Lys Trp Arg Leu Gly Gly Ile Arg Ser Tyr Asn  
 245 250 255  
 Pro Glu Gln Ile Met Leu Ser Ala Ala Val Arg Arg Thr Ser Arg Asp  
 260 265 270  
 Val Ser Ile Met Lys Glu Lys Leu Ile Phe Ser Glu Ile Ser Asp Gly  
 275 280 285  
 Val Glu Val Ser Asp Val Leu Ser Val Cys Ser Asp Asn Ser Tyr Asp  
 290 295 300  
 Thr Tyr Ala Asn Ser Thr Ala Thr Pro Val Gly Pro Arg Gly Gly Gly  
 305 310 315 320  
 Ser His Thr Lys Thr Leu Ile Glu Arg Ser Gly Arg  
 325 330

&lt;210&gt; 17

&lt;211&gt; 1170

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

```

aagtttggtc cccgagttcg gagcctagga gccccccgcg gctgcggcgc aggtgccctc 60
ggcctgagtc gggatggagc tgcttgctgt gaacctgaag gtgattctcc taggtcactg 120
gctgctgaca acctggggct gcattgtatt ctcaggetcc tatgcctggg ccaacttcac 180
catcctggcc ttgggcgtgt gggctgtggc tcagcgggac tccatcgacg ccataagcat 240
gtttctgggt ggcttgctgg ccaccatctt cctggacatc gtgcacatca gcattcttcta 300
cccgcgggtc agcctcacgg acacggggccg ctttggcgtg ggcattggcca tcctcagctt 360
gctgctcaag ccgctctcct gctgcttcgt ctaccacatg taccgggagc gcgggggttt 420
ccttgggtct tctcaggacc gtagtgccta ccagacgatt gactcagcag aggcgcccgc 480
agatcccttt gcagtcccag agggcaggag tcaagatgcc cgagggtact gaagccagcc 540
acgtgcgcc cgccctgcc ccgggccttc ctcgtgcctg ggaggtcggt ctagggatgc 600
tcttgacctc cgtctcttgg acctaatgat gaatgtgtcc ccagctcagg gattgcctga 660
accaagaggg caggagcccc catgggcccgc ccagtaccat gcacactcct gtcccgaact 720
ccctgaggcc tccccctcct tcagggcacc cactggttcc caggctggaa ccagggtctc 780
tctttacctc ctaccccatg gtggcaccac agaggccctc agccgagtcc tgcttgagt 840
ttgcaagctc aggcctttaa ggactgtga tgccccctca ggccctcccc aagtttgctg 900
ggctttggtg gaagccctga gagcttcagg tcctgctcag cccgaggagc agtctgacat 960
gggagtgagg ccctgtcctt ctactgcctt ggtcacatgg tgcctaggga tgcagggtg 1020
gaggccagag gtgtcagcaa cactgtgacc caccacaacc tccagcctcc cttttcagag 1080
cacagcatta aagtttgggg aattctgtaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa

```

1170

&lt;210&gt; 18

&lt;211&gt; 152

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens



&lt;400&gt; 18

Met Glu Leu Pro Ala Val Asn Leu Lys Val Ile Leu Leu Gly His Trp  
 1 5 10 15

Leu Leu Thr Thr Trp Gly Cys Ile Val Phe Ser Gly Ser Tyr Ala Trp  
 20 25 30

Ala Asn Phe Thr Ile Leu Ala Leu Gly Val Trp Ala Val Ala Gln Arg  
 35 40 45

Asp Ser Ile Asp Ala Ile Ser Met Phe Leu Gly Gly Leu Leu Ala Thr  
 50 55 60

Ile Phe Leu Asp Ile Val His Ile Ser Ile Phe Tyr Pro Arg Val Ser  
 65 70 75 80

Leu Thr Asp Thr Gly Arg Phe Gly Val Gly Met Ala Ile Leu Ser Leu  
 85 90 95

Leu Leu Lys Pro Leu Ser Cys Cys Phe Val Tyr His Met Tyr Arg Glu  
 100 105 110

Arg Gly Gly Phe Leu Gly Ser Ser Gln Asp Arg Ser Ala Tyr Gln Thr  
 115 120 125

Ile Asp Ser Ala Glu Ala Pro Ala Asp Pro Phe Ala Val Pro Glu Gly  
 130 135 140

Arg Ser Gln Asp Ala Arg Gly Tyr  
 145 150

&lt;210&gt; 19

&lt;211&gt; 4144

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

ctccttgggc aggtgaggat aacagccctg tggcgagctg gtcctgagca tctgagttag 60  
 cagtgatgcg gagaaatgac cttggtggga ggggacgtcc acagacgtga cccggacgtc 120  
 gggacagggt gatctggggt caaggggagt gtttagaata cttgttggca tccatgacgc 180  
 agtgggtgtga gcgtgagata tccagcatcg cacctggaga gctgtgctgt ctgctcctt 240  
 ccttcctgcc tcaggaggag tgcagaact acgtgagagt cctgatcgtc gccggccgga 300  
 aggtgttcat gtgtggaacc aatgcctttt ccccatgtg caccagcaga cagggtggga 360  
 acctcagccg gactattgag aagatcaatg gtgtggcccg ctgcccctat gaccacgcc 420  
 acaactccac agctgtcatc tcctcccagg gggagctcta tgcagccacg gtcacgcact 480  
 tctcaggtcg ggacctgcc atctaccgca gcctgggcag tgggccaccg ctctgcactg 540  
 cccaatataa ctccaagtgg cttaatgagc caaacttcgt ggcagcctat gatattgggc 600  
 tgtttgcata cttcttctcg cgggagaacg cagtggagca cgactgtgga cgcaccgtgt 660  
 actctcgctg ggcgcgctg tgcaagaatg acgtgggggg ccgattcctg ctggaggaca 720  
 catggaccac attcatgaag gcccggtcca actgtccccg cccgggagag gtccccttct 780  
 actataacga gctgcagagt gccttccact tgccggagca ggacctcatc tatggagttt 840  
 tcacaacca cgtaaacagc attgcggctt ctgctgtctg cgccttcaac ctgagtgcta 900  
 tctcccaggc tttcaatggc ccatttcgct accaggagaa ccccgagggt gcctgggtcc 960  
 ccatagccaa ccccatcccc aatttccagt gtggcaccct gcctgagacc ggtcccaacg 1020  
 agaacctgac ggagcgagc ctgcaggacg cgcagcgctt cttcctgatg agcgaggccg 1080  
 tgcagccggt gacacccgag ccctgtgtca cccaggacag cgtgcgcttc tcacacctcg 1140  
 tgggtggacct ggtgcaggct aaagacacgc tctaccatgt actctacatt ggcaccgagt 1200  
 cgggcacccat cctgaaggcg ctgtccacgg cgagccgcag cctccacggc tgctacctgg 1260

```

aggagctgca cgtgctgccc cccggggcgc gcgagccctc gcgcagcctg cgcacccctgc 1320
acagcgcccc cgcgctcttc gtggggctga gagacggcgt cctgcgggtc ccactggaga 1380
ggtgcgcccc ctaccgcagc cagggggcat gcctgggggc cggggacccg tactgtggct 1440
gggacgggaa gcagcaacgt tgcagcacac tcgaggacag ctccaacatg agcctctgga 1500
cccagaacat caccgcctgt cctgtgcgga atgtgacacg ggatgggggc ttcggcccat 1560
ggtcaccatg gcaacatgt gagcacttgg atggggacaa ctcaggctct tgccctgtgc 1620
gagctcgatc ctgtgattcc cctcgacccc gctgtggggg ccttgactgc ctggggccag 1680
ccatccacat cgccaactgc tccaggaatg gggcgtggac cccgtggtca tcgtgggcgc 1740
tgtgcagcac gtccctgtggc atcggtctcc aggtccgcca gcgaagtgc agcaaccctg 1800
ctccccgcca cgggggcccgc atctgcgtgg gcaagagccg ggaggaacgg ttctgtaatg 1860
agaacacgcc ttgcccggtg cccatcttct gggcttcctg gggctcctgg agcaagtgc 1920
gcagcaactg tggagggggc atgcagtcgc ggcgtcgggc ctgcgagaac ggcaactcct 1980
gcctgggctg cggcgtggag ttcaagacgt gcaacccgga gggctgcccc gaagtgcggc 2040
gcaacacccc ctggagcgcg tggctgcccg tgaacgtgac gcaggcgagg gcacggcagg 2100
agcagcgggt ccgcttcacc tgccgcgcgc ccttgacaga cccgcacggc ctgcagttcg 2160
gcaggagaaag gaccgagacg aggaacctgc ccgcggacgg ctccggctcc tgcgacaccg 2220
acgcccctgt ggaggacctc ctgcgcagcg ggagcacctc cccgcacacg gtgagcgggg 2280
gctggggccc ctggggcccgc tggctgctct gctcccggga ctgcgagctg ggcttcgcgc 2340
tccgcaagag aacgtgcact aacccggagc cccgcaacgg gggcctgccc tgcgtgggcg 2400
atgtgcgca gtaccaggac tgcaaccccc aggcctgccc agttcggggg gcttggtcct 2460
gctggacctc atggtctcca tgctcagctt cctgtggtgg gggtcactat caacgcaccc 2520
gttctctgac cagccccgca cctccccag gtgaggacat ctgtctcggg ctgcacacgg 2580
aggaggcact atgtgccaca caggcctgcc cagaaggctg gtcgcccctg tctgagtggg 2640
gtaagtgcac tgacgacgga gcccagagcc gaagccggca ctgtgaggag ctctctccag 2700
ggtccagcgc ctgtgctgga aacagcagcc agagccgccc ctgcccctac agcgagattc 2760
ccgtcatcct gccagcctcc agcatggagg aggccaccgg ctgtgcaggg ttcaatctca 2820
tccacttggg gccacgggc atctcctgct tcttgggctc tgggctcctg accctagcag 2880
tgtacctgtc ttgccagcac tgccagcgtc agtcccagga gtccacactg gtccatcctg 2940
ccaccccaaa ccatttgac tacaaggcg gaggcacccc gaagaatgaa aagtacacac 3000
ccatggaatt caagaccctg aacaagaata acttgatccc tgatgacaga gccaaactct 3060
accatttgca gcagaccaat gtgtacacga ctacttacta cccaagcccc ctgaacaaac 3120
acagcttccg gccgaggcc tcacctggac aacggtgctt cccaacacg tgataccgcc 3180
gtcctgggga cttgggcttc ttgccttcat aaggcacaga gcagatggag atgggacagt 3240
ggagccagtt tggttttctc cctctgcact aggccaaagaa cttgctgcct tgccctgtggg 3300
gggtcccatc cggcttcaga gagctctggc tggcattgac catgggggaa agggctggtt 3360
tcaggctgac atatggccgc aggtccagtt cagcccaggt ctctcatggt tatcttccaa 3420
ccactgtca cgtgacact atgctgcat gcctgggctg tggacctact gggcatttga 3480
ggaattggag aatggagatg gcaagagggc aggcctttaa gtttgggtg gagacaactt 3540
cctgtggccc ccacaagctg agtctggcct tctccagctg gccccaaaaa aggcctttgc 3600
tacatcctga ttatctctga aagtaataaa tcaagtggct ccagtagctc tggattttct 3660
gccagggctg ggccattgtg gtgctgcccc agtatgacat gggaccaagg ccagcgacag 3720
ttatccacct ctgectggaa gtctatactc taccagggc atccctctgg tcagaggcag 3780
tgagtactgg gaactggagg ctgacctgtg cttagaagtc ctttaactct ggctggtaca 3840
ggcctcagcc ttgcccctaa tgcacgaaag gtggccagg agagaggatc aatgccatag 3900
gaggcagaag tctggcctct gtgcctctat ggagactatc ttccagttgc tgctcaacag 3960
agtgtgtggc tgagacctgc ttgggagtct ctgctggccc ttcatctgtt caggaacaca 4020
cacacacaca cactcacaca cgcacacaca atcacaattt gctacagcaa caaaaaagac 4080
attgggctgt ggcattatta attaaagatg atatccagtc aaaaaaaaaa aaaaaaaaaa 4140
aaaaa 4144

```

&lt;210&gt; 20

&lt;211&gt; 999

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

```

Met Thr Gln Trp Cys Glu Arg Glu Ile Ser Ser Ile Ala Pro Gly Glu
  1             5             10             15

```

```

Leu Cys Cys Leu Leu Leu Ser Phe Leu Pro Gln Glu Glu Cys Gln Asn

```

20	25	30
Tyr Val Arg Val Leu Ile Val Ala Gly Arg Lys Val Phe Met Cys Gly 35 40 45		
Thr Asn Ala Phe Ser Pro Met Cys Thr Ser Arg Gln Val Gly Asn Leu 50 55 60		
Ser Arg Thr Ile Glu Lys Ile Asn Gly Val Ala Arg Cys Pro Tyr Asp 65 70 75 80		
Pro Arg His Asn Ser Thr Ala Val Ile Ser Ser Gln Gly Glu Leu Tyr 85 90 95		
Ala Ala Thr Val Ile Asp Phe Ser Gly Arg Asp Pro Ala Ile Tyr Arg 100 105 110		
Ser Leu Gly Ser Gly Pro Pro Leu Arg Thr Ala Gln Tyr Asn Ser Lys 115 120 125		
Trp Leu Asn Glu Pro Asn Phe Val Ala Ala Tyr Asp Ile Gly Leu Phe 130 135 140		
Ala Tyr Phe Phe Leu Arg Glu Asn Ala Val Glu His Asp Cys Gly Arg 145 150 155 160		
Thr Val Tyr Ser Arg Val Ala Arg Val Cys Lys Asn Asp Val Gly Gly 165 170 175		
Arg Phe Leu Leu Glu Asp Thr Trp Thr Thr Phe Met Lys Ala Arg Leu 180 185 190		
Asn Cys Ser Arg Pro Gly Glu Val Pro Phe Tyr Tyr Asn Glu Leu Gln 195 200 205		
Ser Ala Phe His Leu Pro Glu Gln Asp Leu Ile Tyr Gly Val Phe Thr 210 215 220		
Thr Asn Val Asn Ser Ile Ala Ala Ser Ala Val Cys Ala Phe Asn Leu 225 230 235 240		
Ser Ala Ile Ser Gln Ala Phe Asn Gly Pro Phe Arg Tyr Gln Glu Asn 245 250 255		
Pro Arg Ala Ala Trp Leu Pro Ile Ala Asn Pro Ile Pro Asn Phe Gln 260 265 270		
Cys Gly Thr Leu Pro Glu Thr Gly Pro Asn Glu Asn Leu Thr Glu Arg 275 280 285		
Ser Leu Gln Asp Ala Gln Arg Leu Phe Leu Met Ser Glu Ala Val Gln 290 295 300		
Pro Val Thr Pro Glu Pro Cys Val Thr Gln Asp Ser Val Arg Phe Ser 305 310 315 320		
His Leu Val Val Asp Leu Val Gln Ala Lys Asp Thr Leu Tyr His Val 325 330 335		
Leu Tyr Ile Gly Thr Glu Ser Gly Thr Ile Leu Lys Ala Leu Ser Thr		

340	345	350
Ala Ser Arg Ser Leu His Gly Cys Tyr Leu Glu Glu Leu His Val Leu		
355	360	365
Pro Pro Gly Arg Arg Glu Pro Leu Arg Ser Leu Arg Ile Leu His Ser		
370	375	380
Ala Arg Ala Leu Phe Val Gly Leu Arg Asp Gly Val Leu Arg Val Pro		
385	390	400
Leu Glu Arg Cys Ala Ala Tyr Arg Ser Gln Gly Ala Cys Leu Gly Ala		
405	410	415
Arg Asp Pro Tyr Cys Gly Trp Asp Gly Lys Gln Gln Arg Cys Ser Thr		
420	425	430
Leu Glu Asp Ser Ser Asn Met Ser Leu Trp Thr Gln Asn Ile Thr Ala		
435	440	445
Cys Pro Val Arg Asn Val Thr Arg Asp Gly Gly Phe Gly Pro Trp Ser		
450	455	460
Pro Trp Gln Pro Cys Glu His Leu Asp Gly Asp Asn Ser Gly Ser Cys		
465	470	475
Leu Cys Arg Ala Arg Ser Cys Asp Ser Pro Arg Pro Arg Cys Gly Gly		
485	490	495
Leu Asp Cys Leu Gly Pro Ala Ile His Ile Ala Asn Cys Ser Arg Asn		
500	505	510
Gly Ala Trp Thr Pro Trp Ser Ser Trp Ala Leu Cys Ser Thr Ser Cys		
515	520	525
Gly Ile Gly Phe Gln Val Arg Gln Arg Ser Cys Ser Asn Pro Ala Pro		
530	535	540
Arg His Gly Gly Arg Ile Cys Val Gly Lys Ser Arg Glu Glu Arg Phe		
545	550	555
Cys Asn Glu Asn Thr Pro Cys Pro Val Pro Ile Phe Trp Ala Ser Trp		
565	570	575
Gly Ser Trp Ser Lys Cys Ser Ser Asn Cys Gly Gly Gly Met Gln Ser		
580	585	590
Arg Arg Arg Ala Cys Glu Asn Gly Asn Ser Cys Leu Gly Cys Gly Val		
595	600	605
Glu Phe Lys Thr Cys Asn Pro Glu Gly Cys Pro Glu Val Arg Arg Asn		
610	615	620
Thr Pro Trp Thr Pro Trp Leu Pro Val Asn Val Thr Gln Gly Gly Ala		
625	630	635
Arg Gln Glu Gln Arg Phe Arg Phe Thr Cys Arg Ala Pro Leu Ala Asp		
645	650	655
Pro His Gly Leu Gln Phe Gly Arg Arg Arg Thr Glu Thr Arg Thr Cys		

660					665					670									
Pro	Ala	Asp	Gly	Ser	Gly	Ser	Cys	Asp	Thr	Asp	Ala	Leu	Val	Glu	Asp				
675					680					685									
Leu	Leu	Arg	Ser	Gly	Ser	Thr	Ser	Pro	His	Thr	Val	Ser	Gly	Gly	Trp				
690					695					700									
Ala	Ala	Trp	Gly	Pro	Trp	Ser	Ser	Cys	Ser	Arg	Asp	Cys	Glu	Leu	Gly				
705					710					715					720				
Phe	Arg	Val	Arg	Lys	Arg	Thr	Cys	Thr	Asn	Pro	Glu	Pro	Arg	Asn	Gly				
725					730					735									
Gly	Leu	Pro	Cys	Val	Gly	Asp	Ala	Ala	Glu	Tyr	Gln	Asp	Cys	Asn	Pro				
740					745					750									
Gln	Ala	Cys	Pro	Val	Arg	Gly	Ala	Trp	Ser	Cys	Trp	Thr	Ser	Trp	Ser				
755					760					765									
Pro	Cys	Ser	Ala	Ser	Cys	Gly	Gly	Gly	His	Tyr	Gln	Arg	Thr	Arg	Ser				
770					775					780									
Cys	Thr	Ser	Pro	Ala	Pro	Ser	Pro	Gly	Glu	Asp	Ile	Cys	Leu	Gly	Leu				
785					790					795					800				
His	Thr	Glu	Glu	Ala	Leu	Cys	Ala	Thr	Gln	Ala	Cys	Pro	Glu	Gly	Trp				
805					810					815									
Ser	Pro	Trp	Ser	Glu	Trp	Ser	Lys	Cys	Thr	Asp	Asp	Gly	Ala	Gln	Ser				
820					825					830									
Arg	Ser	Arg	His	Cys	Glu	Glu	Leu	Leu	Pro	Gly	Ser	Ser	Ala	Cys	Ala				
835					840					845									
Gly	Asn	Ser	Ser	Gln	Ser	Arg	Pro	Cys	Pro	Tyr	Ser	Glu	Ile	Pro	Val				
850					855					860									
Ile	Leu	Pro	Ala	Ser	Ser	Met	Glu	Glu	Ala	Thr	Gly	Cys	Ala	Gly	Phe				
865					870					875					880				
Asn	Leu	Ile	His	Leu	Val	Ala	Thr	Gly	Ile	Ser	Cys	Phe	Leu	Gly	Ser				
885					890					895									
Gly	Leu	Leu	Thr	Leu	Ala	Val	Tyr	Leu	Ser	Cys	Gln	His	Cys	Gln	Arg				
900					905					910									
Gln	Ser	Gln	Glu	Ser	Thr	Leu	Val	His	Pro	Ala	Thr	Pro	Asn	His	Leu				
915					920					925									
His	Tyr	Lys	Gly	Gly	Gly	Thr	Pro	Lys	Asn	Glu	Lys	Tyr	Thr	Pro	Met				
930					935					940									
Glu	Phe	Lys	Thr	Leu	Asn	Lys	Asn	Asn	Leu	Ile	Pro	Asp	Asp	Arg	Ala				
945					950					955					960				
Asn	Phe	Tyr	Pro	Leu	Gln	Gln	Thr	Asn	Val	Tyr	Thr	Thr	Thr	Tyr	Tyr				
965					970					975									
Pro	Ser	Pro	Leu	Asn	Lys	His	Ser	Phe	Arg	Pro	Glu	Ala	Ser	Pro	Gly				

980

985

990

Gln Arg Cys Phe Pro Asn Ser  
995

<210> 21  
<211> 2820  
<212> DNA  
<213> Homo sapiens

<400> 21  
agaatgggag ctgtcagtta tcattcagga caaaggcaat cctcagctac ataccaaagt 60  
ccttctgaag tgcattgatct ttgaatatgc agagtcgggtg acaagtacag caatgacttc 120  
agtaagccag gcatccttgg atgtctccat gataataatt atttccttag gagcaatttg 180  
tgtagtggtg ctggttatta tgggtgtatt tgcaactagg tgtaaccgag agaagaaaga 240  
cactagatcc tataactgca ggggtggcga atcaacttac cagcaccacc caaaaaggcc 300  
atcccgccag attcacaag gggacatcac attggtgcct accataaatg gactctgcc 360  
catcagatct catcacagat cgtctccatc ttcattctct acccttagaaa gagggcagat 420  
gggcagccgg cagagtcaca acagtcacca gtcactcaac agtttgggtg caatctcatc 480  
aaaccacgtg ccagagaatt tctcattaga actcaccac gccactcctg ctggtgaggt 540  
ctctcagctt ctttcaatgc ttcaccagg gcaatatac ccaagaccaa gtttccgag 600  
aaacaaatat tccaggagct acagatatgc ctttcaagac atggacaaat ttagcttgaa 660  
agacagtggtg cgtggtgaca gtgaggcagg agacagtgt tatgatttgg ggcgagattc 720  
tccaatagat aggtctgttg gtgaaggatt cagcgacct tttctcacag atggaagaat 780  
tccagcagct atgagactct gcacggagga gtgcagggtc ctgggacact ctgaccagt 840  
ctggatgcca ccactgccct caccgtcttc tgattatagg agtaacatgt tcatccagg 900  
ggaagaattc ccaacgcaac ccacgagca gcatccacat cagagtcttg aggatgacg 960  
tcagcctgca gattccggtg aaaagaagaa gagtttttcc accttggaa aggactcccc 1020  
aaacgatgag gacactgggg ataccagcac atcatctctg ctctcggaaa tgagcagtg 1080  
gttccagcgt ctcttaccgc tttccctgga cacctattct gaatgcagtg aggtggatcg 1140  
gtccaaactcc ctggagcgca ggaagggacc cttgccagcc aaaactgttg gttaccaca 1200  
gggggttagc gcatgggcag ccagtagcga ttttcaaaat cccaccacca actgtggg 1260  
gccacttggg actcactcca gtgtgcagcc ttttcaaaa tggctgccag ccatggagga 1320  
gatccctgaa aattatgagg aagatgatt tgacaatgtg ctcaaccacc tcaatgatgg 1380  
gaacacgaa ctcatggatg ccagtgaact ggtggcagag attaacaaac tgcttcaaga 1440  
tgtccgccag agctaggaga ttttagcgaa gcatttttgt tccatgtat atggaaatag 1500  
ggaacaacaa caacaacaaa aaacctgaa agaactggca ttgccaaaata gttgcattta 1560  
tcataaatgt gtctgtgtat attgaatatt aaatactgta tttctgtatg tacacaatgc 1620  
aagtgtgatt attttaatct gtatttttaa aatacatttg taccttatat ttatgtgtaa 1680  
tttaacaaac aaattttatt tttttactcc catgacagac atgtttttcc tagtcgtgta 1740  
gaaactagcc actgttcaaa tctgatacac tattcaacca caaagtgtaa aggactgct 1800  
tagattagtt ttgttgggga agaattatta tgtgtatga acaacccac tgaagcata 1860  
tacaattctt aattccatta agtgatccca ctttttttca ataactttt agaaattaa 1920  
aatcattaaa attgttaagc tattttattg ttattttctc tactttctac tagccccaat 1980  
agttgaactc ttataggaaa atcgaaagat aaagtgaag tttatttcag gactgagaaa 2040  
tatcttgaag gttatttatt agatgactat ctcaaatgaa ctttttatag acaatgatga 2100  
aaacagaatt aaagtcaatg tttcctgact ccaggcccc tactattcca ggccatcaca 2160  
ctggcctgtt cgggagaata tttctctcac aatattatta tctacttata attatggtaa 2220  
acaataaatt ttattccatc cttgtagtat gaaacatgct ccaaggaaat ggaatctgtc 2280  
ctttaaatgg ataacagtat gtgttctaata ggcataaaat attactggat aaaaacagtt 2340  
gtgtcagtgct ctctcctaag gtagtaata taattgactt attctgaacc cattctattt 2400  
tgaatctccc ctttctctc acaatacttg aacattttaa tcttttggaa tattgtcttt 2460  
ctttgttata actattcatt tttagctttt gtctccagtg catgatctca tatttttgct 2520  
tttattttta gtataagaac atttataaaa tcatattttt gttactgcaa ttgttttatt 2580  
tggtgtgtgg caaatgagaa atcctttatt tattgtgctg tgatctctct gtgtggaatg 2640  
ccttgggtgag agagatgctt attatgacta ttatcatttc tgaccaagct tctattaatg 2700  
ttattttctaa taatacacta tcttgattgt actctccaga aaatttttct gtcagtgaat 2760  
ataaaagaaa aattaaagta aagctaagga actgtcaaaa aaaaaaaaaa aaaaaaaaaa 2820

<210> 22  
 <211> 460  
 <212> PRT  
 <213> Homo sapiens

<400> 22

```

Met Ile Phe Glu Tyr Ala Glu Ser Val Thr Ser Thr Ala Met Thr Ser
  1             5             10             15

Val Ser Gln Ala Ser Leu Asp Val Ser Met Ile Ile Ile Ile Ser Leu
      20             25             30

Gly Ala Ile Cys Ala Val Leu Leu Val Ile Met Val Leu Phe Ala Thr
      35             40             45

Arg Cys Asn Arg Glu Lys Lys Asp Thr Arg Ser Tyr Asn Cys Arg Val
      50             55             60

Ala Glu Ser Thr Tyr Gln His His Pro Lys Arg Pro Ser Arg Gln Ile
      65             70             75             80

His Lys Gly Asp Ile Thr Leu Val Pro Thr Ile Asn Gly Thr Leu Pro
      85             90             95

Ile Arg Ser His His Arg Ser Ser Pro Ser Ser Ser Pro Thr Leu Glu
      100            105            110

Arg Gly Gln Met Gly Ser Arg Gln Ser His Asn Ser His Gln Ser Leu
      115            120            125

Asn Ser Leu Val Thr Ile Ser Ser Asn His Val Pro Glu Asn Phe Ser
      130            135            140

Leu Glu Leu Thr His Ala Thr Pro Ala Val Glu Val Ser Gln Leu Leu
      145            150            155            160

Ser Met Leu His Gln Gly Gln Tyr Gln Pro Arg Pro Ser Phe Arg Gly
      165            170            175

Asn Lys Tyr Ser Arg Ser Tyr Arg Tyr Ala Leu Gln Asp Met Asp Lys
      180            185            190

Phe Ser Leu Lys Asp Ser Gly Arg Gly Asp Ser Glu Ala Gly Asp Ser
      195            200            205

Asp Tyr Asp Leu Gly Arg Asp Ser Pro Ile Asp Arg Leu Leu Gly Glu
      210            215            220

Gly Phe Ser Asp Leu Phe Leu Thr Asp Gly Arg Ile Pro Ala Ala Met
      225            230            235            240

Arg Leu Cys Thr Glu Glu Cys Arg Val Leu Gly His Ser Asp Gln Cys
      245            250            255

Trp Met Pro Pro Leu Pro Ser Pro Ser Ser Asp Tyr Arg Ser Asn Met
      260            265            270

Phe Ile Pro Gly Glu Glu Phe Pro Thr Gln Pro Gln Gln Gln His Pro
      275            280            285

```

His Gln Ser Leu Glu Asp Asp Ala Gln Pro Ala Asp Ser Gly Glu Lys  
 290 295 300  
 Lys Lys Ser Phe Ser Thr Phe Gly Lys Asp Ser Pro Asn Asp Glu Asp  
 305 310 315 320  
 Thr Gly Asp Thr Ser Thr Ser Ser Leu Leu Ser Glu Met Ser Ser Val  
 325 330 335  
 Phe Gln Arg Leu Leu Pro Pro Ser Leu Asp Thr Tyr Ser Glu Cys Ser  
 340 345 350  
 Glu Val Asp Arg Ser Asn Ser Leu Glu Arg Arg Lys Gly Pro Leu Pro  
 355 360 365  
 Ala Lys Thr Val Gly Tyr Pro Gln Gly Val Ala Ala Trp Ala Ala Ser  
 370 375 380  
 Thr His Phe Gln Asn Pro Thr Thr Asn Cys Gly Pro Pro Leu Gly Thr  
 385 390 395 400  
 His Ser Ser Val Gln Pro Ser Ser Lys Trp Leu Pro Ala Met Glu Glu  
 405 410 415  
 Ile Pro Glu Asn Tyr Glu Glu Asp Asp Phe Asp Asn Val Leu Asn His  
 420 425 430  
 Leu Asn Asp Gly Lys His Glu Leu Met Asp Ala Ser Glu Leu Val Ala  
 435 440 445  
 Glu Ile Asn Lys Leu Leu Gln Asp Val Arg Gln Ser  
 450 455 460

&lt;210&gt; 23

&lt;211&gt; 1219

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

gtgggccattc ctcggtacag actagtcctg gtccttgggt gtgggcagtg ggggaggaac 60  
 caactgggtcg aggtttcaga gccaaacctt gcctttgggt ggtgagtcct tgccccccag 120  
 gcctgcgctc cacgatgccc ttcaccttg gcaatctcag ggccatcctg ggtagtaacc 180  
 ccactcctct ctgctcccg cgcacactgt ggctctcact ctgggctcaa cccctgcaac 240  
 cctccaggag cccgacagca gccagctgcc tgcactgtcg cctccgtaag ctccaacttc 300  
 cagacccaga agtcctctct ctccctctg ttggaaaaag cctaaaagaa ttagctttca 360  
 gattcctcta gccctgctc cattcccacc cagtccttct gaagaggaat gagcaatata 420  
 tctgagctgg atttctctct agtcctttct ccagacaaat ccttcttaaa gcaaaagtcc 480  
 tggctgagca cctgtccttg gggaccgatc tgccgtgtga ccaggggag aaagtcccg 540  
 aaagcctggt ccaccaattc tgcttctgtg ttgtgaatcc agtctgcttt ccattagaaa 600  
 accgcttcgg cacttatggt cactttaata aatctagtat gtaaaaaag aaagaaagaa 660  
 aagaaacaga aaaacgtgca ggcacatgta aaatacaatg ctctctgtaa gataaatatt 720  
 tgcctttttt tctaaaagg gtacgtatcc tgtatgtgaa attgtctgta gaaagtttct 780  
 atgttcttaa atggcaatac attccaaaaa ttgtactgta gatagtaca gcaaccgcac 840  
 tgggatgggg tagttttgcc tgtaatttta tttaaactcc agtttccaca cttgcatctt 900  
 gcaatgttgg tatggtatat atcagtgcaa aagaaaaaac aaaacaaaaa caaaaaaaa 960  
 aacaaaaatc cagcgaggtc taaagcacag agtctgacgt acaaaaggaa aaatgctcag 1020  
 tattgaagtg tgtgaccttt gttgtaaatt acatctgtac tgtgaatgag aagtttttac 1080  
 aagtataata attgccttta ttacagctct ggctgagtg tgcagctgag gatatttttt 1140  
 aaaaaaaaaa gaattagcat gttggaataa atttgaaaat cccaaaaaaa aaaaaaaaaa 1200



aaaaaaaaa aaaaaaaaaa

1219

&lt;210&gt; 24

&lt;211&gt; 78

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 24

Met Pro Phe Thr Leu Gly Asn Leu Arg Ala Ile Leu Gly Ser Asn Pro  
 1 5 10 15

Thr Pro Leu Cys Ser Arg Pro His Leu Trp Leu Ser Leu Trp Ala Gln  
 20 25 30

Pro Leu Gln Pro Ser Arg Ser Pro Thr Ala Ala Ser Cys Leu His Cys  
 35 40 45

Arg Leu Arg Lys Leu Gln Leu Pro Asp Pro Glu Val Pro Leu Leu Pro  
 50 55 60

Ser Val Gly Lys Ser Leu Lys Glu Leu Ala Ser Arg Phe Leu  
 65 70 75

&lt;210&gt; 25

&lt;211&gt; 2411

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 25

ggccagaagg cggggagcca gaggcgccag gaccctagcg tggcgctcca gcacccaga 60  
 ccgtggcggc gcctcgccct agggaagagc aagggaagaa ctttatttga accgcgaaca 120  
 ttttttggtc actgagatcg agtctccag tgctttggct tcccgcctct ttatcgtggg 180  
 tttgatccct gagctgctct cctttccga acctcccggt gtgcagccta gagccctccc 240  
 gcgcggctga ctccagagta gaggaaggga ggcggcctcc ggctgggtccc ccgaagccct 300  
 cgctgccccg cagatgcgga tggccagcca gtagcggggt gtggccccgc gtcccgggag 360  
 cgcacagcaa tgcaggcgct taacattacc ccggagcagt tctctcggt gctgcgggac 420  
 cacaacctga cgcgggagca gttcatcgct ctgtaccggc tgcgaccgct cgtctacacc 480  
 ccagagctgc cgggacgcgc caagctggcc ctctgtctca ccggcgtgct catcttcgcc 540  
 ctggcgctct ttggcaatgc tctgggtgtc tacgtgtgta cccgcagcaa ggccatgcgc 600  
 accgtcacca acatctttat ctgctccttg gcgctcagt acctgctcat cactctcttc 660  
 tgcattcccc tcaccatgct ccagaacatt tccgacaact ggctgggggg tgctttcatt 720  
 tgcaagatgg tgccatttgt ccagtctacc gctgtgtgta cagaaatcct cactatgacc 780  
 tgcattgctg tggaaaggca ccagggactt gtgcatecct ttaaaatgaa gtggcaatac 840  
 accaaccgaa gggtttcac aatgctaggt gtggtctggc tgggtggcagt catcgtagga 900  
 tcacccatgt ggcacgtgca acaacttgag atcaaatatg acttcctata tgaaaaggaa 960  
 cacatctgct gcttagaaga gtggaccagc cctgtgcacc agaagatcta caccaccttc 1020  
 atccttgta tcctcttcct cctgcctctt atggtgatgc ttattctgta cagtaaaatt 1080  
 ggttatgaac tttggataaa gaaaagagtt ggggatgggt cagtgcctcg aactattcat 1140  
 ggaaaagaaa tgtccaaaat agccaggaag aagaaacgag ctgtcattat gatggtgaca 1200  
 gtggtggctc tctttgctgt gtgctgggca ccattccatg ttgtccatat gatgattgaa 1260  
 tacagtaatt ttgaaaagga atatgatgat gtcacaaatca agatgatttt tgctatcgtg 1320  
 caaattattg gattttccaa ctccatctgt aatccattg tctatgcatt tatgaatgaa 1380  
 aacttcaaaa aaaatgtttt gtctgcagtt tgttattgca tagtaataaa aaccttctct 1440  
 ccagcacaaa ggcattgaaa ttcaggaatt acaatgatgc ggaagaaagc aaagtcttcc 1500  
 ctgagagaga atccagtgga ggaaaccaa ggagaagcat tcagtgatgg caacattgaa 1560  
 gtcaaattgt gtgaacagac agaggagaag aaaaagctca aacgacatct tgctctcttt 1620  
 aggtctgaac tggctgagaa ttctccttta gacagtgggc attaattata acaatatctt 1680  
 cataattaat gcccttcaga ttgtaaccca aagagaaaat tattttgagc aaaggtcaaa 1740  
 tactcttttt attcttaaga tgatgacaag aagaaaacaa atcatgtttc cattaataaa 1800

```

tgacacgagg ctagtccaag tgcagtgatg tttacaacca attgatcaca atcattttaac 1860
agattttctgt gttccttctc attcctactg cttcacttga ctagccttaa aaaagcaaca 1920
tggaaggcca ggcacggtgg ctcatgcctg taatcccagc actttgggag gcctagacgg 1980
gcggatcacg aggtcaggag atcaaaacca tcctggctaa cacggtgaaa ccccatctct 2040
gctaaaaata caaaaattag ccgggcgtgg tggcgggcac ctgtagtccc agctacttgg 2100
gagcctcagg cgggagaatg gtgtgaaccc gggaggcgga gcttgcagtg atccgagatc 2160
atgccactgc actccagcct gggcgaaaga gcgagactcc ccgtctcaaa aaaaattttt 2220
ttgaaaaatt cgtaaaccat acttttaaga ttatttcagt ggatttttaa aaatcttgta 2280
cagaaatcag ggttcttagc tagcagtttt tctcccacgc agtcactgta atgtgactat 2340
gtattgctag attgaataag aaaataaaat aatatcttct tccttgaaaa aaaaaaaaaa 2400
aaaaaaaaaa a 2411

```

<210> 26

<211> 431

<212> PRT

<213> Homo sapiens

<400> 26

```

Met Gln Ala Leu Asn Ile Thr Pro Glu Gln Phe Ser Arg Leu Leu Arg
  1             5             10             15

Asp His Asn Leu Thr Arg Glu Gln Phe Ile Ala Leu Tyr Arg Leu Arg
          20             25             30

Pro Leu Val Tyr Thr Pro Glu Leu Pro Gly Arg Ala Lys Leu Ala Leu
      35             40             45

Val Leu Thr Gly Val Leu Ile Phe Ala Leu Ala Leu Phe Gly Asn Ala
      50             55             60

Leu Val Phe Tyr Val Val Thr Arg Ser Lys Ala Met Arg Thr Val Thr
      65             70             75             80

Asn Ile Phe Ile Cys Ser Leu Ala Leu Ser Asp Leu Leu Ile Thr Phe
          85             90             95

Phe Cys Ile Pro Val Thr Met Leu Gln Asn Ile Ser Asp Asn Trp Leu
      100             105             110

Gly Gly Ala Phe Ile Cys Lys Met Val Pro Phe Val Gln Ser Thr Ala
      115             120             125

Val Val Thr Glu Ile Leu Thr Met Thr Cys Ile Ala Val Glu Arg His
      130             135             140

Gln Gly Leu Val His Pro Phe Lys Met Lys Trp Gln Tyr Thr Asn Arg
      145             150             155             160

Arg Ala Phe Thr Met Leu Gly Val Val Trp Leu Val Ala Val Ile Val
          165             170             175

Gly Ser Pro Met Trp His Val Gln Gln Leu Glu Ile Lys Tyr Asp Phe
      180             185             190

Leu Tyr Glu Lys Glu His Ile Cys Cys Leu Glu Glu Trp Thr Ser Pro
      195             200             205

Val His Gln Lys Ile Tyr Thr Thr Phe Ile Leu Val Ile Leu Phe Leu
      210             215             220

```

Leu Pro Leu Met Val Met Leu Ile Leu Tyr Ser Lys Ile Gly Tyr Glu  
 225 230 235 240  
 Leu Trp Ile Lys Lys Arg Val Gly Asp Gly Ser Val Leu Arg Thr Ile  
 245 250 255  
 His Gly Lys Glu Met Ser Lys Ile Ala Arg Lys Lys Lys Arg Ala Val  
 260 265 270  
 Ile Met Met Val Thr Val Val Ala Leu Phe Ala Val Cys Trp Ala Pro  
 275 280 285  
 Phe His Val Val His Met Met Ile Glu Tyr Ser Asn Phe Glu Lys Glu  
 290 295 300  
 Tyr Asp Asp Val Thr Ile Lys Met Ile Phe Ala Ile Val Gln Ile Ile  
 305 310 315 320  
 Gly Phe Ser Asn Ser Ile Cys Asn Pro Ile Val Tyr Ala Phe Met Asn  
 325 330 335  
 Glu Asn Phe Lys Lys Asn Val Leu Ser Ala Val Cys Tyr Cys Ile Val  
 340 345 350  
 Asn Lys Thr Phe Ser Pro Ala Gln Arg His Gly Asn Ser Gly Ile Thr  
 355 360 365  
 Met Met Arg Lys Lys Ala Lys Phe Ser Leu Arg Glu Asn Pro Val Glu  
 370 375 380  
 Glu Thr Lys Gly Glu Ala Phe Ser Asp Gly Asn Ile Glu Val Lys Leu  
 385 390 395 400  
 Cys Glu Gln Thr Glu Glu Lys Lys Lys Leu Lys Arg His Leu Ala Leu  
 405 410 415  
 Phe Arg Ser Glu Leu Ala Glu Asn Ser Pro Leu Asp Ser Gly His  
 420 425 430

&lt;210&gt; 27

&lt;211&gt; 1945

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

atcatgccac tcatttcaga acttgagcaa aacagggcag tcaggatctg atgtctttct 60  
 ggtctcccta agaaaactaa gctcttgagg gacagccctt ggcaatgctt tcctatctgc 120  
 tgagcatggt gaccttcctt aggacttcca gagttcagtt ccttctggca gagaggtttt 180  
 ctttctccat gccatatgga tgtgactcaa atgaggggtc ccacagcttt tcctggctac 240  
 cacttgctgt gaccttatac atgttgggtt ttgctcttaa agaggagagc aggaagaaag 300  
 gttggtttca gaaaccaaga gggtcggcag tggacgcgta cattttgtca cggagtccac 360  
 agagctgagc ttttgagcag actctgagaa gtatcattgc ttgtgttgaa agaatacaac 420  
 aggatttaag tttctcttta aaaattgcac tgaagaaagg ccgggcgagg tggctcccc 480  
 tgtaatccca gcgctttggg aggccgaggc ggggggatca cgagggtcaag agatcgagac 540  
 catectggcc aacatggtga aaccccgctt ctaataaaaa tacaaaaatt agccgggcat 600  
 ggtgacgtgc acctgtagtc ccagctacta gataggctga ggcaggagaa ttgcttgaat 660  
 ccgggaggcg gaggttgagc tgagccgaga tcgtgccact gaactccaac ctgccaatag 720  
 agcgagactc cgtctcaaaa aaaaaaaaaa aaaagaaaga aatagcattg aagaaaatac 780  
 cgcacatcag aggaaagctt attttctgca tgggtgtctt tcaaagatag aatatttgaa 840

```

gcattgttttc tagcgattgt gtgaatgagg gtgagctggc tgaggcatcg ctcaagctgg 900
gggggtggtgt gtaagaagca cgtggagcca caagaggcac ctccatatagt cagctaaggg 960
cttcccttttc tgcgcccagc ttttgggtga aggggtgattt ctactagaca catctgtgct 1020
tcagtcatag atgttaatag aggaagcagt tttcctgctg cagattcctg aatagagttg 1080
ctgaaagagt ctacttctgg actcgggaag ttgaaggcca gtctgtgtag aaaggctgag 1140
gcaacgggga aagacctgac agctagttac atacgctctg acatagtact cccatgatgg 1200
cttccagtga cacatgtgct gatagaattc taaacctctg gaatttcctt gctggcgact 1260
tctatggccg ttgactgtac agggtaacct gatgccagat gctatgggag tgatgagaac 1320
tagagcattg cagcatggag gaaactgtga ggcaccagat cctgtgcttc tgcaggccat 1380
tttctgaaaa cccctgttag gaaggttgga tttggcgtga cttgcttgag caagagtcct 1440
ggggagagat tttgagttt aatttaacgg tatatccaga gctaacagtg actcaactcg 1500
tctagtctct caagtcagat gtatacttag agtctctctg tgaagggttt ggggtctgagc 1560
tgtatagtat gtcaaaactgc cagtaagcca gcccctcacc ctctgataga tattccttta 1620
atgcaccaga cttcatgttt gataaatgat taatgggtga aattgtttct cttcttttgt 1680
gttttcccag ttaatagatg gtcactgttt ccacaatgtt ttatacttcc agctttttgt 1740
aacttaacta taattactta attttatttt tttaaagctt gttgtggtct aatgagaagt 1800
atttttcagt gcataatgtt tttctgagct tctgtaaatg ccatcccaat gtgggttggt 1860
tttgttgaac agaaacaaa ataaatttca aaatgttaaa aaaaaaaaaa aaaaaaaaaa 1920
aaaaaaaaaa aaaaaaaaaa aaaaaa 1945

```

&lt;210&gt; 28

&lt;211&gt; 87

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 28

```

Met Leu Ser Tyr Leu Leu Ser Met Val Thr Phe Leu Arg Thr Ser Arg
 1             5             10             15

Val Gln Phe Leu Leu Ala Glu Arg Phe Ser Phe Ser Met Pro Tyr Gly
          20             25             30

Cys Asp Ser Asn Glu Gly Ser His Ser Phe Ser Trp Leu Pro Leu Ala
          35             40             45

Val Thr Leu Tyr Met Leu Gly Phe Ala Leu Lys Glu Glu Ser Arg Lys
          50             55             60

Lys Gly Trp Phe Gln Lys Pro Arg Gly Ser Ala Val Asp Ala Tyr Ile
          65             70             75             80

Leu Ser Arg Ser Pro Gln Ser
          85

```

&lt;210&gt; 29

&lt;211&gt; 2184

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 29

```

ggggcgccctc tggggctccg agcccggcgg gaccatgttc accagcaccg gctccagtgg 60
gctctacaag gcgcctctgt cgaagagcct tctgctggtc cccagtgcc cttccctcct 120
gctcgccctc ctctgcctc actgccagaa gctctttgtg tatgaccttc acgcagtcaa 180
gaacgacttc cagatttggg ggttgatag tggaagaata atttgccttg atttgaaaga 240
tactttttgc agtagtctgc ttatttataa ttttaggata tttgaaagaa gatatggaag 300
cagaaaattt gcatcctttt tgctgggttc ctgggttttg tcagccttat ttgactttct 360
cctcattgaa gctatgcagt atttcttttg catcactgca gctagtaatt tgccttctgg 420
attcctggca cctgtgtttg ctctgtttgt accattttac tgctccatac caagagtcca 480
agtggcacaa attctgggtc cgttggtccat cacaaacaag acattgattt atatattggg 540

```

```

actgcagctt ttcacctctg gttcctacat ctggattgta gccataagtg gacttatgtc 600
cggctctgtc tacgacagca aaatgttcca ggtgcatcag gtgctctgca tccccagctg 660
gatggcaaaa ttcttttctt ggacacttga acccatcttc tcttcttcag aaccaccag 720
cgaagccaga attgggatgg gagccacgct ggacatccag agacagcaga gaatggagct 780
gctggaccgg cagctgatgt tctctcagtt tgcacaaggg aggcgacaga gacagcagca 840
gggaggaatg atcaattgga atcgtctttt tcctccttta cgtcagcgac aaaacgtaaa 900
ctatcagggc ggtcggcagt ctgagccagc agcgccccct ctagaagttt ctgaggaaca 960
ggtcgcccg ctcattgaga tgggattttc cagaggtgat gctttggaag ccctgagagc 1020
ttcaaacaat gacctcaatg tcgccaccaa ctctctgctg cagcactgat agtcccaggc 1080
caacactggg accggaccgg cagccgagtg acagtgcgtg gtccccacca tcagatcagc 1140
ccggggaccg agcatctctg gtgctgatgt tcttggtgga agagggaggt tccaccgcac 1200
ccctgccctc aaccgcaaga ctggtgccgt tttagtgtgg agataagttt gccattacat 1260
tagcatgtat ttctatctta tatttttatt gggcattttc cctaggttgg agagtcagca 1320
ctcgttttga atgtgtttaa aatgcattaa aatggaagat ttctgcaggc agttgaatgg 1380
cactccagat ggggaattgc tgtaaccctc ttactgtaac atgtcatctc ctgcgtcgtg 1440
atggggagag ggtaattgta cttcacaaag gacatgtcag atccttcttc atggactttt 1500
ttagtactct ttttttctct caaacttggt ttcgaaatct ctgggagtga gggagaaaac 1560
gggagctgaa tcctccccc aagctgttcca ggccagagga ctctgcagta gcttctccta 1620
catctagtaa caaagaatgg tgataaccat gcactgggtc aaggttctgg agttctccat 1680
gaaacttggg ttaattttgc tcagagtatc cagagttagc cactaggctg cgggtgaaat 1740
gggatggaga agaacaacag caggcttctt ggagccacat gggctgacta gggcactctg 1800
tggctggcat ggcatgggct cagcccagga agaggagaaa cgatcccttg cctgccctc 1860
cctgtggcag ggctaactgc ctggccctcc tggctcgcag ccagccagcc cctgggcagc 1920
aggttctctc cagggcttgg gtcttcaacc tgtggcgaca ggagggcagg cagactgtgg 1980
aggacaggat gcaggtcagg gagaggggag gcaggggtgg accgccatga gcatgaaaag 2040
acccgaagca agttgactct tgcaatgtgc aactgttatg ttctgcaaaa tgagcaacga 2100
tgtatcaaat tgatgcaaat ttagatgttg atacttacaa taaagttttt aatgtgtttt 2160
aaaaaaaaa aaaaaaaaaa aaaa
2184

```

<210> 30

<211> 344

<212> PRT

<213> Homo sapiens

<400> 30

```

Met Phe Thr Ser Thr Gly Ser Ser Gly Leu Tyr Lys Ala Pro Leu Ser
 1             5             10             15

Lys Ser Leu Leu Leu Val Pro Ser Ala Leu Ser Leu Leu Leu Ala Leu
          20             25             30

Leu Leu Pro His Cys Gln Lys Leu Phe Val Tyr Asp Leu His Ala Val
          35             40             45

Lys Asn Asp Phe Gln Ile Trp Arg Leu Ile Cys Gly Arg Ile Ile Cys
          50             55             60

Leu Asp Leu Lys Asp Thr Phe Cys Ser Ser Leu Leu Ile Tyr Asn Phe
          65             70             75             80

Arg Ile Phe Glu Arg Arg Tyr Gly Ser Arg Lys Phe Ala Ser Phe Leu
          85             90             95

Leu Gly Ser Trp Val Leu Ser Ala Leu Phe Asp Phe Leu Leu Ile Glu
          100            105            110

Ala Met Gln Tyr Phe Phe Gly Ile Thr Ala Ala Ser Asn Leu Pro Ser
          115            120            125

Gly Phe Leu Ala Pro Val Phe Ala Leu Phe Val Pro Phe Tyr Cys Ser

```

130 135 140  
 Ile Pro Arg Val Gln Val Ala Gln Ile Leu Gly Pro Leu Ser Ile Thr  
 145 150 155 160  
 Asn Lys Thr Leu Ile Tyr Ile Leu Gly Leu Gln Leu Phe Thr Ser Gly  
 165 170 175  
 Ser Tyr Ile Trp Ile Val Ala Ile Ser Gly Leu Met Ser Gly Leu Cys  
 180 185 190  
 Tyr Asp Ser Lys Met Phe Gln Val His Gln Val Leu Cys Ile Pro Ser  
 195 200 205  
 Trp Met Ala Lys Phe Phe Ser Trp Thr Leu Glu Pro Ile Phe Ser Ser  
 210 215 220  
 Ser Glu Pro Thr Ser Glu Ala Arg Ile Gly Met Gly Ala Thr Leu Asp  
 225 230 235 240  
 Ile Gln Arg Gln Gln Arg Met Glu Leu Leu Asp Arg Gln Leu Met Phe  
 245 250 255  
 Ser Gln Phe Ala Gln Gly Arg Arg Gln Arg Gln Gln Gln Gly Gly Met  
 260 265 270  
 Ile Asn Trp Asn Arg Leu Phe Pro Pro Leu Arg Gln Arg Gln Asn Val  
 275 280 285  
 Asn Tyr Gln Gly Gly Arg Gln Ser Glu Pro Ala Ala Pro Pro Leu Glu  
 290 295 300  
 Val Ser Glu Glu Gln Val Ala Arg Leu Met Glu Met Gly Phe Ser Arg  
 305 310 315 320  
 Gly Asp Ala Leu Glu Ala Leu Arg Ala Ser Asn Asn Asp Leu Asn Val  
 325 330 335  
 Ala Thr Asn Phe Leu Leu Gln His  
 340

&lt;210&gt; 31

&lt;211&gt; 2880

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 31

ccggcgctccg ggcgcgctgg agaggacgcg aggagccatg agggcgccagc ctgcgaaggt 60  
 ggcggcgctg ctgctcgggc tgctcttgga gtgcacagaa gccaaaaagc attgctggta 120  
 ttctgaagga ctctatccaa cctattatat atgccgctcc tacgaggact gctgtggctc 180  
 caggtgctgt gtgcgggccc tctccataga gaggctgtgg tacttctggt tccttctgat 240  
 gatgggctg cttttctgct gcggagccgg cttcttcac cggaggcgca tgtaccccc 300  
 gccgctgac gaggagccag cttcaatgt gtccacacc aggcagcccc caaatccccg 360  
 cccaggagcc cagcagccgg ggccgcccta ttacaccgac ccaggaggac cggggatgaa 420  
 ccctgtcggg aattccatgg caatggcttt ccaggtecca cccaactcac cccaggggag 480  
 tgtggcctgc ccgccccctc cagcctactg caacacgcct ccgccccctg acgaacaggt 540  
 agtgaaggcc aagtagtggg gtgcccacgt gcaagaggag agacaggaga gggcctttcc 600  
 ctggcctttc tgcttctgt gatgttact tccaggaacg gtctcgtggg ctgctaaggg 660  
 cagttcctct gatataccta cagcaagcac agctctcttt caggctttcc atggagtaca 720

```

atatatgaac tcacactttg tctcctctgt tgcttctgtt tctgacgcag tctgtgctct 780
cacatggtag tgtggtgaca gtccccgagg gctgacgtcc ttacggtggc gtgaccagat 840
ctacaggaga gagactgaga ggaagaaggc agtgctggag gtgcagggtg catgtagagg 900
ggccaggccg agcatccagc gcaagcatcc ttctgcccgg gtattaatag gaagcccat 960
gccggggcgg tcagccgatg aagcagcagc cgactgagct gagcccagca ggtcatctgc 1020
tccagcctgt cctctcgtca gccttctctt tccagaagct gttggagaga cattcaggag 1080
agagcaagcc ccttgtcatg ttctgtctc tggtcatatc ctaaagatag acttctctctg 1140
caccgccagg gaagggtagc acgtgacgtc ctcaccgcag gatggggcct agaatcaggc 1200
ttgccttgga ggcctgacag tgatctgaca tccactaagc aaatttattt aaattcatgg 1260
gaaatcactt cctgccccaa actgagacat tgcattttgt gagctcttgg tctgatttgg 1320
agaaaggact gttaccattt tttttggtgt gtttatggaa gtgcatgtag agcgtcctgc 1380
cctttgaaat cagactgggt gtgtgtcttc cctggacatc actgcctctc cagggcattc 1440
tcaggccggg ggtgtctcctt cctcaggca gctccagtgg tgggttctga aggggtgctt 1500
caaacggggg cacatctggc tgggaagtca catggactct tccagggaga gagaccagct 1560
gaggcgctct tctctgaggt tgtgttgggt ctaagcgggt gtgtgctggg ctccaaggag 1620
gaggagcttg ctgggaaaag acaggagaag tactgactca actgcactga ccatgttgtc 1680
ataattagaa taaagaagaa gtggctcgaa atgcacattc ctggatagga atcacagctc 1740
acccaggat ctcacaggta gtctcctgag tagttgacgg ctacggggga gctagtctccg 1800
ccgcatagtt atagtgttga tgtgtgaacg ctgacctgtc ctgtgtgcta agagctatgc 1860
agcttagctg aggcgcctag attactagat gtgctgtatc acggggaatg aggtgggggt 1920
gcttattttt taatgaacta atcagagcct cttgagaaat tgttactcat tgaactggag 1980
catcaagaca tctcatggaa gtggatacgg agtgatttgg tgtccatgct ttctactctg 2040
aggacattta atcggagaac ctcttgggga attttgtggg agacacttgg gaacaaaaca 2100
gacaccttgg gaatgcagtt gcaagcacag atgctgccac cagtgtctct gaccacctg 2160
gtgtgactgc tgactgccag cgtgttacct cccatgctgc aggcctccat ctaaatgaga 2220
caacaaagca caatgttcac tgtttacaac caagacaact gcgtgggtcc aaacactcct 2280
cttctctcag gtcatttgggt ttgcattttt aatgtcttta tttttgttaa tgaaaaagca 2340
cactaagctg cccctggaat cgggtgcagc tgaataggca cccaaaagtc cgtgactaaa 2400
tttcgtttgt ctttttgata gcaaattatg ttaagagaca gtgatggcta gggctcaaca 2460
atthtgtatt cccatgtttg tgtgagacag agtttgtttt cccttgaact tgggttagaat 2520
tgtgtactg tgaacgttga tcttgcatac ggaagtcccg cttcgggtgac atttctctggc 2580
cattcttgggt tccattgtgt ggatgggtgg ttgtgccac ttctggagt gagacagctc 2640
ctgggtgtgta gaattcccg agcgtccgtg gttcagagta aacttgaagc agatctgtgc 2700
atgcttttct ctgcaacaat tggctcgttt ctcttttttg ttctcttttg ataggatcct 2760
gtttctctat gtgtcaaaat aaaaataaat ttgggcaaaa aaaaaaaaaa aaaaaaaaaa 2820
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2880

```

&lt;210&gt; 32

&lt;211&gt; 172

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 32

```

Met Arg Arg Gln Pro Ala Lys Val Ala Ala Leu Leu Leu Gly Leu Leu
  1              5              10              15

```

```

Leu Glu Cys Thr Glu Ala Lys Lys His Cys Trp Tyr Phe Glu Gly Leu
      20              25              30

```

```

Tyr Pro Thr Tyr Tyr Ile Cys Arg Ser Tyr Glu Asp Cys Cys Gly Ser
      35              40              45

```

```

Arg Cys Cys Val Arg Ala Leu Ser Ile Gln Arg Leu Trp Tyr Phe Trp
      50              55              60

```

```

Phe Leu Leu Met Met Gly Val Leu Phe Cys Cys Gly Ala Gly Phe Phe
      65              70              75              80

```

```

Ile Arg Arg Arg Met Tyr Pro Pro Pro Leu Ile Glu Glu Pro Ala Phe
      85              90              95

```

Asn Val Ser Tyr Thr Arg Gln Pro Pro Asn Pro Gly Pro Gly Ala Gln  
 100 105 110  
 Gln Pro Gly Pro Pro Tyr Tyr Thr Asp Pro Gly Gly Pro Gly Met Asn  
 115 120 125  
 Pro Val Gly Asn Ser Met Ala Met Ala Phe Gln Val Pro Pro Asn Ser  
 130 135 140  
 Pro Gln Gly Ser Val Ala Cys Pro Pro Pro Pro Ala Tyr Cys Asn Thr  
 145 150 155 160  
 Pro Pro Pro Pro Tyr Glu Gln Val Val Lys Ala Lys  
 165 170

&lt;210&gt; 33

&lt;211&gt; 3406

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

taaagaacaa tcctttaagg gagaacctag aagccattca acaagggttaa aatcttcagg 60  
 cttccgagga ttgtgtagac agatcagagg cacgtttccc acaactgcga agaggcgctg 120  
 aggcaattct gcaagaagat ttgtgggttt tggaaaagaa gctatggaaa acggaggggc 180  
 aggcactctg cagataaggc aagtcctgct tttctttgtt ttgctgggaa tgtctcaggc 240  
 gggctctgaa actgggaact ttttgggtgat ggaggaattg cagagcggga gctttgtagg 300  
 aaatttggca aagaccctgg gactcgaggt gactgagctg tcttcgcggg gggctcgggt 360  
 ggtttctaata gatacaaaag agtgtttgca gctggacaca aacactgggg atttgcctct 420  
 gagagaaatg ctagacaggg aggagctctg tggctccaat gagccttggtg tgcgtgattt 480  
 ccaagtgtta atgaaaaaac ccacgcagtt tttacaaatt gagctccagg tcagggatatt 540  
 aatgatcac tctcccgctt tcttggaaaa agaaatgctc ttagaaatcc cagagaacag 600  
 tcctgttggt gctgtgttct tgcttgaaag tgcaaaggat ttagatgtag gaatcaatgc 660  
 tgtaaaaagc tacacaataa atccgaactc tcatttccac gttaaaataa gactcaatcc 720  
 agacaatagg aaataccctg agttagttct ggacaaggcg ctggattatg aagagcgccc 780  
 ggagctcagt ttcattcctca ctgctctgga tggcgggtcc cctcccagggt ctggaactgc 840  
 cttgtgcagg gtgggtggtg tagatattaa tgacaactcc cctgagtttg agcaggcttt 900  
 ttatgaggtg aagattctgg agaatagcat ccttggctcc ctggttgtga ccgtctcagc 960  
 ctgggattta gactctggaa caaacagtga actatcctat accttttccc atgcctcaga 1020  
 agatattcgc aagacatttg aaattaatca aaagtctggt gacattactt taacagcacc 1080  
 tttggatttt gaagcaattg agtcatactc aataatcatt caagccacag atgggggagg 1140  
 actttttgga aaatctacag tcagaattca ggtgatggat gtaaacgaca acgctcctga 1200  
 aatcactgtg tcatcaatta ccagtccaat cccagaaaac actccagaga ctgtggttat 1260  
 ggttttcagg atacgagaca gagactctgg ggacaacgga aagatggttt gttctatccc 1320  
 ggaggacatc ccattcgtgc taaaaatctt ggtaaataat tactacactt tggaaacaga 1380  
 gagaccgtg gagagagaga gcagagccga gtacaacatc accatcaccg tcaccgactt 1440  
 ggggaccccc aggtctaaaaa ccgagcacaa cataaccgtg ctggtctccg acgtcaatga 1500  
 caacgcccc gccttcaccc aaacttccta cgcctgttgc gtccgcgaga acaacagccc 1560  
 cgccctgcac atcggcagca tcagcgccac agacagagac tcggggacca acgcccagggt 1620  
 caactactcg ctgctgccgt cccaggaccc gcacctgccc ctgcctccc tggctcctcat 1680  
 caacgcggaac aacggccacc tgtttgccct caggctcgtg gactacgagg ccctgcaggg 1740  
 gttccagttc cgcgtgggcg ccacagacca cggctccccg gctttgagca gcgaggcgct 1800  
 ggtgcgcgtg ctggtgctgg acgccaacga caactcgcct ttcgtgctgt acccgctgca 1860  
 gaacggctcc gcgcccgtga ccgagctggt gccctgggcg gccgagccgg gctacctggt 1920  
 gaccaagggt gtggcggtgg acggtgactc gggccagaac gcctggctgt cgtaccagct 1980  
 gctcaaggcc acggagcccg ggcatttcgg cgtgtgggcg cacaatggcg aggtgcgcac 2040  
 cgccaggctg ctgagcgagc gcgacggcg caagcacagg ctgggtggtgc tggtaaggga 2100  
 caatggcgag cctccgcgct cggccaccgc cacgtgcac gtgctcctgg tggacggctt 2160  
 ctcccagccc tacctgcctc tcccggaggc ggccccggcc caggcccagg ccgactcgct 2220



```

cactgtctac ctggtggtgg cgttggcctc agtgtcgtcg ctcttcctct tctcgggtgct 2280
cctgttcgtg gcggtgcggc tgtgcaggag gagcagggcg gccccggtcg gtcgctgctc 2340
ggtgcctgag ggcccccttc caggacatct ggtggacgtg agtggcaccg ggaccctgtc 2400
ccagagctac cactatgagg tgtgtgtgac tggaggctcc aggtcaaata agttcaaatt 2460
tctgaaacca attatcccca acttcctacc ccagagcaca ggtagtgaag tcgaagaaaa 2520
tccccctatt cagaataatt tgggtttctg ataaagaatg aaaaataaaa cctgtgttta 2580
tgaatacatt tataattagg aacttatcgt gaggtgcctg taaagtagta tttttgatca 2640
cttcaaatac atactcttca agtcaagaaa taaatttctt tacatagaaa aggatacaga 2700
tttagtacca agaacacttc acaaagcagg aaatgtgcat gtgtaatggt ttatgtcaaa 2760
caattatgct taatataaag tctattaagt ggtaagtctt gtttgagata ttttaaattg 2820
ctttccattg ttttcaatat ttactgtgac ttttgtttcc tgagttgatt agaattgctgt 2880
tcgagtatac ctaccctagt ttcagaagca tagattgtag tgtacctttt taaactttat 2940
ttttttaaaa aaagtgtgtt tatgaatcat acactatttt cacactttta atctcagaag 3000
aaacatatgt gacatggtat tttagtaatg accaaataga cggctctaga gattcagtaa 3060
gttcactaag gtccactaac taataagtga caaaactgag catccatcct agatctgcct 3120
gactctaagt cagtgacttt gctccattc cactactgtt ttgtcattgg atatcacctg 3180
gcaagtttct gcctaactaa agagaagaaa agtttttacc gtattcatac tactgttcaa 3240
tctttattta gaaataaact ttatctatga tttcattttc ttataaacca gtaatcttgc 3300
ttttctgggt aaattttcag ctattattac taatgctctg atctgcccac atcttaagta 3360
aaaaacaaaa ttgaaagagc aaaaaaaaaa aaaaaaaaaa aaaaaa 3406

```

&lt;210&gt; 34

&lt;211&gt; 795

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

```

Met Glu Asn Gly Gly Ala Gly Thr Leu Gln Ile Arg Gln Val Leu Leu
  1                      5                      10                      15

Phe Phe Val Leu Leu Gly Met Ser Gln Ala Gly Ser Glu Thr Gly Asn
      20                      25                      30

Phe Leu Val Met Glu Glu Leu Gln Ser Gly Ser Phe Val Gly Asn Leu
      35                      40                      45

Ala Lys Thr Leu Gly Leu Glu Val Ser Glu Leu Ser Ser Arg Gly Ala
      50                      55                      60

Arg Val Val Ser Asn Asp Asn Lys Glu Cys Leu Gln Leu Asp Thr Asn
      65                      70                      75                      80

Thr Gly Asp Leu Leu Leu Arg Glu Met Leu Asp Arg Glu Glu Leu Cys
      85                      90                      95

Gly Ser Asn Glu Pro Cys Val Leu Tyr Phe Gln Val Leu Met Lys Asn
      100                      105                      110

Pro Thr Gln Phe Leu Gln Ile Glu Leu Gln Val Arg Asp Ile Asn Asp
      115                      120                      125

His Ser Pro Val Phe Leu Glu Lys Glu Met Leu Leu Glu Ile Pro Glu
      130                      135                      140

Asn Ser Pro Val Gly Ala Val Phe Leu Leu Glu Ser Ala Lys Asp Leu
      145                      150                      155                      160

Asp Val Gly Ile Asn Ala Val Lys Ser Tyr Thr Ile Asn Pro Asn Ser
      165                      170                      175

```

His Phe His Val Lys Ile Arg Val Asn Pro Asp Asn Arg Lys Tyr Pro  
 180 185 190  
 Glu Leu Val Leu Asp Lys Ala Leu Asp Tyr Glu Glu Arg Pro Glu Leu  
 195 200 205  
 Ser Phe Ile Leu Thr Ala Leu Asp Gly Gly Ser Pro Pro Arg Ser Gly  
 210 215 220  
 Thr Ala Leu Val Arg Val Val Val Val Asp Ile Asn Asp Asn Ser Pro  
 225 230 235 240  
 Glu Phe Glu Gln Ala Phe Tyr Glu Val Lys Ile Leu Glu Asn Ser Ile  
 245 250 255  
 Leu Gly Ser Leu Val Val Thr Val Ser Ala Trp Asp Leu Asp Ser Gly  
 260 265 270  
 Thr Asn Ser Glu Leu Ser Tyr Thr Phe Ser His Ala Ser Glu Asp Ile  
 275 280 285  
 Arg Lys Thr Phe Glu Ile Asn Gln Lys Ser Gly Asp Ile Thr Leu Thr  
 290 295 300  
 Ala Pro Leu Asp Phe Glu Ala Ile Glu Ser Tyr Ser Ile Ile Ile Gln  
 305 310 315 320  
 Ala Thr Asp Gly Gly Gly Leu Phe Gly Lys Ser Thr Val Arg Ile Gln  
 325 330 335  
 Val Met Asp Val Asn Asp Asn Ala Pro Glu Ile Thr Val Ser Ser Ile  
 340 345 350  
 Thr Ser Pro Ile Pro Glu Asn Thr Pro Glu Thr Val Val Met Val Phe  
 355 360 365  
 Arg Ile Arg Asp Arg Asp Ser Gly Asp Asn Gly Lys Met Val Cys Ser  
 370 375 380  
 Ile Pro Glu Asp Ile Pro Phe Val Leu Lys Ser Ser Val Asn Asn Tyr  
 385 390 395 400  
 Tyr Thr Leu Glu Thr Glu Arg Pro Leu Asp Arg Glu Ser Arg Ala Glu  
 405 410 415  
 Tyr Asn Ile Thr Ile Thr Val Thr Asp Leu Gly Thr Pro Arg Leu Lys  
 420 425 430  
 Thr Glu His Asn Ile Thr Val Leu Val Ser Asp Val Asn Asp Asn Ala  
 435 440 445  
 Pro Ala Phe Thr Gln Thr Ser Tyr Ala Leu Phe Val Arg Glu Asn Asn  
 450 455 460  
 Ser Pro Ala Leu His Ile Gly Ser Ile Ser Ala Thr Asp Arg Asp Ser  
 465 470 475 480  
 Gly Thr Asn Ala Gln Val Asn Tyr Ser Leu Leu Pro Ser Gln Asp Pro  
 485 490 495

His Leu Pro Leu Ala Ser Leu Val Ser Ile Asn Ala Asp Asn Gly His  
 500 505 510  
 Leu Phe Ala Leu Arg Ser Leu Asp Tyr Glu Ala Leu Gln Gly Phe Gln  
 515 520 525  
 Phe Arg Val Gly Ala Thr Asp His Gly Ser Pro Ala Leu Ser Ser Glu  
 530 535 540  
 Ala Leu Val Arg Val Leu Val Leu Asp Ala Asn Asp Asn Ser Pro Phe  
 545 550 555 560  
 Val Leu Tyr Pro Leu Gln Asn Gly Ser Ala Pro Cys Thr Glu Leu Val  
 565 570 575  
 Pro Trp Ala Ala Glu Pro Gly Tyr Leu Val Thr Lys Val Val Ala Val  
 580 585 590  
 Asp Gly Asp Ser Gly Gln Asn Ala Trp Leu Ser Tyr Gln Leu Leu Lys  
 595 600 605  
 Ala Thr Glu Pro Gly Leu Phe Gly Val Trp Ala His Asn Gly Glu Val  
 610 615 620  
 Arg Thr Ala Arg Leu Leu Ser Glu Arg Asp Ala Ala Lys His Arg Leu  
 625 630 635 640  
 Val Val Leu Val Lys Asp Asn Gly Glu Pro Pro Arg Ser Ala Thr Ala  
 645 650 655  
 Thr Leu His Val Leu Leu Val Asp Gly Phe Ser Gln Pro Tyr Leu Pro  
 660 665 670  
 Leu Pro Glu Ala Ala Pro Ala Gln Ala Gln Ala Asp Ser Leu Thr Val  
 675 680 685  
 Tyr Leu Val Val Ala Leu Ala Ser Val Ser Ser Leu Phe Leu Phe Ser  
 690 695 700  
 Val Leu Leu Phe Val Ala Val Arg Leu Cys Arg Arg Ser Arg Ala Ala  
 705 710 715 720  
 Pro Val Gly Arg Cys Ser Val Pro Glu Gly Pro Phe Pro Gly His Leu  
 725 730 735  
 Val Asp Val Ser Gly Thr Gly Thr Leu Ser Gln Ser Tyr His Tyr Glu  
 740 745 750  
 Val Cys Val Thr Gly Gly Ser Arg Ser Asn Lys Phe Lys Phe Leu Lys  
 755 760 765  
 Pro Ile Ile Pro Asn Phe Leu Pro Gln Ser Thr Gly Ser Glu Val Glu  
 770 775 780  
 Glu Asn Pro Pro Phe Gln Asn Asn Leu Gly Phe  
 785 790 795

&lt;210&gt; 35

&lt;211&gt; 3809

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 35

```

ggggaccgga gtggggagcg cggcgtggag gtgccaccgg gcgcgggtgg cggagagatc 60
agaagcctct tccccaaagg gagccaacct cagcggggac ccgggctcag ggacgcggcg 120
gcggcgcccg cgactgcagt ggctggacga tggcagcgtc cgccggagcc gggcggtga 180
ttgcagcccc agacagcccg cgctggctgt ggtcgggtgt ggccggcgcg cttgggctct 240
tgacagctgg agtatcagcc ttggaagtat atacgccaaa agaaatcttc gtggcaaatg 300
gtacacaaag gaagctgacc tgcaagttca agtctactag tacgactggc ggggtgacct 360
cagtctcctg gagcttccag ccagaggggg ccgacactac tgtgtcgttt ttccactact 420
cccaagggca agtgtacctt gggaattatc caccatttaa agacagaatc agctgggctg 480
gagaccttga caagaaagat gcatcāatca acatagaaaa tatgcagttt atacacaatg 540
gcacctatat ctgtgatgtc aaaaaccctc ctgacatcgt tgtccagcct ggacacatta 600
ggctctatgt cgtagaaaaa gagaatttgc ctgtgttttc agtttgggta gtgggtgggca 660
tagttactgc tgtggtccta ggtctcactc tgctcatcag catgattctg gctgtcctct 720
atagaaggaa aaactctaaa cgggattaca ctggggccca gtcataatg cacagttaga 780
ccactccggc ggacatcaca gtgacaagat taacaagtca gagtctgtgg tgtatgcgga 840
tatccgaaag aattaagaga atacctagaa catatcctca gcaagaaaca aaaccaaaact 900
ggactctcgt gcagaaaatg tagcccata ccacatgtag ccttggagac ccaggcaagg 960
acaagtacac gtgtactcac agaggagag aaagatgtgt acaaaggata tgtataaata 1020
ttctatttag tcatcctgat atgaggagcc agtggtgcat gatgaaaaga tggtagtatt 1080
ctacatatgt acccattgtc ttgctgtttt tgtactttct tttcaggta tttacaattg 1140
ggagatttca gaaacattcc ttccaccatc atttagaaat ggtttgcctt aatggagaca 1200
atagcagatc ctgtagtatt tccagtagac atggcctttt aatctaaggg cttaagactg 1260
attagtctta gcatttactg tagttggagg atggagatgc tatgatggaa gcatacccag 1320
ggtggccttt agcacagtat cagtaccatt tatttgtctg ccgcttttaa aaaataacca 1380
ttggctatgc cacttgaaaa caatttgaga agtttttttg aagtttttct cactaaaaata 1440
tggggcaatt gttagcctta catgttgtgt agacttactt taagtttgca cccttgaaat 1500
gtgtcatatc aatttcttga ttcataatag caagatttagc aaaggataaa tgccgaaggt 1560
cacttcattc tggacacagt tggatcaata ctgattaagt agaaaatcca agctttgctt 1620
gagaactttt gtaacgtgga gagtaaaaag tatcggtttt attcttttgc gatgtccttt 1680
ctgcttgaaa taacagtcac catacagcta aaggagagga gtttcttttc ttctaagtag 1740
gcagaaatgg tatcattatg ttgccgtctt ccaatctccc agagctcgtc ctctagagaa 1800
tcaccttctt tcgctttttt ttttttttga ggtagagtct cactatgttg cccagactag 1860
ccttgaactc ttgggctcaa gtgattctcc ctccctcagcc tcccagtag ctggaacgaa 1920
ctatagttag accactgcag ctggcaagaa tcaccttttt tataaagcgt cagtcatgct 1980
tccagcaaga ggcagatca gtcattggct tataacagct tcattggtgc tcaaagactg 2040
ttgagggtta tgagagccta gattagacag tttggctgtc cttccctaaa acttggtttc 2100
tcctattcac tactccccac cgcacttaaa atctatgagt ttttactttt tactgggaat 2160
ggaaagtgtg gtgaagatca ttcaaacact atgttgtcat ttctccatt ttctgaattt 2220
ttttttaaatt tccccccctt ttaaaattgt tcgaaagccc acagttatgg aaagaattac 2280
tgtctagatg gtctgcagaa cgtgtttggg gtgagtggga gtgaggggca atgttacttt 2340
ttctccttgt agtttggagt ccattatgag ctgctgcttt ttctctcat cttgtcatct 2400
tctggggatg tttgaaggct gagttccaac agaattcaca aagggaataa aacaggattg 2460
agatttttag gtgtgcacaa ggtggtaaga taaagggcat atgagcttca aaactaatgc 2520
tgttgcatac atgaagcctt ttgttttttg aggagctatt tttgttattc ttgtaacgct 2580
ccaccttaca tgccacgtct gtgtgagtca acagggatca ggtttggtca ccacacatgt 2640
ctgaagctgg gcagcgtctg ctctgtgttc tgtgtggaat ggagaaaaaa acgctgccc 2700
tgctgccttc catgttcata ggcccagccc aagagagtga cacacagtgc tggccctgag 2760
acatttccac aaagtgttca actctgcctt gcatcctaaa actttttggg catctatttt 2820
gaaaactata ggagcctttg gaaggcctct tatgtttgga ggggaagggt gttgagattg 2880
tcaccatcct tcaagctgag actcctgggt agcctttgcc accatgaaaa ccacatagct 2940
gaccagggct gtgcttgagg tacagaggac acacatcgta gacaggcctg tgtcatgttt 3000
ccttacagtc gttttttaca gagaaaagg gcatgtttt ttactgctt tctcaacagt 3060
tcctgtgaat aaatgaaaca ttctggagct cctgagagc aagagccttc acttctctct 3120
gcggtgccgg gaccatgtgt tgggtgaagct ggtgctgtgg gggccactca ctgaaatgac 3180
acctggaggc ctgttcctcc cttaccactc ccttcccag cccgacttct tggcctcctg 3240
cccaaccaga cactcaaac tctgtcagcg ccttggcatt ctggcagaga atcctcacca 3300
gttctcacca accttcccc caggcaaggg cagctgccag catggtgtct tgccaggaca 3360

```

```

ggtttccttg aaggaagctg ctcacactga gatgagcctc tcagggcagg acctcttccc 3420
aagccctgca caccaccccc tgcagccctt ttggctcccc ttttccctgt gcctcagcac 3480
tcctttcctg gttgcagata acgaactaag gttgcctaaa gggcagatct gccctctcca 3540
tgtcttcgtc ctggcaaaaca gggctcgtctt aaaattatgc gctaattctg tatgggagca 3600
ctcaaaaggc attacttaga gattgaaatt tcaaactatc tctagttttt caatggaaat 3660
atatcagcta gggaaaaacc atcaagctca ttattatttt ttgatcttca gttgtatttt 3720
tgtgaatatt ttaatacatc tttttcaatt tctgaaaaaa aaaaaaaaaa aaaaaaaaaa 3780
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa

```

3809

&lt;210&gt; 36

&lt;211&gt; 209

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 36

```

Met Ala Ala Ser Ala Gly Ala Gly Ala Val Ile Ala Ala Pro Asp Ser
  1             5             10             15

Arg Arg Trp Leu Trp Ser Val Leu Ala Ala Ala Leu Gly Leu Leu Thr
      20             25             30

Ala Gly Val Ser Ala Leu Glu Val Tyr Thr Pro Lys Glu Ile Phe Val
      35             40             45

Ala Asn Gly Thr Gln Gly Lys Leu Thr Cys Lys Phe Lys Ser Thr Ser
      50             55             60

Thr Thr Gly Gly Leu Thr Ser Val Ser Trp Ser Phe Gln Pro Glu Gly
      65             70             75             80

Ala Asp Thr Thr Val Ser Phe Phe His Tyr Ser Gln Gly Gln Val Tyr
      85             90             95

Leu Gly Asn Tyr Pro Pro Phe Lys Asp Arg Ile Ser Trp Ala Gly Asp
      100            105            110

Leu Asp Lys Lys Asp Ala Ser Ile Asn Ile Glu Asn Met Gln Phe Ile
      115            120            125

His Asn Gly Thr Tyr Ile Cys Asp Val Lys Asn Pro Pro Asp Ile Val
      130            135            140

Val Gln Pro Gly His Ile Arg Leu Tyr Val Val Glu Lys Glu Asn Leu
      145            150            155            160

Pro Val Phe Pro Val Trp Val Val Val Gly Ile Val Thr Ala Val Val
      165            170            175

Leu Gly Leu Thr Leu Leu Ile Ser Met Ile Leu Ala Val Leu Tyr Arg
      180            185            190

Arg Lys Asn Ser Lys Arg Asp Tyr Thr Gly Ala Gln Ser Tyr Met His
      195            200            205

Ser

```

&lt;210&gt; 37

&lt;211&gt; 1954

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 37

```

gagacttggg ctggagccgc cctgggtgtc agcgggctcg gctcccgcgc acgctccggc 60
cgtcgcgcag cctcggcacc tgcaggtccg tgcgtcccgc ggctggcgcc cctgactccg 120
tcccggccag ggagggccat gatttccctc ccggggcccc tggtgaccaa cttgctgcgg 180
tttttgttcc tggggctgag tgcctcgcgc ccccccctcg gggcccagct gcaactgcac 240
ttgcccgcga accggttgca ggcggtggag ggaggggaag tggtgcttcc agcgtggtac 300
accttgcacg gggaggtgtc ttcattcccag ccatgggagg tgccttttgt gatgtggttc 360
ttcaaacaga aagaaaagga ggatcagggt ttgtcctaca tcaatggggt cacaacaagc 420
aaacctggag tatccttggg ctactccatg ccctcccga acctgtccct gcggctggag 480
ggtctccagg agaaagactc tggcccctac agctgctccg tgaatgtgca agacaaacaa 540
ggcaaatcta gggggcacag catcaaaacc ttagaactca atgtactggg tcctccagct 600
cctccatcct gccgtctcca ggggtgtgcc catgtggggg caaacgtgac cctgagctgc 660
cagttctcaa ggagtaagcc cgctgtccaa taccagtggg atcggcagct tccatccttc 720
cagactttct ttgcaccagc attagatgtc atccgtgggt ctttaagcct caccaacctt 780
tcgtcttcca tggctggagt ctatgtctgc aagggccaca atgaggtggg cactgcccac 840
tgtaattgta cgctggaagt gagcacaggg cctggagctg cagtgggtgc tggagctgtt 900
gtgggtaccc tgggtggact ggggttgcgt gctgggctgg tcctcttcta ccaccgccc 960
ggcaaggccc tggaggagcc agccaatgat atcaaggagg atgccattgc tccccggacc 1020
ctgcccctggc ccaagagctc agacacaatc tccaagaatg ggacccttcc ctctgtcacc 1080
tccgcacgag ccctccggcc accccatggc cctcccaggc ctgggtgcatt gacccccacg 1140
cccagtcctc ccagccaggc cctgccctca ccaagactgc ccacgacaga tggggcccac 1200
cctcaaccaa tatcccccat ccctgggtggg gtttcttctc ctggcttgag ccgcatgggt 1260
gctgtgcctg tgatgggtgcc tgcctcagag caagctggct ctctgggtatg atgacccac 1320
cactcattgg cttaaaggatt tgggtctctc ccttctata agggtcacct ctacgacaga 1380
ggcctgagtc atgggaaaga gtcacactcc tgacccttag tactctgccc ccacctctc 1440
ttactgtggg aaaaccatct cagtaagacc taagtgtcca ggagacagaa ggagaagagg 1500
aagtggatct ggaattggga ggagcctcca cccaccctg actcctcctt atgaagccag 1560
ctgctgaaat tagctactca ccaagagtga ggggcagaga cttccagtc ctgagtctcc 1620
caggccccct tgatctgtac cccaccctca tctaaccaca cccttggtc ccactccagc 1680
tcctgttatt gatataacct gtcaggctgg cttgggtagg ttttactggg gcagaggata 1740
gggaatctct tattaact aacatgaaat atgtgtgtt ttcatttgca aatttaaata 1800
aagatacata atgtttgtat gaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1860
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1920
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa                                     1954

```

&lt;210&gt; 38

&lt;211&gt; 390

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 38

```

Met Ile Ser Leu Pro Gly Pro Leu Val Thr Asn Leu Leu Arg Phe Leu
  1                   5                   10                   15

Phe Leu Gly Leu Ser Ala Leu Ala Pro Pro Ser Arg Ala Gln Leu Gln
      20                   25                   30

Leu His Leu Pro Ala Asn Arg Leu Gln Ala Val Glu Gly Gly Glu Val
      35                   40                   45

Val Leu Pro Ala Trp Tyr Thr Leu His Gly Glu Val Ser Ser Ser Gln
      50                   55                   60

Pro Trp Glu Val Pro Phe Val Met Trp Phe Phe Lys Gln Lys Glu Lys
      65                   70                   75                   80

Glu Asp Gln Val Leu Ser Tyr Ile Asn Gly Val Thr Thr Ser Lys Pro

```

85					90					95					
Gly	Val	Ser	Leu	Val	Tyr	Ser	Met	Pro	Ser	Arg	Asn	Leu	Ser	Leu	Arg
			100					105					110		
Leu	Glu	Gly	Leu	Gln	Glu	Lys	Asp	Ser	Gly	Pro	Tyr	Ser	Cys	Ser	Val
		115					120					125			
Asn	Val	Gln	Asp	Lys	Gln	Gly	Lys	Ser	Arg	Gly	His	Ser	Ile	Lys	Thr
	130						135					140			
Leu	Glu	Leu	Asn	Val	Leu	Val	Pro	Pro	Ala	Pro	Pro	Ser	Cys	Arg	Leu
	145					150					155				160
Gln	Gly	Val	Pro	His	Val	Gly	Ala	Asn	Val	Thr	Leu	Ser	Cys	Gln	Ser
				165					170					175	
Pro	Arg	Ser	Lys	Pro	Ala	Val	Gln	Tyr	Gln	Trp	Asp	Arg	Gln	Leu	Pro
			180					185					190		
Ser	Phe	Gln	Thr	Phe	Phe	Ala	Pro	Ala	Leu	Asp	Val	Ile	Arg	Gly	Ser
		195					200					205			
Leu	Ser	Leu	Thr	Asn	Leu	Ser	Ser	Ser	Met	Ala	Gly	Val	Tyr	Val	Cys
	210						215				220				
Lys	Ala	His	Asn	Glu	Val	Gly	Thr	Ala	Gln	Cys	Asn	Val	Thr	Leu	Glu
	225					230					235				240
Val	Ser	Thr	Gly	Pro	Gly	Ala	Ala	Val	Val	Ala	Gly	Ala	Val	Val	Gly
				245					250					255	
Thr	Leu	Val	Gly	Leu	Gly	Leu	Leu	Ala	Gly	Leu	Val	Leu	Leu	Tyr	His
		260						265						270	
Arg	Arg	Gly	Lys	Ala	Leu	Glu	Glu	Pro	Ala	Asn	Asp	Ile	Lys	Glu	Asp
		275					280					285			
Ala	Ile	Ala	Pro	Arg	Thr	Leu	Pro	Trp	Pro	Lys	Ser	Ser	Asp	Thr	Ile
	290					295						300			
Ser	Lys	Asn	Gly	Thr	Leu	Ser	Ser	Val	Thr	Ser	Ala	Arg	Ala	Leu	Arg
	305					310					315			320	
Pro	Pro	His	Gly	Pro	Pro	Arg	Pro	Gly	Ala	Leu	Thr	Pro	Thr	Pro	Ser
				325					330					335	
Leu	Ser	Ser	Gln	Ala	Leu	Pro	Ser	Pro	Arg	Leu	Pro	Thr	Thr	Asp	Gly
			340					345						350	
Ala	His	Pro	Gln	Pro	Ile	Ser	Pro	Ile	Pro	Gly	Gly	Val	Ser	Ser	Ser
	355						360					365			
Gly	Leu	Ser	Arg	Met	Gly	Ala	Val	Pro	Val	Met	Val	Pro	Ala	Gln	Ser
	370					375					380				
Gln	Ala	Gly	Ser	Leu	Val										
	385						390								

<210> 39  
 <211> 1933  
 <212> DNA  
 <213> Homo sapiens

<400> 39  
 ggggtggggc caggaggaag atggcggcgt cgcagctgc cgctgagctc caggcttctg 60  
 ggggtccgcg gcacccagtg tgtctgttgg tgttgggaat ggcgggatcc gggaaaacca 120  
 cttttgtaca gaggctcaca ggacacctgc atgcccagg cactccaccg tatgtgatca 180  
 acctggatcc agcagtacat gaagtccctt ttcttgccaa tattgatatt cgtgatactg 240  
 taaagtataa agaagtaatg aaacaatatg gacttggacc caatggcggc atagtgcct 300  
 cactcaatct ctttgtctacc agatttgatc aggtgatgaa atttattgag aaggcccaga 360  
 acatgtccaa atatgtgttg attgacacac ctggacagat tgaggatttc acctggtcag 420  
 cttctgggac aattatcact gaagcccttg catcctcatt tccaacagtt gtcattctatg 480  
 taatggacac atcgagaagt accaaccctg tgaccttcat gtccaacatg ctctatgcct 540  
 gcagcatctt atacaaaacc aagctgcctt tcattgtggt catgaataaa actgacatca 600  
 ttgaccacag ctttgcagtg gaatggatgc aggatgttga ggctttccaa gatgccttga 660  
 atcaagagac tacatacgtc agtaacctga ctctgtcaat gagcctggtg ttagatgagt 720  
 tttacagctc actcaggggtg gtgggtgtct ctgctgttct gggtagctga ttagatgaac 780  
 tctttgtgca agttaccagt gctgcccgaag aatatgaaag ggagtatcgt cctgaatatg 840  
 aacgtctgaa aaaatcactg gccaacgcag agagccaaca gcagagagaa caactggaac 900  
 gccttcgaaa agatatgggt tctgtagcct tggatgcagg gactgccaaa gacagcttat 960  
 ctctgtgtct gcaccttctt gatttgatcc tgactcgagg aaccttggat gaagaggatg 1020  
 aggaagcaga cagcgatact gatgacattg accacagagt tacagaggaa agccatgaag 1080  
 agccagcatt ccagaatttt atgcaagaat cgatggcaca atactggaag agaaacaata 1140  
 aataggagac tttagcacac ttacttgtt tctagaagtc cagaattttg gacctccacg 1200  
 tgaagaagact gttcttacct ctgaactggg ggctcccata agggataatt ttctcagag 1260  
 tagcaaaagt tctcttatta gagaaatctt gtgactcaga tgaagtcagg gatagaagac 1320  
 ccttggacct ggcaggttaa tgctgattat tccttggcct ttcccttgta tttatgcaag 1380  
 gaaggatata ctgagctgat actcttccaa gcctacaact tcaagtttta tcatttgaac 1440  
 tcaagtactt ttgctgctga ggaatggaat caaaagaacg tagtctcctg gtaaccacct 1500  
 cagatctcta ttattaggct agatgtatag cctctactcc cccagcttct tgctcttgac 1560  
 cctgcaactg aagttgccct tctattagca gccaaagaaa agggaaacat gagcttatcc 1620  
 agaacggttg cagagtctcc ttggcaatca accaacgttg ctatgaaata tgcctcacac 1680  
 tgtatagctc attataggac gtcagggttg ttgaaaaaag tgggcaagac atgattaatg 1740  
 aatcagaatc ctgtttcatt ggtgacttgg ataaagactt tttaatttta actttgaaaa 1800  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1860  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1920  
 aaaaaaaaaa aaa 1933

<210> 40  
 <211> 374  
 <212> PRT  
 <213> Homo sapiens

<400> 40  
 Met Ala Ala Ser Ala Ala Ala Glu Leu Gln Ala Ser Gly Gly Pro  
 1 5 10 15  
 Arg His Pro Val Cys Leu Leu Val Leu Gly Met Ala Gly Ser Gly Lys  
 20 25 30  
 Thr Thr Phe Val Gln Arg Leu Thr Gly His Leu His Ala Gln Gly Thr  
 35 40 45  
 Pro Pro Tyr Val Ile Asn Leu Asp Pro Ala Val His Glu Val Pro Phe  
 50 55 60  
 Pro Ala Asn Ile Asp Ile Arg Asp Thr Val Lys Tyr Lys Glu Val Met  
 65 70 75 80



Lys Gln Tyr Gly Leu Gly Pro Asn Gly Gly Ile Val Thr Ser Leu Asn  
                                     85                                    90                                    95  
 Leu Phe Ala Thr Arg Phe Asp Gln Val Met Lys Phe Ile Glu Lys Ala  
                                     100                                    105                                    110  
 Gln Asn Met Ser Lys Tyr Val Leu Ile Asp Thr Pro Gly Gln Ile Glu  
                                     115                                    120                                    125  
 Val Phe Thr Trp Ser Ala Ser Gly Thr Ile Ile Thr Glu Ala Leu Ala  
                                     130                                    135                                    140  
 Ser Ser Phe Pro Thr Val Val Ile Tyr Val Met Asp Thr Ser Arg Ser  
                                     145                                    150                                    155                                    160  
 Thr Asn Pro Val Thr Phe Met Ser Asn Met Leu Tyr Ala Cys Ser Ile  
                                     165                                    170                                    175  
 Leu Tyr Lys Thr Lys Leu Pro Phe Ile Val Val Met Asn Lys Thr Asp  
                                     180                                    185                                    190  
 Ile Ile Asp His Ser Phe Ala Val Glu Trp Met Gln Asp Phe Glu Ala  
                                     195                                    200                                    205  
 Phe Gln Asp Ala Leu Asn Gln Glu Thr Thr Tyr Val Ser Asn Leu Thr  
                                     210                                    215                                    220  
 Arg Ser Met Ser Leu Val Leu Asp Glu Phe Tyr Ser Ser Leu Arg Val  
                                     225                                    230                                    235                                    240  
 Val Gly Val Ser Ala Val Leu Gly Thr Gly Leu Asp Glu Leu Phe Val  
                                     245                                    250                                    255  
 Gln Val Thr Ser Ala Ala Glu Glu Tyr Glu Arg Glu Tyr Arg Pro Glu  
                                     260                                    265                                    270  
 Tyr Glu Arg Leu Lys Lys Ser Leu Ala Asn Ala Glu Ser Gln Gln Gln  
                                     275                                    280                                    285  
 Arg Glu Gln Leu Glu Arg Leu Arg Lys Asp Met Gly Ser Val Ala Leu  
                                     290                                    295                                    300  
 Asp Ala Gly Thr Ala Lys Asp Ser Leu Ser Pro Val Leu His Pro Ser  
                                     305                                    310                                    315                                    320  
 Asp Leu Ile Leu Thr Arg Gly Thr Leu Asp Glu Glu Asp Glu Glu Ala  
                                     325                                    330                                    335  
 Asp Ser Asp Thr Asp Asp Ile Asp His Arg Val Thr Glu Glu Ser His  
                                     340                                    345                                    350  
 Glu Glu Pro Ala Phe Gln Asn Phe Met Gln Glu Ser Met Ala Gln Tyr  
                                     355                                    360                                    365  
 Trp Lys Arg Asn Asn Lys  
                                     370

&lt;210&gt; 41

<211> 2795  
 <212> DNA  
 <213> Homo sapiens

<400> 41

```

aaaaagctcg gcatggctga cgacgcgggt ttggagaccc cgctgtgttc cgagcagttc 60
ggctccgggg aggcacgggg ctgccgcgcc gccgcggacg ggagcctgca gtgggaggtc 120
gggggctggc gctgggtggg gctctccagg gccttcacgg tcaaacctga aggacgagat 180
gcggggcgaag tgggggcttc cggggccccc tcaccgcccc tctccgggct ccaggccctg 240
ttcctgcctc agggcttccc tgatagcgtc agcccggact acttgcccta ccagctgttg 300
gattccgtgc aggcgtttgc ttccagcctc tccggctccc tagccacca ggagctctt 360
ctgggcatag ggggtgggaa cgcaaaagcc actgtttcag ctgccacggc cactggctc 420
gtgaaagatt caactggcat gctgggcccgc atcgtctttg cctggtggaa agggagcaaa 480
ctggactgca atgccaagca gtggaggctt ttgctggaca tcctcaatga cgtagccatg 540
ttccttgaga ttatggctcc tgtataccca atctgtttca ccatgaccgt ctccaccagc 600
aacctagcca agtgcacgtc gagggttgc ggtggggcca ctccggctgc cctgaccgtg 660
caccaggtgc ggaggaacaa catggctgac gtgtcagcca aggacagcag ccaggagacg 720
ctggtgaacc tggcggggct cttggtcagc ctctgatgc tccctctggt gtcagggttc 780
cctggcttca gccttgatg ttctcttctc ctactgccc tccacatcta cgccaactac 840
cgcgcggtcc gagccctggt catggagacc ttgaacgaag gccggctccg gctggctctg 900
aagcactacc ttcagagggg agaggtactc gaccacactg cagccaatcg catggagccg 960
ctgtggacag gtttctggcc agctccgtct ctatccctgg gggctcccctt acaccgctt 1020
gtctccagtg tctttgagct gcagcagctg gttgaggggc accaagaatc ctacctctc 1080
tgctgggacc agtcacaaaa ccaggtagc gtagttctga accagaaggc agggcccaag 1140
accatcctaa gggccgccc acatgggctg atgcttgggg ccctgcaggg agatggaccc 1200
ctccacagc agctggagga gctgaggaac cgggtgcggg caggctcctaa gaaagagagc 1260
tgggtcgtcg tcaaggagac acacgaagtg ttggacatgc tgttcccaaa gttcttga 1320
ggactgcagg atgccggctg gaagaccgag aagcaccagc tagaggtgga tgagtggag 1380
gccacatggc ttctgtctcc cgaaaagaag gtcttgtgag cagcccagac ggaggcccaa 1440
gcccagggca ggaacctgga gcaaggacac ttgggccaca gcaggacagg ggaaggcag 1500
ctttattttt ccttagggca actgcagcgg gtgggccagg ccctcatggg aagtgactgc 1560
caatcagatg cagtggccc caggcagagg aaggccggga gaaggggagc caggacctc 1620
tcaccccact gccccttccc cttttctggg gagcaccgca ggctctcac ccccacttc 1680
tgtgaggctg tggcttatgg tgtccaacgc agttggtctt aggcatagaa gcccagagg 1740
aacacggcca ctgccatcat gagcagggca ttgaggttga ccacacggg ccagctcggg 1800
tcctcgctga tgtctccag ccgctgggt gctgcgctg cctcctctg ggtaggggc 1860
ggaggactgc ccaccccacc tctgtcatt ccacaaaacc agagcaggca ctggcggaa 1920
aggcttggtg ccggggcctg gggctctgag ggaaattgag gccctgcagt tagtttgcg 1980
gaactcagct cctccagccc cactctccag catggtgccc taccattcat ctccatggc 2040
ctctctgggc acccattctg tacaggagat gaggagcctt gctgctcatc agcatccag 2100
tcctccggtt cctccttctg atgccggaga ctgaagacca gccgggtggg ctggggagg 2160
tgggagcacg aacgaggtg gagttctgtc ccccatgcc tggccctaaa gtctcttgca 2220
caccagctcg tcactgectg cctacccac ctctgtccag tctacacacc cagcccagg 2280
ttaactcatg ccaactccac cctacatggc tgcctgtgc cctcgggata aacccaagc 2340
ccctgagctt gtgtttaaag ccgttggcct tgctcccca gctttgtcag ctcaggctc 2400
tctacacca gatgtagcg cttgtgacac tggcctggca gtctgtcga cagtgtctc 2460
tgctgtgtg ctcccacct ttctcctgc tgetgcagaa acccggccat ccttccacac 2520
ccagatctct tgtctgtcc tcaccccacc ctgccaccat cagccctgcc tggagccacc 2580
tgccctttg caacaaaacc aaacctttt gtgggcgttc aagatggat tgtgccacc 2640
agtcagatcc tgtgttttga gtcccaaagg ccatgccaa gattggctt gggaggctt 2700
aatcaccaac ccatcaacat caagcctccc ccaggccgg tcaaataaat gtatttaa 2760
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 2795

```

<210> 42  
 <211> 468  
 <212> PRT  
 <213> Homo sapiens

<400> 42

Met Ala Asp Asp Ala Gly Leu Glu Thr Pro Leu Cys Ser Glu Gln Phe

1	5	10	15
Gly Ser Gly Glu Ala Arg Gly Cys Arg Ala Ala Ala Asp Gly Ser Leu	20	25	30
Gln Trp Glu Val Gly Gly Trp Arg Trp Trp Gly Leu Ser Arg Ala Phe	35	40	45
Thr Val Lys Pro Glu Gly Arg Asp Ala Gly Glu Val Gly Ala Ser Gly	50	55	60
Ala Pro Ser Pro Pro Leu Ser Gly Leu Gln Ala Val Phe Leu Pro Gln	65	70	75
Gly Phe Pro Asp Ser Val Ser Pro Asp Tyr Leu Pro Tyr Gln Leu Trp	85	90	95
Asp Ser Val Gln Ala Phe Ala Ser Ser Leu Ser Gly Ser Leu Ala Thr	100	105	110
Gln Ala Val Leu Leu Gly Ile Gly Val Gly Asn Ala Lys Ala Thr Val	115	120	125
Ser Ala Ala Thr Ala Thr Trp Leu Val Lys Asp Ser Thr Gly Met Leu	130	135	140
Gly Arg Ile Val Phe Ala Trp Trp Lys Gly Ser Lys Leu Asp Cys Asn	145	150	155
Ala Lys Gln Trp Arg Leu Phe Ala Asp Ile Leu Asn Asp Val Ala Met	165	170	175
Phe Leu Glu Ile Met Ala Pro Val Tyr Pro Ile Cys Phe Thr Met Thr	180	185	190
Val Ser Thr Ser Asn Leu Ala Lys Cys Ile Val Ser Val Ala Gly Gly	195	200	205
Ala Thr Arg Ala Ala Leu Thr Val His Gln Ala Arg Arg Asn Asn Met	210	215	220
Ala Asp Val Ser Ala Lys Asp Ser Ser Gln Glu Thr Leu Val Asn Leu	225	230	235
Ala Gly Leu Leu Val Ser Leu Leu Met Leu Pro Leu Val Ser Gly Cys	245	250	255
Pro Gly Phe Ser Leu Gly Cys Phe Phe Phe Leu Thr Ala Leu His Ile	260	265	270
Tyr Ala Asn Tyr Arg Ala Val Arg Ala Leu Val Met Glu Thr Leu Asn	275	280	285
Glu Gly Arg Leu Arg Leu Val Leu Lys His Tyr Leu Gln Arg Gly Glu	290	295	300
Val Leu Asp Pro Thr Ala Ala Asn Arg Met Glu Pro Leu Trp Thr Gly	305	310	315
Phe Trp Pro Ala Pro Ser Leu Ser Leu Gly Val Pro Leu His Arg Leu			320

```
<210> 43
<211> 2980
<212> DNA
<213> Homo sapiens
```

<400> 43							
aagaaaaact	tttttgttat	tgggctttcc	aggtggagtt	cagaaccagt	gactcacact	60	
tctcagtcct	gggagcaatt	tatttgtctac	ttggagggggt	tgtaagaaaa	gccagtgaga	120	
aagcagatac	ccccacaaac	acagatccac	tgttgagcccc	caaaacctgt	cctgtccccc	180	
tcttttaaga	ctccagccac	ccctcttggg	ctctctactt	ccacggggca	catgctgatg	240	
ccctgtgtgt	ggctgctctg	gtggtgggtgg	tgctgctgct	ccggctggta	ctgctatgga	300	
ttgtgtgtccc	cagcccccca	gatgttgctc	caccaggggtc	tcttcaagtg	ccgctgcccgc	360	
atgctcttca	atgacctgaa	ggttttctta	ctgctggcgcc	ctcttcaagc	gccctgtccc	420	
atgcacggcg	accctcagac	ccccggtttg	cgggccaata	acacctctcc	ggctctgggc	480	
gccgggggggt	gggcaggctg	gagggggcccc	cgagagtagtg	tgggcaggga	gccccctctc	540	
gtgccacctc	cacccccctt	gccaccttct	tctgtggaag	atgactgggg	tggcccagcc	600	
acagagccac	ctgcctcgct	gctcagcagt	gcctcctcag	atgacttctg	taaggagaag	660	
accgaggatc	gctactcact	gggcagcagc	ttggacagtg	gtatgaggac	cccactctgc	720	
cgcactctgct	tccagggggc	agaacagggg	gagctgctga	gcccattgcc	ctgtgatggc	780	
tccgtcaagt	gcacacacca	gccttgcctc	atacagttga	tcagcgagcg	gggtctgtgc	840	
agctgcgagc	tgtgctacta	caagtaccac	gtcatcgcca	taagcacaaa	aaatcctctg	900	
cagtggcagg	ccatctctct	gacgggtcatt	gagaagggttc	aggttgcgac	cgcatctctg	960	
ggctccctct	tctctcatgc	cagtattttc	tggctcatct	ggtaacctct	cagccccctg	1020	
gcagaatggc	agcgccaaga	cttctctctc	catagctgct	acgggatgta	tggcttcatg	1080	
gacgtgggtg	gcataggttc	catcatccat	gaaggacctc	cggtgtaccg	catcttataa	1140	
cgggtggcagg	ctgtcaacca	gcagtggaag	gtgctgaact	atgacaagac	aaaagacctg	1200	
gaggatcaaa	aggcaggagg	caggaccaac	ccccggacct	cctcatccac	ccaggccaat	1260	
atccccctct	cggagaggga	cagccgcaggc	accctgtccc	ctgagcagag	ccctgtcccag	1320	
gctcccgccc	accctcaggc	ccctctgttc	catcaccact	gtgcttatac	catctctgac	1380	
atcctgagtc	acttgagacc	tcatgaacag	cgaagctccc	caggcagcac	ccgaagactg	1440	

```

gtcatgagag tcacgacagt gtgagagcag aggccccgaa ggaaggccat gaccaccact 1500
gagggcccg agcaggtggt ggaggtgcag tggcaccccc ggagccaaca gagggagcag 1560
gcagagggtg ggggacctgg cgggagccct ggggtagtgt cagagcggga gtgaggctgg 1620
tgcaggagca gttctgctat ttccaatcag tcaatgccac tctccacaac aacaatgaaa 1680
accaacacca actcaacaac aaagtgcaat acaggtgaa cctggcccaa cagaaaaacc 1740
ctgccccaat gcacctgcag gcaaggtacc cgaagaagca gaggtgagg gcaggcaaaag 1800
cctgtgtgac tgtggcagt cgggaggcca agggggccaa gaggaagc atctgtggtc 1860
tgcctgtctc tcacctgtt tggttttgtt tctcctggg ctgtgttctg caggcagcca 1920
gaaaaggagg aggcacgggt gagctggcag ggacacactg cctttggggc tcctgggctc 1980
atttgatgg gcaagattcg ctgacaaatg gctgtgggga tgggtgggtg gatggtcagg 2040
gagggatgct cagggagggg tatgctggtg tgagcagcca gagggagagt gtgtctcctt 2100
cctgaaggaa cttccaaatg gaactcccga ttccaggtgg gctaaaagag ggcttaggtt 2160
tggaagaggg tgtccttctg tgccttgtt aatttatttt atagtattt ggttcaaaga 2220
tgtttacagg acacacacac acacacacac acacacacac ccctagagaa 2280
aagtacagat ttccagtggg tatttcaagc acagttctgc tgcgtgggtc tcagctttgg 2340
aagctgtcaa tcccggagca actttcccaa ctaccaacc ccaccatggc caggacatgt 2400
gcaatgccag ccccttcttg tcttggcaca tgcacagacc cagtcccctc acggtagggc 2460
acccctgacc tacgggcttc caagagagca gctgcagtgg ttgggaggag cttgaccagt 2520
gtgccccaa gaggtagta gagccaatc taagtattcc ttgctgctg gaacctccc 2580
tgtttggaac cctcccaaaa gaggcagtca ggctgatgct cagtgccttg tgctccctgc 2640
tccttcccgc gtagccaggt gggcccaagg gtgcctggca gggagcacta cccctggacc 2700
cctcctgctc gctctgggga ccctgccagg gaaggccact ggtgtttcac ctgcaaagt 2760
tctggtgtc actgcacagt ggtcgcgtca tccatgggta ttaaaaggac actgtcaagt 2820
acttttttaa actagttttt agggttttt aaaactctct gttgtttgta atattctctt 2880
aaaagcttga aaataaaact tctttcccta ccaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2940
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2980

```

&lt;210&gt; 44

&lt;211&gt; 410

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 44

```

Met Leu Met Pro Leu Cys Gly Leu Leu Trp Trp Trp Trp Cys Cys Cys
  1             5             10             15

```

```

Ser Gly Trp Tyr Cys Tyr Gly Leu Cys Ala Pro Ala Pro Gln Met Leu
      20             25             30

```

```

Arg His Gln Gly Leu Leu Lys Cys Arg Cys Arg Met Leu Phe Asn Asp
      35             40             45

```

```

Leu Lys Val Phe Leu Leu Arg Arg Pro Pro Gln Ala Pro Leu Pro Met
      50             55             60

```

```

His Gly Asp Pro Gln Pro Pro Gly Leu Ala Ala Asn Asn Thr Leu Pro
      65             70             75             80

```

```

Ala Leu Gly Ala Gly Gly Trp Ala Gly Trp Arg Gly Pro Arg Glu Val
      85             90             95

```

```

Val Gly Arg Glu Pro Pro Pro Val Pro Pro Pro Pro Pro Leu Pro Pro
      100            105            110

```

```

Ser Ser Val Glu Asp Asp Trp Gly Gly Pro Ala Thr Glu Pro Pro Ala
      115            120            125

```

```

Ser Leu Leu Ser Ser Ala Ser Ser Asp Asp Phe Cys Lys Glu Lys Thr
      130            135            140

```

Glu Asp Arg Tyr Ser Leu Gly Ser Ser Leu Asp Ser Gly Met Arg Thr  
 145 150 155 160  
 Pro Leu Cys Arg Ile Cys Phe Gln Gly Pro Glu Gln Gly Glu Leu Leu  
 165 170 175  
 Ser Pro Cys Arg Cys Asp Gly Ser Val Lys Cys Thr His Gln Pro Cys  
 180 185 190  
 Leu Ile Lys Trp Ile Ser Glu Arg Gly Cys Trp Ser Cys Glu Leu Cys  
 195 200 205  
 Tyr Tyr Lys Tyr His Val Ile Ala Ile Ser Thr Lys Asn Pro Leu Gln  
 210 215 220  
 Trp Gln Ala Ile Ser Leu Thr Val Ile Glu Lys Val Gln Val Ala Ala  
 225 230 235 240  
 Ala Ile Leu Gly Ser Leu Phe Leu Ile Ala Ser Ile Ser Trp Leu Ile  
 245 250 255  
 Trp Ser Thr Phe Ser Pro Ser Ala Arg Trp Gln Arg Gln Asp Leu Leu  
 260 265 270  
 Phe Gln Ile Cys Tyr Gly Met Tyr Gly Phe Met Asp Val Val Cys Ile  
 275 280 285  
 Gly Leu Ile Ile His Glu Gly Pro Ser Val Tyr Arg Ile Phe Lys Arg  
 290 295 300  
 Trp Gln Ala Val Asn Gln Gln Trp Lys Val Leu Asn Tyr Asp Lys Thr  
 305 310 315 320  
 Lys Asp Leu Glu Asp Gln Lys Ala Gly Gly Arg Thr Asn Pro Arg Thr  
 325 330 335  
 Ser Ser Ser Thr Gln Ala Asn Ile Pro Ser Ser Glu Glu Glu Thr Ala  
 340 345 350  
 Gly Thr Pro Ala Pro Glu Gln Gly Pro Ala Gln Ala Ala Gly His Pro  
 355 360 365  
 Ser Gly Pro Leu Ser His His His Cys Ala Tyr Thr Ile Leu His Ile  
 370 375 380  
 Leu Ser His Leu Arg Pro His Glu Gln Arg Ser Pro Pro Gly Ser Ser  
 385 390 395 400  
 Arg Glu Leu Val Met Arg Val Thr Thr Val  
 405 410

&lt;210&gt; 45

&lt;211&gt; 3666

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 45

aagacaagta ttcctgaaat gtgtcacctg gctctgttgg ccaagccagc agtgccacaa 60  
 gagaagcagc tgccatggcc cagtgcacacc tggagtgttt gacgatgctt tgtgtccatg 120

```

tgccccctaa ccaggccttt ctacacagcca catagcatcc ccaggcctgc tgcactgggc 180
cagccgcagc ctccgagccc agcctgaggg gcctctcact gggatcccca gctttgtccc 240
tgagatgagg gggctgatcc taagagtgtg ggtggggatg cctctgcccc ggggggtctg 300
gtggtgttgg gctctggccc aggcctcctg gctctgtgtc cttccataga ctccagggtc 360
ctcacagtca ttcacaaacc aactgaatgt tgggtgtctt tggcctcttg tctcagggtt 420
ggaggtatta aagccatgga agagcagtct ctccctgggt gcaacagcct gtcttctgtg 480
gcctctagct ccatcctcag agtttgggat aggggctgaa tcttttgcct gttctccctc 540
ccagggtctg ccccagagt cagtacttca ggttgagggg agtgggaggt tggctctgga 600
ggggctgccg agtggcagga ggaaggggct caggagcagg ggggtggctcc actgcagcgg 660
ccagactctg ttgattacag atgattattt caagggtgtg gtgtgtgtgc gcgtgtgtgt 720
gtgtgcgtgt gtgtattttt aaataacagc tctattgaga tatagttcac ataccataga 780
attcaccat tgaaagtgt caattcaatg gtttctagta tgttcacaga gttgtataat 840
gaacaccaca gtttaatttta gaacatttct tctattctc ttacaatgtg taaaaataaa 900
taaagggttt tttaaaaaaa cattttcatc acacaaaaa tgagctcctc gatggatgtt 960
atccctcatt tgtcccaaaa ccccccagcc ctggcaacca ctaatctact ttctatctct 1020
ggatttgctt atcatggaca tttcttatga acggagtgc acaccatgtg gcattttgtg 1080
tctggtttct tccactgagc atcatgattt caagtctatg taggagcatg catctgagct 1140
tcattccttc tcatggtgca atcatcttcc actgtgtgta taagaccaca ttttgtcaat 1200
ccacacatcc atggatcaca agtgcatttt taaagggtgc ttttacatct caaacacagt 1260
tctcagtttg cgaggggggc agaggacaat acagaagtga ctgtaagcgg cataaactgt 1320
agcttcccag aggggctcca ggggctcggg ggagaggtcg aggtaggag gcaatgtctc 1380
tcctttgagc agccactgag agggagaagg agttgcagag atcaggaagg agaggattta 1440
tttcagttt tctggcagtt tttctcact ctttcggaga cctaggatgg gttaggagag 1500
ggcatggcaa aggcagaggt ggctctgcta gaggaggccg caggtccgca ggccgaacct 1560
ggggagccag ccatgtcttg cctgaggccc tgccttgagg agaagataaa gccttgagga 1620
gccttttggg ccgatctggg aaccagcaa gactagatta gcctgactga tgacagacct 1680
gggatgagga tgagggcagt taagtatttg ttaactctt ctttgtcttt tcaaacagcg 1740
atttagtaat cctgtttgag gctgcagtgt ggcaatgctt tccagaggat ggagtccttt 1800
ttgtttgttt tgaaaaata gagataaggt ttactatgt tgcctaggct ggtctctac 1860
tcggcctcaa gccatcctcc tgcctgggcc tcccaaaagt tggggattac aggcgtgagc 1920
catggtgcca gactggagtc cttttattaa aattaactgc cctgctcagc tttctgctgg 1980
gccaccccag agccaatcct tggttcttgg gcccaaggct ggaccaggg gttgcaggaa 2040
acagtctgta gcatccaagt ggggcctgtc gtaccactc cagtgtgtag gtgcagaacg 2100
ctctttggg gatctctctg ctgggccacc ttactccagg gatccctcag ttttcaaac 2160
aaagcaagag ggaaggaag aatggagaaa cagctcagtg ttgactctct tccccctggt 2220
gagtgctgtg ctggggcctc tgtgcacatc ataccacttc ccccttgaat cagccccaca 2280
aggcagggtg agagatgagg actcagggtg caaggaggtc tcacagcttg gaaatggatc 2340
aggacagctc tgattctcca aggccaaagt cttctctata tcatgaggca gcccaaaaat 2400
gtgtttctac acatattctg tgtagaaact caggccaggc tcctatatgc tctgggtgac 2460
agcgggaagg tgccttccag tgtgtatcag tgtggcccat ctgggagtca taatgtcatt 2520
gcctcttcca aaatatcagc aaagctgacc ctagtgttg tagtgctcaa atgcacctct 2580
ctgcctttgt gccataggga aatggatttg gtccctgggg tggtgatacc aacttctgtc 2640
tcaattccat gaccctcact cagcaccccc accccaggcc tgtgcagagg aggaataacc 2700
gtccttgaga accatccagc cctagggtga gaaattagat ttatggactt aatcagttct 2760
tcttccctgt cagtgtgtgt gagaatgaaa tgggggaaat tgataagcct cagtcagtgt 2820
ttcctgcgag acaaggcagc tttggtagct ggtgggactg acgccttctt cacctgtgtc 2880
ctctgcctct gagtactat aaattgcgaa ccccatgagg tggcacctgc agccacagat 2940
cgcttaggca ccttgacag gggccaccta atgaactggc tgcaggacgg caatcctgcc 3000
gtgaagccag gcacctcggg aggggcaagg tggcagaacc gagagagcag atgctccttg 3060
ctgtttctgt aaggccacca gtggctgggt ggccctggcc tggtagcaag gtggttaaccg 3120
tgtgagtggc tgaatcctgc accacttttc ctgagcctcc ttcctccttg attcctgcaa 3180
ctatattctc actacatggc tcaccaccaa ttcgttctct ctgccccaa gaggctggac 3240
aagaggctca ggtcagtcg ggtacgcagg aacctatgtc tgtaaaatga aaatgcctgc 3300
cttcaagggc tgtcttgagg ttgaaaggag atggtctaca tcaagggtta ggaagaccac 3360
aatgtggctg tgcgggccac ctccaggctg gcttcagtct cccatcccc a tagtcatacc 3420
cctgtgaagt ccttcccat actgaatcgg gctacctgtg tatcttatag ggtgttgaa 3480
atgacagatt gtgaattcca agggcccata aaagacagcg tgggctgggc gtggtggctc 3540
acgcctgtaa tcccagcact ttgggaggcc gagccaagtg gatcacttga ggtcaggagt 3600
ctagaccag cctggccagc atggtgaaaa cctatctcta ctaaaaaaa aaaaaaaaaa 3666
aaaaaa

```

<210> 46  
 <211> 143  
 <212> PRT  
 <213> Homo sapiens

<400> 46  
 Met Val Pro Asp Trp Ser Pro Phe Ile Lys Ile Asn Cys Pro Ala Gln  
           1                  5                  10                  15  
 Leu Ser Ala Gly Pro Pro Gln Ser Gln Ser Leu Val Leu Gly Pro Lys  
                   20                  25                  30  
 Ala Gly Pro Arg Gly Cys Arg Lys Gln Ser Val Ala Ser Lys Trp Gly  
           35                  40                  45  
 Leu Ser Tyr Pro Leu Gln Cys Val Gly Ala Glu Arg Ser Leu Gly Asp  
           50                  55                  60  
 Phe Ser Ala Gly Pro Pro Tyr Ser Arg Asp Pro Ser Val Phe Lys Thr  
           65                  70                  75                  80  
 Lys Gln Glu Gly Lys Glu Glu Trp Arg Asn Ser Ser Val Leu Thr Leu  
                   85                  90                  95  
 Phe Pro Leu Val Ser Ala Val Leu Gly Pro Leu Cys Thr Ser Tyr His  
           100                  105                  110  
 Phe Pro Leu Glu Ser Ala Pro Gln Gly Arg Val Arg Asp Glu Asp Ser  
           115                  120                  125  
 Gly Cys Lys Glu Val Ser Gln Leu Gly Asn Gly Ser Gly Gln Leu  
           130                  135                  140

<210> 47  
 <211> 2052  
 <212> DNA  
 <213> Homo sapiens

<400> 47  
 gactgtggctc gggagtaggc agcggcgccg cgctccgctct cgcccgcctct cgcccgcctcg 60  
 ccagccggct ctcctccgc cgcaggaccc gcgcgcccgc ctcggggggcc atgcagcgcc 120  
 tggccatgga cctgcggatg ttgtcccggg agctctccct ctacctggaa caccaagtcc 180  
 ggggtggggtt cttcggctcg ggggtgggct tatcccttat cctgggcttc agcgtcgctt 240  
 atgccttcta ctacctgagc agcattgccca agaaacccca gttagtgacc gggggtgaga 300  
 gtttcagccg cttccttcaa gaccactgtc ccgtgggttac agaaacgtac taccgcacgg 360  
 tctgggtgctg ggagggtcga ggacagaccc tgcttagacc tttcatcact tcgaagcccc 420  
 cgggtgcagta caggaatgaa cttattaaaa ctgcagatgg aggacagatt tcactggact 480  
 ggtttgataa tgataacagt acgtgttata tggatgccag caccagacct actatcttat 540  
 tgttgccctg cctcacggga acaagcaagg agtcatatat ccttcatatg atccatctta 600  
 gtgaagaatt aggatacaga tgtgtggttt ttaacaacag aggagtggcg ggggagaatc 660  
 tcttgacgcc aaggacttat tgttggtgcta acactgaaga cttggagaca gttattcacc 720  
 atgtacacag cctgtaccct tctgtccctt tcctggcagc aggggtttca atgggaggaa 780  
 tgctgcttct aaattacttg ggcaaaattg ggtccaaaac gcctttgatg gcagctgcaa 840  
 ctttttcctg tgggttgaac accttcgctt gctcagagtc attggaaaaa ccactgaact 900  
 ggctactttt taattactat ttgacaacct gccttcagtc ttcagttaat aagcaccgac 960  
 atatgtttgt aaaacaagtt gatattggatc atgtcatgaa ggctaaatcc atcagagagt 1020  
 ttgataagcg attcacttca gtcattgttg gataccaaac aattgatgat tattatactg 1080  
 atgccagtc gagtcctaga ctgaagtcag taggaattcc agtattgtgt ctaaattctg 1140



```

tggatgatgt tttctcacc agtcatgcta ttccaataga aactgctaag caaaatccta 1200
atgttgcttt ggtccttact tcttatggag gccatattgg ttttctggag ggaatctggc 1260
caagacagtc cacttacatg gatcgtgtct tcaagcaatt tgtgcaagcc atggttgagc 1320
atggacatga actctcttaa catgtagttc ttgggtgca ttttgtctga accacaattg 1380
tgaaggcagc tcagcttagt gcacaaatth taactgttgt atataaagca aataagccag 1440
cagatgggtg aagaggtcca gaatgatatg caaaaactac tttttagaga aacaaaacaa 1500
ctttgtagca acaaattaaa tatagtatta gattgttact tacgtagatt ttatttttac 1560
tatgccttac caagtacatc cttaaacaaa gtagtatgta catgaaattg cacttaacca 1620
aaactattgt gtaaaacaaa ttttaattcc tcagggtttt aatttaaact agtatttttt 1680
tagattattt gttttagggt atttaattgt actttaataa ctactaagaa atattggcta 1740
tttcaatgta agttataagg tggtagatcc ctaagggtat ttatagttga tgataacatg 1800
aaaactgaaa taagataaaa tacaacgtgc taaatctttt atgtattcta actttaaaag 1860
acaagtgcga caaagttaga ctgacttcta tatgtgctct tttactctga taatattaaa 1920
ttaggactaa cttatgtttt ataattgatta taatttacat gcttattttt aaaatagtat 1980
atgtggacac atatatatca ttatattaaa ataaattcta ccatttttaa aaaaaaaaaa 2040
aaaaaaaaaa aa 2052

```

&lt;210&gt; 48

&lt;211&gt; 409

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 48

```

Met Gln Arg Leu Ala Met Asp Leu Arg Met Leu Ser Arg Glu Leu Ser
  1             5             10             15

Leu Tyr Leu Glu His Gln Val Arg Val Gly Phe Phe Gly Ser Gly Val
          20             25             30

Gly Leu Ser Leu Ile Leu Gly Phe Ser Val Ala Tyr Ala Phe Tyr Tyr
  35             40             45

Leu Ser Ser Ile Ala Lys Lys Pro Gln Leu Val Thr Gly Gly Glu Ser
  50             55             60

Phe Ser Arg Phe Leu Gln Asp His Cys Pro Val Val Thr Glu Thr Tyr
  65             70             75             80

Tyr Pro Thr Val Trp Cys Trp Glu Gly Arg Gly Gln Thr Leu Leu Arg
          85             90             95

Pro Phe Ile Thr Ser Lys Pro Pro Val Gln Tyr Arg Asn Glu Leu Ile
          100             105             110

Lys Thr Ala Asp Gly Gly Gln Ile Ser Leu Asp Trp Phe Asp Asn Asp
          115             120             125

Asn Ser Thr Cys Tyr Met Asp Ala Ser Thr Arg Pro Thr Ile Leu Leu
          130             135             140

Leu Pro Gly Leu Thr Gly Thr Ser Lys Glu Ser Tyr Ile Leu His Met
          145             150             155             160

Ile His Leu Ser Glu Glu Leu Gly Tyr Arg Cys Val Val Phe Asn Asn
          165             170             175

Arg Gly Val Ala Gly Glu Asn Leu Leu Thr Pro Arg Thr Tyr Cys Cys
          180             185             190

Ala Asn Thr Glu Asp Leu Glu Thr Val Ile His His Val His Ser Leu

```

195                      200                      205  
 Tyr Pro Ser Ala Pro Phe Leu Ala Ala Gly Val Ser Met Gly Gly Met  
 210                      215                      220  
 Leu Leu Leu Asn Tyr Leu Gly Lys Ile Gly Ser Lys Thr Pro Leu Met  
 225                      230                      235                      240  
 Ala Ala Ala Thr Phe Ser Val Gly Trp Asn Thr Phe Ala Cys Ser Glu  
 245                      250                      255  
 Ser Leu Glu Lys Pro Leu Asn Trp Leu Leu Phe Asn Tyr Tyr Leu Thr  
 260                      265                      270  
 Thr Cys Leu Gln Ser Ser Val Asn Lys His Arg His Met Phe Val Lys  
 275                      280                      285  
 Gln Val Asp Met Asp His Val Met Lys Ala Lys Ser Ile Arg Glu Phe  
 290                      295                      300  
 Asp Lys Arg Phe Thr Ser Val Met Phe Gly Tyr Gln Thr Ile Asp Asp  
 305                      310                      315                      320  
 Tyr Tyr Thr Asp Ala Ser Pro Ser Pro Arg Leu Lys Ser Val Gly Ile  
 325                      330                      335  
 Pro Val Leu Cys Leu Asn Ser Val Asp Asp Val Phe Ser Pro Ser His  
 340                      345                      350  
 Ala Ile Pro Ile Glu Thr Ala Lys Gln Asn Pro Asn Val Ala Leu Val  
 355                      360                      365  
 Leu Thr Ser Tyr Gly Gly His Ile Gly Phe Leu Glu Gly Ile Trp Pro  
 370                      375                      380  
 Arg Gln Ser Thr Tyr Met Asp Arg Val Phe Lys Gln Phe Val Gln Ala  
 385                      390                      395                      400  
 Met Val Glu His Gly His Glu Leu Ser  
 405

&lt;210&gt; 49

&lt;211&gt; 2505

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

ggacaggagg ctcaaggggg cggaggcggc gttgccgggc tctccggaag gagacgtggc 60  
 ggcggttggg ccggtgatac ccgggcgctt tatagtcccg ccgcctcctc ctccacctcc 120  
 tctcctcct cctcctcctc tggagcagag gaggttgttg cgggtggctgg agaaagcggc 180  
 ggcggaggat ggaggaagga ggcggcggcg tacggagtct ggtcccgggc gggccggtgt 240  
 tactggtcct ctgcggcctc ctggaggcgt ccggcggcgg cggagccctt cctcaactca 300  
 gcgatgacat ccctttccga gtcaactggc ccggcaccga gttctctctg ccacaaactg 360  
 gagttttata taaagaagat aattatgtca tcatgacaac tgcacataaa gaaaaatata 420  
 aatgcatact tccccttgtg acaagtgggg atgaggaaga agaaaaggat tataaaggcc 480  
 ctaatccaag agagcttttg gagccactat ttaaacaag cagttgttcc tacagaattg 540  
 agtcttattg gacttacgaa gtatgtcatg gaaaacacat tcggcagtac catgaagaga 600  
 aagaaactgg tcagaaaata aatattcacg agtactacct tgggaatatg ttggccaaga 660  
 accttctatt tgaaaaagaa cgagaagcag aagaaaagga aaaatcaaat gagattccca 720

```

ctaaaaatat cgaagggtcag atgacacccat actatcctgt gggaatggga aatggtacac 780
cttgtagttt gaaacagaac cggcccagat caagtactgt gatgtacata tgtcatcctg 840
aatctaagca tgaattctt tcagtagctg aagttacaac ttgtgaatat gaagttgtca 900
ttttgacacc actcttgtgc agtcattcta aatataggtt cagagcatct cctgtgaaatg 960
acatatattt tcaatcactg ccaggatctc catttaagcc cctcaccctg aggcagctgg 1020
agcagcagga agaaatacta aggggtgcctt ttaggagaaa taaagaggaa gatttgcaat 1080
caactaaaga agagagattt ccagcgatcc acaagtcgat tgctattggc tctcagccag 1140
tgctcactgt tgggacaacc cacatatcca aattgacaga tgaccaactc ataaaagagt 1200
ttcttagtgg ttcttactgc ttctgtgggg gtgtcggttg gtggaaatat gaattctgct 1260
atggcaaaaca tgtacatcaa taccatgagg acaaggatag tgggaaaacc tctgtggttg 1320
tcgggacatg gaaccaagaa gagcatattg aatgggctaa gaagaatact gctagagctt 1380
atcatcttca agacgatggt acccagacag tcaggatggt gtcacatttt tatggaaatg 1440
gagatatttg tgatataact gacaaaccaa gacaggtgac tgtaaaacta aagtgcгааag 1500
aatcagattc acctcatgct gttactgtat atatgctaga gcctcactcc tgtcaatata 1560
ttcttggggg tgaatctcca gtgatctgta aaatcttaga tacagcagat gaaaatggac 1620
ttctttctct cccaactaa aggatattaa agttagggga aagaaaagat cattgaaagt 1680
catgataatt tctgtccac tgtgtctcat tatagagttc tcagccattg gacctcttct 1740
aaaggatggt ataaaatgac tctcaaccac tttgtgaata catatgtgta tataagaggt 1800
tattgataaa cttctgaggg agacatttgt ctgccttttt ttcatTTTTt ttgtgtctta 1860
taaaactgact gtttttcttt gcttggatac tgtgattcca aaataaatct catccaagca 1920
agtttagatc cagcctaatac aaatgtcata attgttgtag ctattgaaag tttttaaata 1980
atagatttat tatgtaaatt atagtatatg taagtagcta atgaagtaaa gatcatgaag 2040
aaagaaattg atagggtgaa atgagagacc atgtaaaata tgtaaattct agtacctgaa 2100
atcctttcaa cagattttta tatagcaact gctctctgca agtagttaa ctagaaactg 2160
ggcacatggt agaggctcac atgggagttg tctcaccct tgtaaatctc aagaaactct 2220
tatttataat aggttgcttc tctctcagaa cttttatcta ttactttttt cttcttatga 2280
gtatgtttac tctcagagta tctatctgat gtagacagtt ggtgatgctt ctgagactca 2340
gaatggttta ctctaacaaa acactgtgct gtctatccct tgtacttgcc tactgtaata 2400
tggatttcac ttctgaacag ttacagcac aatatatttatt ttaaagtgaa taaaatgtcc 2460
acaagcagtg ttgtcatgta aaaaaaaaaa aaaaaaaaaa aaaaaa 2505

```

<210> 50

<211> 483

<212> PRT

<213> Homo sapiens

<400> 50

```

Met Glu Glu Gly Gly Gly Val Arg Ser Leu Val Pro Gly Gly Pro
  1              5              10              15

Val Leu Leu Val Leu Cys Gly Leu Leu Glu Ala Ser Gly Gly Gly Arg
      20              25              30

Ala Leu Pro Gln Leu Ser Asp Asp Ile Pro Phe Arg Val Asn Trp Pro
      35              40              45

Gly Thr Glu Phe Ser Leu Pro Thr Thr Gly Val Leu Tyr Lys Glu Asp
      50              55              60

Asn Tyr Val Ile Met Thr Thr Ala His Lys Glu Lys Tyr Lys Cys Ile
      65              70              75              80

Leu Pro Leu Val Thr Ser Gly Asp Glu Glu Glu Lys Asp Tyr Lys
      85              90              95

Gly Pro Asn Pro Arg Glu Leu Leu Glu Pro Leu Phe Lys Gln Ser Ser
      100              105              110

Cys Ser Tyr Arg Ile Glu Ser Tyr Trp Thr Tyr Glu Val Cys His Gly
      115              120              125

```

Lys His Ile Arg Gln Tyr His Glu Glu Lys Glu Thr Gly Gln Lys Ile  
 130 135 140  
 Asn Ile His Glu Tyr Tyr Leu Gly Asn Met Leu Ala Lys Asn Leu Leu  
 145 150 155 160  
 Phe Glu Lys Glu Arg Glu Ala Glu Glu Lys Glu Lys Ser Asn Glu Ile  
 165 170 175  
 Pro Thr Lys Asn Ile Glu Gly Gln Met Thr Pro Tyr Tyr Pro Val Gly  
 180 185 190  
 Met Gly Asn Gly Thr Pro Cys Ser Leu Lys Gln Asn Arg Pro Arg Ser  
 195 200 205  
 Ser Thr Val Met Tyr Ile Cys His Pro Glu Ser Lys His Glu Ile Leu  
 210 215 220  
 Ser Val Ala Glu Val Thr Thr Cys Glu Tyr Glu Val Val Ile Leu Thr  
 225 230 235 240  
 Pro Leu Leu Cys Ser His Pro Lys Tyr Arg Phe Arg Ala Ser Pro Val  
 245 250 255  
 Asn Asp Ile Phe Cys Gln Ser Leu Pro Gly Ser Pro Phe Lys Pro Leu  
 260 265 270  
 Thr Leu Arg Gln Leu Glu Gln Gln Glu Glu Ile Leu Arg Val Pro Phe  
 275 280 285  
 Arg Arg Asn Lys Glu Glu Asp Leu Gln Ser Thr Lys Glu Glu Arg Phe  
 290 295 300  
 Pro Ala Ile His Lys Ser Ile Ala Ile Gly Ser Gln Pro Val Leu Thr  
 305 310 315 320  
 Val Gly Thr Thr His Ile Ser Lys Leu Thr Asp Asp Gln Leu Ile Lys  
 325 330 335  
 Glu Phe Leu Ser Gly Ser Tyr Cys Phe Arg Gly Gly Val Gly Trp Trp  
 340 345 350  
 Lys Tyr Glu Phe Cys Tyr Gly Lys His Val His Gln Tyr His Glu Asp  
 355 360 365  
 Lys Asp Ser Gly Lys Thr Ser Val Val Val Gly Thr Trp Asn Gln Glu  
 370 375 380  
 Glu His Ile Glu Trp Ala Lys Lys Asn Thr Ala Arg Ala Tyr His Leu  
 385 390 395 400  
 Gln Asp Asp Gly Thr Gln Thr Val Arg Met Val Ser His Phe Tyr Gly  
 405 410 415  
 Asn Gly Asp Ile Cys Asp Ile Thr Asp Lys Pro Arg Gln Val Thr Val  
 420 425 430  
 Lys Leu Lys Cys Lys Glu Ser Asp Ser Pro His Ala Val Thr Val Tyr  
 435 440 445

Met Leu Glu Pro His Ser Cys Gln Tyr Ile Leu Gly Val Glu Ser Pro  
 450 455 460

Val Ile Cys Lys Ile Leu Asp Thr Ala Asp Glu Asn Gly Leu Leu Ser  
 465 470 475 480

Leu Pro Asn

<210> 51

<211> 2986

<212> DNA

<213> Homo sapiens

<400> 51

```

atgggagacc ctcgatgtgc agagccttcc cctgggagaa ggagctgaaa gacaaacacc 60
ccagcttggt ccaggcattg ctggagatgg atctgctgac cgtgccaaagg aaccaaagt 120
aatctgtatc agaaatcggg gggaagatat ttgagaaggc tgtaaagaga ctctctagca 180
ttgatggctc tcaccaaatt agctctatcg tcccccttct gacggattcc agctgctgtg 240
gataccataa agcatcctac taccttgtag tcttttatga gactggatta aatgttcctc 300
gggatcagct gcagggcatg ttgtatagtt tggttggagg ccaggggagt gagaggctgt 360
cttcaatgaa tcttgggtat aaacactacc aggggtattga caactacccc ctggactggg 420
aactgtcgtg tgccctactac agcaacattg ccaccaagac accccttgac cagcacacac 480
tgcaaggaga tcaggcatat gttgaaacaa ttagactaaa agatgatgaa atactcaagg 540
tacaacccaa agaagatgga gatgtcttta tgtggttgaa gcatgaagct acccgaggca 600
atgcagcagc tcagcaacga ttggcccaga tgctgttctg ggggcagcaa ggtgtggcca 660
agaatccccg agcagcaatt gagtggtagc ccaaggggcg cctggagacg gaggatcctg 720
cgtaaatcta tgactatgcc attgtgctat tcaagggtca aggagtaaaa aagaacagac 780
ggcttgccct agagctgatg aagaaagcag cttccaaggg attgcatcag gcagtcaatg 840
gcctgggatg gtattaccac aaattcaaga aaaattacgc caaagcagca aagtactggt 900
taaaagcaga agaaatgggg aacccagatg cgtcatacaa tcttggagtc ctgcatttgg 960
atggcatctt ccctggagtt cctggaagga atcaaaacttt agctgggtgaa tatttccata 1020
aggctgcgca aggtggacac atggaagggg ccttgtgggt ttctctctac tatatcacag 1080
gcaacctgga gacattccct agagatccct agaaagctgt tgtatgggca aaacatgtag 1140
ctgagaaaaa tggctacttg ggccatgtca tccgcaaagg cctcaatgcc tacctggaag 1200
gttcatggca tgaagctttg ctgtattatg ttttagcagc agaaactgga attgaagtgt 1260
cacagacaaa ttagcacac atctgtgagg agaggccaga cctggccagg agatacttgg 1320
gtgttaactg tgtttggaga tactataatt tctctgtttt tcaaactgat gctccttcc 1380
ttgcatattt gaagatggga gacctttact actatggcca ccaaaaccag tcacaagacc 1440
tggagttgtc tgtgcagatg tacgcccagg ccgcccctgga tggagactcc cagggtttt 1500
ttaacctggc cctgctaate gaggaaggta cgataatccc acaccatate ttggatttct 1560
tggaaattga ctcaactctc cattctaata acatctccat tctccaggaa ctgtacgaaa 1620
ggtgctggag ccacagtaac gaggagtcc 1680
tgcaacttgc gcttctctgg ggtgctatcc tgcactcagc cctgatctac tttctgggaa 1740
cctttctgct atccatattg atcgccctgga ctgtgcagta ttccagctct gtctcagcaa 1800
gcgatccccc tccaagacca tcccaggcct ccccagacac tgccacgtcc actgcaagtc 1860
cagctgtgac tccagctgca gatgcctctg accaagacca gccacagta actaataacc 1920
cggagccacg tgggtgaact gtgactcca gttctctcca gatgagagag aatcttttca 1980
acagctggta ttgggaagct ggggcccagg catgatcctg ataaacacct taaatgtctt 2040
gtcaactgga tgcaaathtt gcaattgggt tcattttttt taaagtcaaa ttacaaggaa 2100
gtaccagat caggcagtg taataccaaa ggtcatcaaa cacatacaag gaacatcttg 2160
atcatagggc atgtgggaa gtttactggg ccatcacaga cttttgttct agtgattgta 2220
tgtattagga gtcatagcac gccctacggc agatctggat tcttatacac taagatgtgt 2280
cttaagaatc acagtgcgtg cttcatccct ttattgaaga acagaaaatt atgactactc 2340
tacaaggtag ataattttt ggtacctgtg cttgccacag cctgttctct caaagctgaa 2400
ttgatagatt tctctttgac ttccaagacc tagcagttat aaggcacctt gaaataaatt 2460
gtttgtgcct ggaaatgcag ggagggcaat agctttgtaa attggtttac atttttctcc 2520
ttgaattttt ctagggtcct agtgcctccg aatcatttaa tggcattgtc ggatatcttt 2580

```

```

tacatttcaa ttgcaatcca tgaaattaca tttagaagat tcttagtact taactgtagt 2640
cttctccatg aattacacgt tagaatagac tggcagcaac ggaatatgca gcaagtaagc 2700
ctctagctta tagtttcac cctacccctc atgcctgcgt gagtctgtac agggatatgt 2760
gtgtgtgtgt gtgtgtgtgt gtgttagaga ggaagaggaa gagcagaatg tctgtataca 2820
acatggctgc taaggtagtg aataaatcag taatgcaata ttgtgggtcc aaactactct 2880
ttgcactact ttatttacag tagtaaataa aattattttt atacaattga aaaaaaaaaa 2940
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 2986

```

&lt;210&gt; 52

&lt;211&gt; 640

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

```

Met Cys Arg Ala Phe Pro Trp Glu Lys Glu Leu Lys Asp Lys His Pro
  1              5              10              15

Ser Leu Phe Gln Ala Leu Leu Glu Met Asp Leu Leu Thr Val Pro Arg
          20              25              30

Asn Gln Asn Glu Ser Val Ser Glu Ile Gly Gly Lys Ile Phe Glu Lys
      35              40              45

Ala Val Lys Arg Leu Ser Ser Ile Asp Gly Leu His Gln Ile Ser Ser
      50              55              60

Ile Val Pro Phe Leu Thr Asp Ser Ser Cys Cys Gly Tyr His Lys Ala
      65              70              75              80

Ser Tyr Tyr Leu Ala Val Phe Tyr Glu Thr Gly Leu Asn Val Pro Arg
          85              90              95

Asp Gln Leu Gln Gly Met Leu Tyr Ser Leu Val Gly Gly Gln Gly Ser
          100              105              110

Glu Arg Leu Ser Ser Met Asn Leu Gly Tyr Lys His Tyr Gln Gly Ile
          115              120              125

Asp Asn Tyr Pro Leu Asp Trp Glu Leu Ser Tyr Ala Tyr Tyr Ser Asn
          130              135              140

Ile Ala Thr Lys Thr Pro Leu Asp Gln His Thr Leu Gln Gly Asp Gln
          145              150              155              160

Ala Tyr Val Glu Thr Ile Arg Leu Lys Asp Asp Glu Ile Leu Lys Val
          165              170              175

Gln Thr Lys Glu Asp Gly Asp Val Phe Met Trp Leu Lys His Glu Ala
          180              185              190

Thr Arg Gly Asn Ala Ala Ala Gln Gln Arg Leu Ala Gln Met Leu Phe
          195              200              205

Trp Gly Gln Gln Gly Val Ala Lys Asn Pro Glu Ala Ala Ile Glu Trp
          210              215              220

Tyr Ala Lys Gly Ala Leu Glu Thr Glu Asp Pro Ala Leu Ile Tyr Asp
          225              230              235              240

Tyr Ala Ile Val Leu Phe Lys Gly Gln Gly Val Lys Lys Asn Arg Arg

```

245	250	255
Leu Ala Leu Glu Leu Met Lys Lys Ala Ala Ser Lys Gly Leu His Gln 260 265 270		
Ala Val Asn Gly Leu Gly Trp Tyr Tyr His Lys Phe Lys Lys Asn Tyr 275 280 285		
Ala Lys Ala Ala Lys Tyr Trp Leu Lys Ala Glu Glu Met Gly Asn Pro 290 295 300		
Asp Ala Ser Tyr Asn Leu Gly Val Leu His Leu Asp Gly Ile Phe Pro 305 310 315 320		
Gly Val Pro Gly Arg Asn Gln Thr Leu Ala Gly Glu Tyr Phe His Lys 325 330 335		
Ala Ala Gln Gly Gly His Met Glu Gly Thr Leu Trp Cys Ser Leu Tyr 340 345 350		
Tyr Ile Thr Gly Asn Leu Glu Thr Phe Pro Arg Asp Pro Glu Lys Ala 355 360 365		
Val Val Trp Ala Lys His Val Ala Glu Lys Asn Gly Tyr Leu Gly His 370 375 380		
Val Ile Arg Lys Gly Leu Asn Ala Tyr Leu Glu Gly Ser Trp His Glu 385 390 395 400		
Ala Leu Leu Tyr Tyr Val Leu Ala Ala Glu Thr Gly Ile Glu Val Ser 405 410 415		
Gln Thr Asn Leu Ala His Ile Cys Glu Glu Arg Pro Asp Leu Ala Arg 420 425 430		
Arg Tyr Leu Gly Val Asn Cys Val Trp Arg Tyr Tyr Asn Phe Ser Val 435 440 445		
Phe Gln Ile Asp Ala Pro Ser Phe Ala Tyr Leu Lys Met Gly Asp Leu 450 455 460		
Tyr Tyr Tyr Gly His Gln Asn Gln Ser Gln Asp Leu Glu Leu Ser Val 465 470 475 480		
Gln Met Tyr Ala Gln Ala Ala Leu Asp Gly Asp Ser Gln Gly Phe Phe 485 490 495		
Asn Leu Ala Leu Leu Ile Glu Glu Gly Thr Ile Ile Pro His His Ile 500 505 510		
Leu Asp Phe Leu Glu Ile Asp Ser Thr Leu His Ser Asn Asn Ile Ser 515 520 525		
Ile Leu Gln Glu Leu Tyr Glu Arg Cys Trp Ser His Ser Asn Glu Glu 530 535 540		
Ser Phe Ser Pro Cys Ser Leu Ala Trp Leu Tyr Leu His Leu Arg Leu 545 550 555 560		
Leu Trp Gly Ala Ile Leu His Ser Ala Leu Ile Tyr Phe Leu Gly Thr		

	565		570		575
Phe Leu Leu Ser Ile Leu Ile Ala Trp Thr Val Gln Tyr Phe Gln Ser					
580		585		590	
Val Ser Ala Ser Asp Pro Pro Pro Arg Pro Ser Gln Ala Ser Pro Asp					
595	600		605		
Thr Ala Thr Ser Thr Ala Ser Pro Ala Val Thr Pro Ala Ala Asp Ala					
610	615		620		
Ser Asp Gln Asp Gln Pro Thr Val Thr Asn Asn Pro Glu Pro Arg Gly					
625	630	635	640		

<210> 53  
 <211> 1908  
 <212> DNA  
 <213> Homo sapiens

<400> 53

```

gaactcaaga tggcggacga gaggtagaca gcagtgaagc cgccggcacc tccgctgccg 60
cagatgatgg aaggggaacgg gaacggccat gagcactgca gcgattgcga gaatgaggag 120
gacaacagct acaaccgggg gtaacgaaat cctcggagtc caattcccgt ccagcctccc 180
acaacctggg tctctggggg gcggggaagt caccgggagc ttagtctagc cccaccctca 240
tggttcttatt ggctaagtca ctaggattct gcctggcgat tggttattgc ctttgctcatt 300
taccaagggt ggggtgctctg gatattaagg gccggagacg ttcagtactt ttattatata 360
aaagttctat gcagtgttcc actcgtgtgc ttcaggccat tcctgggaaa agagcatgtt 420
ggacgatggg ttctgatggg acaacacagg acagtctagt aacagtacag ttcaaaggga 480
acgttactta tagacgtgtt gtttgaacaa cgtgactagc aataaacgaa tcataaaaca 540
tttcgcagca gtccaccctc agtgaggaaa ctcatcagac tatacaaaact gtatgtataa 600
ttatattgta aaatatcact accagagaat cccaaaatat aaactcacct tataaaccta 660
tgtaattaac ggtatggtaa tgctttgatt tacatagttt ccattgctcc aaataatcca 720
taagtatcct gaacgatata ttaatatata taagtcttcg aacatctttg ccttggtacc 780
tttgacagag agccccaggc ttcaagaaat gagcgatgtt gtttcttggg gagtatttgg 840
aatcctcggg ctccatactg cctatcctgt actttagggt cttgccatta gttactattt 900
gttcctaggg atgaagaaac acaccaggca tttgaatgtt gaattctttt cccaaattag 960
tttttacttc tgtaagccac agcttattca tcttcttcca catggtagt gcctggcact 1020
ggcactcacg ctgtctctcc ccattcttct gtctatattt gtagatata ttttaaaactg 1080
aaagtctgta gagatttttg ggtgtataac ttgctgagtc tgaggtagtc tgattttcgt 1140
agccagcttg ttattcgtag tattggtggc agtgcagata taaaggata aagaccaagg 1200
actgcgtaga agagatgggt ccacaagatt atagtacata tttttattgt atcctttcta 1260
tgtttaggta tgcttagata cacaataact taccattgta ttatagttgc ctgcagtact 1320
cagtaacatg cagcagccta ttgctcctag gctacaaacc tgtatagcat gttactgtaa 1380
tgtacaccat atagcctagg catttagtag gttataccat ctgattttgt gtaagtacac 1440
cctataatgt tggcaaggca aaaatcacct aaccatgcat ttcccagcac atatgcctga 1500
cgttcagtga agcatgactg tatttaagtg gtttataatc tgcttggtgt aaatcttagg 1560
ttaaagatc tagcttttat gttcttcagc cggcattaaa aatcacagga agggccgggc 1620
atggtggctc atgcctgtaa tcccagtagt ttgggaggca gatcacgagg tcaggagatc 1680
gagaccatcc tggctaacac tgtgaaatcc catctctact aaaaaataca aaaatttagc 1740
cgggcggtgg gggggcgcc tgtagtcca actactcggg aggctaaggc aggagaatgg 1800
cgtgaaccca ggaggcagag cttgcagtga gctgagatca cgccactgca ctccagcctg 1860
ggcgacagag cgagactccg actcaaaaaa aaaaaaaaaa aaaaaaaa 1908

```

<210> 54  
 <211> 88  
 <212> PRT  
 <213> Homo sapiens

<400> 54



Met Phe Leu Leu Ala Lys Ser Leu Gly Phe Cys Leu Ala Ile Gly Tyr  
 1 5 10 15

Cys Leu Cys His Leu Pro Arg Val Gly Ala Leu Asp Ile Lys Gly Arg  
 20 25 30

Arg Arg Ser Val Leu Leu Leu Tyr Lys Ser Ser Met Gln Cys Ser Thr  
 35 40 45

Arg Val Leu Gln Ala Ile Pro Gly Lys Arg Ala Cys Trp Thr Met Val  
 50 55 60

Phe Asp Gly Thr Thr Gln Asp Ser Leu Val Thr Val Gln Phe Lys Gly  
 65 70 75 80

Asn Val Thr Tyr Arg Arg Val Val  
 85

&lt;210&gt; 55

&lt;211&gt; 2969

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 55

gagttgagag atggcgggccg ccgcaggttag atcgctcctg ctgctcctct cctctcgggg 60  
 cggcgggcggc gggggcgccg gcggctgcgg ggcgctgact gccggctgct tccctgggct 120  
 gggcgctcagc cgccaccggc agcagcagca ccaccggacg gtacaccaga ggatcgcttc 180  
 ctggcagaat ttgggagctg tttattgcag cactgttgtg ccctctgatg atgttacagt 240  
 ggtttatcaa aatgggttac ctgtgatatc tgtgaggcta ccattcccggc gtgaacgctg 300  
 tcagttcaca ctcaagccta tctctgactc tgttggtgta tttttacgac aactgcaaga 360  
 agaggatcgg ggaattgaca gatttgctat ctattcacca gatgggtgtc gcgttgctgc 420  
 ttcaacagga atagacctcc tcctccttga tgactttaag ctggtcatta atgacttaac 480  
 ataccacgta cgaccaccaa aaagagacct cttaagtcag gaaaatgcag caacgctgaa 540  
 tgatgtaaag acattggtcc agcaactata caccacactg tgcattgagc agcaccagtt 600  
 aaacaaggaa agggagctta ttgaaagact agaggatctc aaagagcagc tggctccctc 660  
 ggaaaaggta cgaattgaga ttagcagaaa agctgagaag aggaccactt tgggtgctatg 720  
 ggggtggcctt gcctacatgg ccacacagtt tggcattttg gcccggtta cctgggtggga 780  
 atattcctgg gacatcatgg agccagtaac atacttcac acttatggaa gtgccatggc 840  
 aatgtatgca tattttgtaa tgacacgcca ggaatatgtt tatccagaag ccagagacag 900  
 acaatactta ctatttttcc ataaaggagc caaaaagtca cgttttgacc tagagaaata 960  
 caatcaactc aaggatgcaa ttgctcaggc agaaatggac cttaagagac tgagagaccc 1020  
 attacaagta catctgcctc tccgacaaat tggtgaaaaa gattgatctg caaaaagcct 1080  
 ctgaatcctg gcagaaggaa cacctgtttg cctttttaat taaagcattg cagggtggaag 1140  
 ctgggagcca tgtggggggt agagcgtttt tacctttaat tataaaacaa aaacagaaag 1200  
 gatctgaggg aagaaggga tgtaaaacc tgaggatcag gcattgtgga atataagctc 1260  
 aaagggtcta gtgaatattg tcttaaccaa gtatctcagt ttctggatga aaatgatgca 1320  
 gttatatagt tgagagattc ataaagagaa aacaatgctg ggggtgttcg tttcttgcat 1380  
 cttctttgca ggtcagcaa aagagtaaca caccagcacc ccactcgact ctatttggtt 1440  
 ttaatttaac tgtccctatt ttgacatag gagtaataa atatactaga aaagcaaat 1500  
 ctcattgatg gctaaaatat cattagcatt tattttaaat tggaccagc ctctgcagag 1560  
 ttaccaggaa tctttccttc cagcatccct ttactgacca cctacctgta cctcttggtt 1620  
 acactttttt ttccatttga taattggaac caactataa ctgtttaata attgacatt 1680  
 tagattatct cttataacct tcttaaatgt ctatatatcc cagtgtctct gatcagtgct 1740  
 taaaaatcac tggcaacct gcattgaggt gttgggtttt ttttgtttta ttaattagtc 1800  
 tttcacagga ggaataattg ccctccttta tatacttatc tattgataat cccctctccc 1860  
 tccagaacac aaatcagagg gaaagggggt gtccagctgt actaccaa atcaggaagatg 1920  
 taagggttac aaattggcta agaactcatg ctctgtagcc atttcaacca gaataatttt 1980  
 attgctaate tgctttgtgt gacagcattc caggccagcc agatgggact gccttgctctg 2040  
 gaggccttgt tcatctcgaa ggacacacac ttccacactg tttgtgagcc ctcccacctc 2100

```

cacaacttca gttgtaaatc aagtgtgtgg atctcaaagg gtgcaattta tctttatata 2160
ggaatacatt tctagggctt ccttcaagcc cactctcttc accctatttt ttcttatctt 2220
aaattgagag aaagagaatt aatcttatac ttgtgcaaaa ctttttctac catatttcca 2280
gatgacatct gcgcttgaag agtcaaagga atctgtgtct aatatacctgt ttttaactgc 2340
tgtaggggca ggatggaaag gatgatgggg gctgccacac cactgattgg ctttttcttt 2400
cacgtgattc atccttcttc attgtggcaa ggagttcttt ctctttttct tcctcctttg 2460
ggatcattgt gtatgaaaag aaaaacttta aatgacaaac ccagactcca ggtgccttgc 2520
aaagggttgaa ggccagccag gattgctgct gctgctgcta ctctgccaa cacccttttc 2580
attggcatga cggaatgaaa ggatgcatgt ctccacttcc tgaccctccg cccacttcc 2640
tctccctcca ccaccccccag tcgtcagctc cttccctcat ttatttttgt taagttgtgt 2700
gaattatttt taacccattt atcctgtttg tgcatagggt ttttaagaag aaacagcaca 2760
gtgcaacgag caaatctttt tgggggtgtg gggaagcaag ggagggagga catggagaaa 2820
agttctttta acaaatagca aactattgaa catgtgtaaa atcctgtatc atttatgaaa 2880
tatgtataaa aagcaatgta cttcttgtaa caataaatac ttattcaatt ttgaaaaaaa 2940
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2969

```

&lt;210&gt; 56

&lt;211&gt; 351

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 56

```

Met Ala Ala Ala Ala Gly Arg Ser Leu Leu Leu Leu Leu Ser Ser Arg
  1                      5                      10                      15

```

```

Gly Gly Gly Gly Gly Gly Ala Gly Gly Cys Gly Ala Leu Thr Ala Gly
      20                      25                      30

```

```

Cys Phe Pro Gly Leu Gly Val Ser Arg His Arg Gln Gln Gln His His
      35                      40                      45

```

```

Arg Thr Val His Gln Arg Ile Ala Ser Trp Gln Asn Leu Gly Ala Val
      50                      55                      60

```

```

Tyr Cys Ser Thr Val Val Pro Ser Asp Asp Val Thr Val Val Tyr Gln
      65                      70                      75                      80

```

```

Asn Gly Leu Pro Val Ile Ser Val Arg Leu Pro Ser Arg Arg Glu Arg
      85                      90                      95

```

```

Cys Gln Phe Thr Leu Lys Pro Ile Ser Asp Ser Val Gly Val Phe Leu
      100                      105                      110

```

```

Arg Gln Leu Gln Glu Glu Asp Arg Gly Ile Asp Arg Val Ala Ile Tyr
      115                      120                      125

```

```

Ser Pro Asp Gly Val Arg Val Ala Ala Ser Thr Gly Ile Asp Leu Leu
      130                      135                      140

```

```

Leu Leu Asp Asp Phe Lys Leu Val Ile Asn Asp Leu Thr Tyr His Val
      145                      150                      155                      160

```

```

Arg Pro Pro Lys Arg Asp Leu Leu Ser His Glu Asn Ala Ala Thr Leu
      165                      170                      175

```

```

Asn Asp Val Lys Thr Leu Val Gln Gln Leu Tyr Thr Thr Leu Cys Ile
      180                      185                      190

```

```

Glu Gln His Gln Leu Asn Lys Glu Arg Glu Leu Ile Glu Arg Leu Glu
      195                      200                      205

```

Asp Leu Lys Glu Gln Leu Ala Pro Leu Glu Lys Val Arg Ile Glu Ile  
 210 215 220  
 Ser Arg Lys Ala Glu Lys Arg Thr Thr Leu Val Leu Trp Gly Gly Leu  
 225 230 235 240  
 Ala Tyr Met Ala Thr Gln Phe Gly Ile Leu Ala Arg Leu Thr Trp Trp  
 245 250 255  
 Glu Tyr Ser Trp Asp Ile Met Glu Pro Val Thr Tyr Phe Ile Thr Tyr  
 260 265 270  
 Gly Ser Ala Met Ala Met Tyr Ala Tyr Phe Val Met Thr Arg Gln Glu  
 275 280 285  
 Tyr Val Tyr Pro Glu Ala Arg Asp Arg Gln Tyr Leu Leu Phe Phe His  
 290 295 300  
 Lys Gly Ala Lys Lys Ser Arg Phe Asp Leu Glu Lys Tyr Asn Gln Leu  
 305 310 315 320  
 Lys Asp Ala Ile Ala Gln Ala Glu Met Asp Leu Lys Arg Leu Arg Asp  
 325 330 335  
 Pro Leu Gln Val His Leu Pro Leu Arg Gln Ile Gly Glu Lys Asp  
 340 345 350

&lt;210&gt; 57

&lt;211&gt; 1727

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 57

gggaaggga tccttgaggc acccgaatgg tccctcaggg tgcagggagg cagaagcctg 60  
 gccacagagg agcctcctaa ggcagcagct gcagcaagcg caccctctcc ccactctccc 120  
 cagccagagg cggcttccag agcagatgct gtttccatcc tccctcgtaa aaccattctc 180  
 gctgctgagc ttgacaatct gggcaaggct tgtggggcgc ttgacaaca gaatctgccc 240  
 tgtgccgcct ggttccgtgg cctccagcat ggcctgcag gcagggcgct gcgggaaccc 300  
 agttgtgctg cccagcccca tgccctcggg tctgctgtgc atgaatgagt gctcacttgt 360  
 ccggggttta ggacgtggtc aagtgaacag cagggtctaa ctgtgcttac ttagccagct 420  
 tcaaacagaa caaaggaaaa atatagaaag caacatctgt tgatcattta gggttttttt 480  
 taaaccacca tgctactttg agtccttcat ggggttttga acagcattta tcaagaagaa 540  
 aatgtgggct ttttcccctc tcccggtgtt tgtttgcct gtagatagag ggaggaaagc 600  
 cgtgcagtgg caggcgggac cccctctggt ggcgggaccc cctcttgagg tggctctgag 660  
 gggccagccg ggacctgtca ctttattatt taaggagtgt gtgtgtagag tcgctggcct 720  
 attaacagta ttgtgtgtgg gttgggtttt tagtttgttc cttctttttg aagtccttcc 780  
 atttcaatcc ttgactctct cttcccctcc cttgccagc tctgttgaat gctgctgtgc 840  
 gcgtgtgagg gccgctctgc acacagggcc cttgggttgt gtgaactgaa attctccctg 900  
 tattttgtgag actgcagga gtccccatct gtagcacagg caatgccagt gccatgctgc 960  
 agcctcagaa accaggcctc tcaactccagc agcaggcaga accgtgtctg tggctcgggtg 1020  
 ctgtccacag ctctgtctgc cttgttcttg ggcttgagct ggatagaggt ggggtctctt 1080  
 cacccttccc gaattcagaa cagacctgt gcctggcccc agtgtgccc ggcaattccc 1140  
 caggccctca ttgggagccc ttggtgttct gaggcaggc gcccaggcag cacatgagca 1200  
 gtgccagggg gctccctgcg tgaggacggc aagggtgcg gtatgtctaa cttattgatg 1260  
 gcaggcagcc ccctgtgccc cctaagcctg gccctggtta ttgctgagct ctgtgctcag 1320  
 tgctgcggcc tggccgtggc tcgtctgttc ctttgggggg cccgggaggg ttgtgggaat 1380  
 cagctcttcac agacagacgt gagccaggcg gaggactcgt tccttgca gaagtcagct 1440  
 caccctgcag tgctcgggtg ggggggggca aggaggggca ggcacacacc atgtctgacc 1500

tgaacccgat tctggggagc atcttcccgc tccggcccca cgacctccac agggttacat 1560  
 tgtaatatat atgccccagc taacctgtct gatgggtggca tcttcctgca gacatttcaa 1620  
 acatgtaact tttatatgaa aaaaaataaa cacagatgaa agctgaaaaa aaaaaaaaaa 1680  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 1727

<210> 58  
 <211> 115  
 <212> PRT  
 <213> Homo sapiens

<400> 58  
 Met Trp Ala Phe Ser Pro Leu Pro Cys Phe Val Cys Pro Val Asp Arg  
 1 5 10 15  
 Gly Arg Lys Ala Val Gln Trp Gln Ala Gly Pro Pro Leu Val Ala Gly  
 20 25 30  
 Pro Pro Leu Ala Val Val Leu Arg Gly Gln Pro Gly Pro Val Thr Leu  
 35 40 45  
 Leu Phe Lys Glu Cys Val Cys Arg Val Ala Gly Leu Leu Thr Val Leu  
 50 55 60  
 Cys Val Gly Trp Val Phe Ser Leu Phe Leu Leu Phe Glu Val Pro Ser  
 65 70 75 80  
 Phe Gln Ser Leu Thr Leu Ser Pro Leu Pro Leu Pro Ser Ser Val Glu  
 85 90 95  
 Cys Cys Cys Ala Arg Val Arg Ala Ala Leu His Thr Gly Pro Leu Gly  
 100 105 110  
 Cys Val Asn  
 115

<210> 59  
 <211> 2617  
 <212> DNA  
 <213> Homo sapiens

<400> 59  
 tactgactcg aggggttccc ttccaaaata tgcagggtc aggcctccaa ttccgggcct 60  
 gtctgctttg cttgtgtttc tctgtccct gttctcccg agggcccagg tggaactcac 120  
 gacagggagg gagacgttc ccaaaacct gcagggtat ttcccagaat ttggttttca 180  
 agtacaaaac tttttgtcct gtaagatata tgcagcctca cagaagcagc ctctgcctcc 240  
 actttaccag ctacgttttt atcttaagca catggggctc ccttagaact tactccactg 300  
 atttaaaaaa aaaaaaactg cctggcagca tctcagtgtc agagttagca cggcacagga 360  
 aaggcccgtg gtgacgaggg tgagggtggc acagtggacc gacgacaaat gagactctgc 420  
 aaatgagact ccagaggggtg aagatctgct gtctccagac atcataggcc atgtgaccca 480  
 ctaggggccg cttaccctcg gccgtccgct ggctgaactg aacgcattcc ctctctccgc 540  
 aactctcccg tgaggctgca cccgtgtggg tagcactgga agcggcactg tttgcattgt 600  
 acataggaag gaaggaagt ctccagcct caccagcacc tggcagcgag tcagagcctg 660  
 tgaggggcac cgaagcagt atgcagtgtc aacctcccag ctgggtgccac tctgccctcg 720  
 ggggctccaa gcattgtaac tcagtcattg gagctgcctc tttggaagtg cagatttatt 780  
 cctgtaataa tctgcctgc ttttacctct cgtccactga ccagcaagtg tgagtcccgg 840  
 tgtcagtcgg cacagtccag tgtccatctg catttgctca tgcagagggg gtgagttggg 900  
 cactccctgt tgttggtttt ccttttgtag cacactgggc agtctcccta taaaacaaaa 960  
 accccacctt ctgtgccttc tgcttttagag cagagctccc cctcccatct cctcagtcct 1020  
 ccctgcaaaa tctgtccacc ggggaaggca gcaggaaccc tgggcagcgg gtgttctggg 1080

```

aaggctagtg acagcagatg tcattccagga acagccacac acggttctcc aggccgccgt 1140
cagcagctca aggtggggta tgagtgagaa gctgaggatc tcgcagcttg ttgctgagca 1200
aggtgcaacc gggctcatgc tgtcatcagc acaagacggg atggcaaggg ctttcagacg 1260
catttccaag agtccagcaa gccaggggga agatgatccc tttgccgaag tgtaccctct 1320
agccaacttt tgggagcgct tctgtttgca aagcgctggg gatgtgcctg tctctgtgtg 1380
acccacgaac gggaaggagg agcactggag taatgacact tctgctgctg ctttgattct 1440
caaggctgat ctttaaaacc ctgccttgcc tgacaggtgc tttaaaggca gtctgcatct 1500
tttcttccct tgggtgtgga gaggtaaaca ctttgatttg ctgaaagctg tatggagtat 1560
atttgaacag ctagtagtta gctttgaaag tggaagtgtg aacagacact acttggtgtg 1620
ctttgggtcc ttacttttac cccacagaa gtctagaggc gtctgttata aagcgttacg 1680
gggcgcctgc atgcaggagg aaggacctgt attagctgga aatcatcagg aaccagctt 1740
gcctccatct ctctgagatg tgctgggtac agcctgcccc tcctagtctt gtccaccggg 1800
aagagccggc tggcggcaga tcccagggg cagagcccct gctggatcct gggagctcat 1860
ctttacctgt gccggagtgg gaactgtgat tccagccggg caggtcagag tggagcagtg 1920
ctaagaggct gttgcaggag aactagacgg gcggggcctg ctgcatctgg atcatgtttc 1980
tgtgctctgc cccgcgctag ggactcaggg tctgggcttc tgccagggtg ggagcagaga 2040
gactgttccc ttgggtggag aggtgtgggc atgagagcca cccattgcca agcagcaaga 2100
atgttctgctg ttttttccag agaggggaac cccactggtt tttgtggaaa caatggaac 2160
ttacagatgc ctgcctggga tgatgaggca cattcagaac aaatgctttt tttttttttg 2220
agacagatgc tcgctctgac gcccaggctg gagtgcagtg gcgcgatctc ggctcactgc 2280
aaactttgcc tcccagggtc aagtgtattc cctacctcag cctcccgagt agctgggatt 2340
acaccaccat gcccgcaaaa tttttgtgtt tttagtagag acggagtctc accatgttgg 2400
ccaggctggt ctggaactcc tgacctcagg tgatccatcc gccttggcct cccaaagtgc 2460
tggtgattaca ggcgggagcc accatgcctg gccagaacaa atgccttttt aaacctttta 2520
agaacatttt taaaatgtct ttttctatgt caaatgtaac gtttattttt ttaaacaata 2580
aaattgattt gccaaaaaaa aaaaaaaaaa aaaaaaa 2617

```

&lt;210&gt; 60

&lt;211&gt; 105

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 60

```

Met Gln Gly Ser Gly Ser Gln Phe Arg Ala Cys Leu Leu Cys Leu Cys
  1             5             10             15

```

```

Phe Ser Cys Pro Cys Ser Pro Gly Gly Pro Arg Trp Asn Ser Arg Gln
      20             25             30

```

```

Gly Gly Arg Arg Phe Pro Lys Thr Cys Arg Ala Ile Ser Gln Asn Leu
  35             40             45

```

```

Val Phe Lys Tyr Lys Thr Phe Cys Pro Val Arg Tyr Met Gln Pro His
  50             55             60

```

```

Arg Ser Ser Leu Cys Leu His Phe Thr Ser Tyr Val Phe Ile Leu Ser
  65             70             75             80

```

```

Thr Trp Gly Ser Leu Arg Thr Tyr Ser Thr Asp Leu Lys Lys Lys Lys
      85             90             95

```

```

Leu Pro Gly Ser Ile Ser Val Ser Glu
      100             105

```

&lt;210&gt; 61

&lt;211&gt; 1457

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 61

```

tttttttttg ctctcgctgg gaatggcgga gggagagaaa aaccaagatt tcactttcaa 60
gatggaaagt ccgtcagact cagctgtggt ttacctagc actcctcagg cctctgcgaa 120
tccatcatct ccctatacaa atagttcccg aaaacaacct atgagtgcga cacttagaga 180
aagattaagg aaaacaagat ttctatttaa ttctctttac aatgtggtga aacgtcttaa 240
agtagagagt gaagaaaatg atcagacctt ttcagagaaa ccagcatctt ccacagagga 300
aaactgtttg gaatttcaag aaagttttaa acatatagac agtgaatttg aagaaaatac 360
aaatttgaaa aatactttga agaattctaa tgtctgtgaa tctcagtcac ttgattctgg 420
atcatgcagt gctctccaaa atgagtttgt gagtgcgaag cttcctaaac aaagattaaa 480
cgctgaaaaa gccaaatttg tgaagcaggt tcaggagaaa gaagaccttc ttcggagggt 540
aaaactagtc aaaatgtata gatcaaagaa tgatctgtct cagttacagt tgtaataaaa 600
gaagtggaga agctgtagcc agctctgtct ttatgagttg cagtcagctg tgtctgaaga 660
gaacaagaaa ctaagcctta ctcaattgat agaccactat gggtttagatg ataaattact 720
acactataac agaagtgaag aagaatttat agatgtttaa ttctgtattt ttgctccaga 780
atatctttga gaatgacaac ttaattaaaa gatacttagg cacttttttt ttttttttga 840
gactgagttt cgctcttgct atcctggctg gagtgtgatg gtgcgatctt gactcactgc 900
aacctctgcc tctcgggttc cagcaattct cctgcctcag cctcccagat agctgagatt 960
acaggcgccc gccaccatgc cgggctaatt tttgcatttt tagtagagac tgggtttcac 1020
cacgttgccc aggtgtgtct cgaactcctg acctcaggtg atccaccgcc taggectccc 1080
aaaaccatta gggctcagag gaaggtatcc caatgaatat caattaaggg cactttaata 1140
tataaattat aaactaagtt ctaaaaggaa aattagtatt ttggatagat ttgtcaaac 1200
gacatttaag tcatgtttaa aaagtcattt gggcagttct ggaactagat ttaatacat 1260
ttgtttttta tgacaaaaag ttttatttta aatgttaaaa attgtccaat ctggtgaatg 1320
tctaacccta aagtttaaaa atttctgcct cctaagttta tgtacctgt ttccatccat 1380
ttaccacata tttccatctg atagtctagc aggtaattaa acttatatgt ccaaaaaaaaa 1440
aaaaaaaaaa aaaaaaa 1457

```

&lt;210&gt; 62

&lt;211&gt; 245

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 62

```

Met Ala Glu Gly Glu Lys Asn Gln Asp Phe Thr Phe Lys Met Glu Ser
  1              5              10              15

Pro Ser Asp Ser Ala Val Val Leu Pro Ser Thr Pro Gln Ala Ser Ala
 20              25              30

Asn Pro Ser Ser Pro Tyr Thr Asn Ser Ser Arg Lys Gln Pro Met Ser
 35              40              45

Ala Thr Leu Arg Glu Arg Leu Arg Lys Thr Arg Phe Ser Phe Asn Ser
 50              55              60

Ser Tyr Asn Val Val Lys Arg Leu Lys Val Glu Ser Glu Glu Asn Asp
 65              70              75              80

Gln Thr Phe Ser Glu Lys Pro Ala Ser Ser Thr Glu Glu Asn Cys Leu
 85              90              95

Glu Phe Gln Glu Ser Phe Lys His Ile Asp Ser Glu Phe Glu Glu Asn
100              105              110

Thr Asn Leu Lys Asn Thr Leu Lys Asn Leu Asn Val Cys Glu Ser Gln
115              120              125

Ser Leu Asp Ser Gly Ser Cys Ser Ala Leu Gln Asn Glu Phe Val Ser
130              135              140

```

Glu Lys Leu Pro Lys Gln Arg Leu Asn Ala Glu Lys Ala Lys Leu Val  
 145 150 155 160  
 Lys Gln Val Gln Glu Lys Glu Asp Leu Leu Arg Arg Leu Lys Leu Val  
 165 170 175  
 Lys Met Tyr Arg Ser Lys Asn Asp Leu Ser Gln Leu Gln Leu Leu Ile  
 180 185 190  
 Lys Lys Trp Arg Ser Cys Ser Gln Leu Leu Leu Tyr Glu Leu Gln Ser  
 195 200 205  
 Ala Val Ser Glu Glu Asn Lys Lys Leu Ser Leu Thr Gln Leu Ile Asp  
 210 215 220  
 His Tyr Gly Leu Asp Asp Lys Leu Leu His Tyr Asn Arg Ser Glu Glu  
 225 230 235 240  
 Glu Phe Ile Asp Val  
 245

<210> 63  
 <211> 4093  
 <212> DNA  
 <213> Homo sapiens

<400> 63  
 tttgttccgg gagccgtcac cacagtaggt ccctcgggtc agtcgggcca gcccctctca 60  
 gtccctcccca acccccacaa ccgcccgcgg ctctgagacg cggcccccggc ggcggcggca 120  
 gcagctgcag catcatctcc accctccagc catggaagac ctggaccagt ctccctctggt 180  
 ctctctctcg gacagccca cccggccgca gcccgcggtc aagtaccagt tcgtgaggga 240  
 gcccgaggac gaggagggaag aagaggagga ggaagaggag gacgaggacg aagacctgga 300  
 ggagctggag gtgctggaga ggaagcccg cgcggggctg tccgcgggcc cagtgccacc 360  
 cgccctgcc gccggcgcg cctgatgga ctctcgaaat gacttcgtgc cgccggcgcc 420  
 cgggggaccc ctgcccggcg cccccccgt cgccccggag cggcagccgt cttgggaccc 480  
 gagcccggtg tcgtcgaccg tgcccgcgcc atccccgctg tctgctgccc cagtctcgcc 540  
 ctccaagctc cctgaggacg acgagcctcc ggcccggcct cccctctctc ccccgccag 600  
 cgtgagcccc caggcagagc ccgtgtggac cccgccagcc ccggctcccc ccgcgcccc 660  
 ctccaccccg gccgcgcca agcgcagggg ctctcggggc tcagtggatg agaccctttt 720  
 tgctcttctc gctgcatctg agcctgtgat acgctctct gcagaaaata tggacttgaa 780  
 ggagcagcca ggaactacta ttctggctgg tcaagaggat ttcccatctg tctgtcttga 840  
 aactgctgct tctcttctct ctctgtctcc tctctcagcc gcttctttca aagaacatga 900  
 ataccttggt aatttgtaa cagtattacc cactgaagga acacttcaag aaaatgtcag 960  
 tgaagcttct aaagaggtct cagagaaggc aaaactctac tcatagatag agatttaaca 1020  
 gagttttcag aattagaata ctcaaaaatg ggatcatcgt tcagtgtctc tccaaaagca 1080  
 gaatctgccg taatagtagc aaatcctagg gaagaaataa tcgtgaaaaa taaagatgaa 1140  
 gaagagaagt tagttagtaa taacatcctt cataatcaac aagagttacc tacagctctt 1200  
 actaaattgg ttaaagagga tgaagttgtg tcttcagaaa aagcaaaaaga cagttttaat 1260  
 gaaaagagag ttgcagtgga agctcctatg agggaggaat atgcagactt caaaccattt 1320  
 gagcgagtat gggaagtga agatagtaag gaagatagt atagtgtggc tgctggaggt 1380  
 aaaatcgaga gcaacttgga aagtaaagt gataaaaaat gttttgcaga tagccttgag 1440  
 caaactaatc acgaaaaaga tagtgagagt agtaaatgat atacttcttt cccagtagc 1500  
 ccagaaggtc taaaggatcg ttcaggagca tatatcacat gtgtccctt taaccagca 1560  
 gcaactgaga gcattgcaac aaacattttt cctttgttag gagatcctt ttcagaaaat 1620  
 aagaccgatg aaaaaaaat agaagaaaag aaggcccaaa tagtaacaga gaagaatact 1680  
 agcaccaaaa catcaaaccc ttttcttgta gcagcacagg attctgagac agattatgtc 1740  
 acaacagata atttaacaaa ggtgactgag gaagtctggt caaacatgcc tgaaggcctg 1800  
 actccagatt tagtacagga agcatgtgaa agtgaattga atgaagttac tggtaaaaag 1860  
 attgcttatg aaacaaaaat ggacttgggt caaacatcag aagttatgca agagtcactc 1920

tatectgcag cacagctttg cccatcattt gaagagtcag aagctactcc ttcaccagtt 1980  
 ttgcctgaca ttgttatgga agcaccattg aattctgcag ttcctagtgc tgggtgcttcc 2040  
 gtgatacagc ccagctcatc accattagaa gcttcttcag ttaattatga aagcataaaa 2100  
 catgagcctg aaaaccccc accatatgaa gaggccatga gtgtatcact aaaaaaagta 2160  
 tcaggaataa aggaagaaat taaagagcct gaaaatatta atgcagctct tcaagaaaca 2220  
 gaagctcctt atatatctat tgcattgtgat ttaattaaag aaacaaagct ttctgctgaa 2280  
 ccagctccgg atttctctga ttattcagaa atggcaaaag ttgaacagcc agtgcctgat 2340  
 cattctgagc tagttgaaga ttcctcacct gattctgaac cagttgactt atttagtgat 2400  
 gattcaatac ctgacgttcc aaaaaacaa gatgaaactg tgatgcttgt gaaagaaagt 2460  
 ctactgaga cttcatttga gtcaatgata gaatatgaaa ataaggaaaa actcagtgtc 2520  
 ttgccacctg agggaggaaa gccatatattg gaattcttta agctcagttt agataacaca 2580  
 aaagataccc tgttacctga tgaagtttca acattgagca aaaaggagaa aattcctttg 2640  
 cagatggagg agctcagtag tgcagtttat tcaaatgatg acttatttat ttctaaggaa 2700  
 gcacagataa gagaaactga aacgttttca gattcatctc caattgaaat tatagatgag 2760  
 ttccctacat tgatcagttc taaaactgat tcattttcta aattagccag ggaatatact 2820  
 gacctagaag tatcccacaa aagtgaattt gtaaatgcc cggatggagc tgggtcattg 2880  
 ccttgccagc aattgcccc tgacctttct ttgaagaaca tacaacccaa agttgaagag 2940  
 aaaatcagtt tctcagatga cttttctaaa aatgggtctg ctacatcaaa ggtgctctta 3000  
 ttgcctccag atgtttctgc tttggccact caagcagaga tagagagcat agttaaaccc 3060  
 aaagtctctg tgaaagaagc tgagaaaaaa cttccttccg atacagaaaa agaggacaga 3120  
 tcaccatctg ctatattttc agcagagctg agtaaaactt cagttgttga cctcctgtac 3180  
 tggagagaca ttaagaagac tggagtgggt tttggtgcca gcctattcct gctgctttca 3240  
 ttgacagtat tcagcattgt gagcgtaaca gcctacattg ccttggccct gctctctgtg 3300  
 accatcagct ttaggatata caagggtgtg atccaagcta tccagaaatc agatgaaggc 3360  
 caccatttca gggcatatct ggaatctgaa gttgctatat ctgaggagtt ggttcagaag 3420  
 tacagtaatt ctgctcttgg tcatgtgaac tgcacgataa aggaactcag gcgcctcttc 3480  
 ttagttagtg atttagttga ttctctgaag tttgcagtgt tgatgtgggt atttacctat 3540  
 gttggtgcct tgtttaatgg tctgacacta ctgattttgg ctctcatttc actcttcagt 3600  
 gttcctgtta tttatgaacg gcatcaggca cagatagatc attatctagg acttgcaaat 3660  
 aagaatgtta aagatgctat ggctaaaaatc caagcaaaaa tccctggatt gaagcgcaaa 3720  
 gctgaatgaa aacgccccaa ataattagta ggagtcatc tttaaagggt atattcattt 3780  
 gattatacgg gggagggtca ggaagaacg aaccttgacg ttgcagtga gtttcacaga 3840  
 tcgtttagtg atctttattt ttagccatgc actgttgtga ggaaaaatta cctgtcttga 3900  
 ctgccatgtg ttcacatctc taagtattgt aagctgctat gtatggattt aaaccgtaat 3960  
 catatctttt tcctatctga ggcactggtg gaataaaaaa cctgtatatt ttactttgtt 4020  
 gcagatagtc ttgccgcac tttgcaagtt gcagagatgg tggagctaga aaaaaaaaaa 4080  
 aaaaaaaaaa aaa 4093

&lt;210&gt; 64

&lt;211&gt; 893

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 64

Met Gly Ser Ser Phe Ser Val Ser Pro Lys Ala Glu Ser Ala Val Ile  
 1 5 10 15

Val Ala Asn Pro Arg Glu Glu Ile Ile Val Lys Asn Lys Asp Glu Glu  
 20 25 30

Glu Lys Leu Val Ser Asn Asn Ile Leu His Asn Gln Gln Glu Leu Pro  
 35 40 45

Thr Ala Leu Thr Lys Leu Val Lys Glu Asp Glu Val Val Ser Ser Glu  
 50 55 60

Lys Ala Lys Asp Ser Phe Asn Glu Lys Arg Val Ala Val Glu Ala Pro  
 65 70 75 80

Met Arg Glu Glu Tyr Ala Asp Phe Lys Pro Phe Glu Arg Val Trp Glu



85	90	95
Val Lys Asp Ser Lys Glu Asp Ser Asp Met Leu Ala Ala Gly Gly Lys 100 105 110		
Ile Glu Ser Asn Leu Glu Ser Lys Val Asp Lys Lys Cys Phe Ala Asp 115 120 125		
Ser Leu Glu Gln Thr Asn His Glu Lys Asp Ser Glu Ser Ser Asn Asp 130 135 140		
Asp Thr Ser Phe Pro Ser Thr Pro Glu Gly Ile Lys Asp Arg Ser Gly 145 150 155 160		
Ala Tyr Ile Thr Cys Ala Pro Phe Asn Pro Ala Ala Thr Glu Ser Ile 165 170 175		
Ala Thr Asn Ile Phe Pro Leu Leu Gly Asp Pro Thr Ser Glu Asn Lys 180 185 190		
Thr Asp Glu Lys Lys Ile Glu Glu Lys Lys Ala Gln Ile Val Thr Glu 195 200 205		
Lys Asn Thr Ser Thr Lys Thr Ser Asn Pro Phe Leu Val Ala Ala Gln 210 215 220		
Asp Ser Glu Thr Asp Tyr Val Thr Thr Asp Asn Leu Thr Lys Val Thr 225 230 235 240		
Glu Glu Val Val Ala Asn Met Pro Glu Gly Leu Thr Pro Asp Leu Val 245 250 255		
Gln Glu Ala Cys Glu Ser Glu Leu Asn Glu Val Thr Gly Thr Lys Ile 260 265 270		
Ala Tyr Glu Thr Lys Met Asp Leu Val Gln Thr Ser Glu Val Met Gln 275 280 285		
Glu Ser Leu Tyr Pro Ala Ala Gln Leu Cys Pro Ser Phe Glu Glu Ser 290 295 300		
Glu Ala Thr Pro Ser Pro Val Leu Pro Asp Ile Val Met Glu Ala Pro 305 310 315 320		
Leu Asn Ser Ala Val Pro Ser Ala Gly Ala Ser Val Ile Gln Pro Ser 325 330 335		
Ser Ser Pro Leu Glu Ala Ser Ser Val Asn Tyr Glu Ser Ile Lys His 340 345 350		
Glu Pro Glu Asn Pro Pro Pro Tyr Glu Glu Ala Met Ser Val Ser Leu 355 360 365		
Lys Lys Val Ser Gly Ile Lys Glu Glu Ile Lys Glu Pro Glu Asn Ile 370 375 380		
Asn Ala Ala Leu Gln Glu Thr Glu Ala Pro Tyr Ile Ser Ile Ala Cys 385 390 395 400		
Asp Leu Ile Lys Glu Thr Lys Leu Ser Ala Glu Pro Ala Pro Asp Phe		

405	410	415
Ser Asp Tyr Ser Glu Met Ala Lys Val Glu Gln Pro Val Pro Asp His		
420	425	430
Ser Glu Leu Val Glu Asp Ser Ser Pro Asp Ser Glu Pro Val Asp Leu		
435	440	445
Phe Ser Asp Asp Ser Ile Pro Asp Val Pro Gln Lys Gln Asp Glu Thr		
450	455	460
Val Met Leu Val Lys Glu Ser Leu Thr Glu Thr Ser Phe Glu Ser Met		
465	470	475
Ile Glu Tyr Glu Asn Lys Glu Lys Leu Ser Ala Leu Pro Pro Glu Gly		
485	490	495
Gly Lys Pro Tyr Leu Glu Ser Phe Lys Leu Ser Leu Asp Asn Thr Lys		
500	505	510
Asp Thr Leu Leu Pro Asp Glu Val Ser Thr Leu Ser Lys Lys Glu Lys		
515	520	525
Ile Pro Leu Gln Met Glu Glu Leu Ser Thr Ala Val Tyr Ser Asn Asp		
530	535	540
Asp Leu Phe Ile Ser Lys Glu Ala Gln Ile Arg Glu Thr Glu Thr Phe		
545	550	555
Ser Asp Ser Ser Pro Ile Glu Ile Ile Asp Glu Phe Pro Thr Leu Ile		
565	570	575
Ser Ser Lys Thr Asp Ser Phe Ser Lys Leu Ala Arg Glu Tyr Thr Asp		
580	585	590
Leu Glu Val Ser His Lys Ser Glu Ile Ala Asn Ala Pro Asp Gly Ala		
595	600	605
Gly Ser Leu Pro Cys Thr Glu Leu Pro His Asp Leu Ser Leu Lys Asn		
610	615	620
Ile Gln Pro Lys Val Glu Glu Lys Ile Ser Phe Ser Asp Asp Phe Ser		
625	630	635
Lys Asn Gly Ser Ala Thr Ser Lys Val Leu Leu Leu Pro Pro Asp Val		
645	650	655
Ser Ala Leu Ala Thr Gln Ala Glu Ile Glu Ser Ile Val Lys Pro Lys		
660	665	670
Val Leu Val Lys Glu Ala Glu Lys Lys Leu Pro Ser Asp Thr Glu Lys		
675	680	685
Glu Asp Arg Ser Pro Ser Ala Ile Phe Ser Ala Glu Leu Ser Lys Thr		
690	695	700
Ser Val Val Asp Leu Leu Tyr Trp Arg Asp Ile Lys Lys Thr Gly Val		
705	710	715
Val Phe Gly Ala Ser Leu Phe Leu Leu Leu Ser Leu Thr Val Phe Ser		
720		

725	730	735
Ile Val Ser Val Thr Ala Tyr Ile	Ala Leu Ala Leu Leu Ser Val Thr	
740	745	750
Ile Ser Phe Arg Ile Tyr Lys Gly Val Ile Gln Ala Ile Gln Lys Ser		
755	760	765
Asp Glu Gly His Pro Phe Arg Ala Tyr Leu Glu Ser Glu Val Ala Ile		
770	775	780
Ser Glu Glu Leu Val Gln Lys Tyr Ser Asn Ser Ala Leu Gly His Val		
785	790	795
Asn Cys Thr Ile Lys Glu Leu Arg Arg Leu Phe Leu Val Asp Asp Leu		
805	810	815
Val Asp Ser Leu Lys Phe Ala Val Leu Met Trp Val Phe Thr Tyr Val		
820	825	830
Gly Ala Leu Phe Asn Gly Leu Thr Leu Leu Ile Leu Ala Leu Ile Ser		
835	840	845
Leu Phe Ser Val Pro Val Ile Tyr Glu Arg His Gln Ala Gln Ile Asp		
850	855	860
His Tyr Leu Gly Leu Ala Asn Lys Asn Val Lys Asp Ala Met Ala Lys		
865	870	875
Ile Gln Ala Lys Ile Pro Gly Leu Lys Arg Lys Ala Glu		
885	890	

&lt;210&gt; 65

&lt;211&gt; 3033

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 65

```

atacttcata gctgtggaag atgacaaaat tttaccatta aattcagctg aaaggaaacc 60
tggtgtggaag catgcaccat atataagcat tgcagggtgat gatcctcctg caagctgtgt 120
gtttagtcaa gttatgaaca tggcagcctt cctagccctt gtggtagctg ttctgcgctt 180
catacaactg aaaccgaagg ttttaaaccg gtggctgaat attagtggat tgggtgctct 240
gtgtctggct tccttcggaa tgaccttact tggtaatttt cagctcacia atgatgaaga 300
aatccataac gtcggaactt ccttgacctt tggatttggc acattgacct gctggatcca 360
ggctgcgctg acactcaagg tcaacatcaa gaatgaagga cggagagttg gaattccacg 420
ggttattctg tcggcatcta tcaactctctg tgtggtcttc tacttcatcc tcatggccca 480
aagcatccac atgtatgcag ccagggtcca gtggggcctg gtcattgtgt tctgtcttta 540
ttttggcacc tttgccgtgg agttccggca ttaccgctat gagattgttt gctctgagta 600
ccaggagaat ttctaagct tctcagaaag cctgtcagaa gcttctgaat atcagactga 660
ccagggttaa accatcagtt ttcccttctg ggtgaggtgg gtgtgacagt gggggagggg 720
ccagtaggac acactcacag gacttgacat agaacctcat ttcacacaca cacacacaca 780
cacattcatg gccacatttg ccaaatagagc ttttcagggc gagttatttc tttaatgaaa 840
aagcacaaagc ccttatgtgt cgaaatacac gctgttacac tgaaaatata tgcacgacag 900
agcaagaagc ttgtgcatga tcaactctta tccgtccctt tcccagcact cctcctctt 960
ccattctctt ccacatgtct caagcaccct accgagtagg gcaggccaaa tgttccttgg 1020
gagtaatgcc aactcccgac gttgccttca ggtccaaagg gcttggaacc agctcgtag 1080
gaagttctga atctggcact aatattcttg agtggataat agtgtatcat agaataggac 1140
ggaaaattgta ttgagatgtg accctgtgtc gcctgtggaa aggcatagtg agaagaactt 1200
tcccacgaaa gcccccttca tcgttgttca gtggtcggct gtgtggatcc caggagagac 1260

```

```

atatgccaca gactgtgaga gcaaagcccg ccgctgtgat ctggacttga tgcactgtga 1320
ctgagaatga tttccaaatg tgaatatgtg tagggacgtg gtctatcagg cctggaacaa 1380
gatgggggca gtaaggtat ggttttagtg ttgctttcat agtatgccat gtacaatggt 1440
ttatatttca tagtttcttt taagtaacta ccatgagtct ctctaagcct catggacaaa 1500
gatgtagacc aaatgcaaga gctgagcttg ctttgggttc aacctgac aaagaaaaac 1560
tgaggtcacc tgcaggctta cgtgggaagc taagaaaata tcacagaaaa atggaaagaa 1620
atatggatgt gcagaggtaa taaaaagagc tgctgctgga catttatcaa agataggcac 1680
cttacagtat cttgttgcaa gtccaaatca ggggagactt tgttaatatg tatcttttgc 1740
tcaaatttag tagggttttt ttttttttgg gattaaagggc ttttcaaatt tggaatttgt 1800
agaagccaat taaaagaata cccttcctaa aaaccaaat gacacagctg gccccctgc 1860
tagccaaccc gcagttccca gttcccacct gctgagagtt cagaactcag accgtagggtg 1920
tgggcttatg agactgtgct tcatgaagaa aagcagaatt ccaaattcaa actgttgagg 1980
tgagccaagt ggacagaccc tccatgagt gactttcttc caagaaatcc ccagatttac 2040
cccatagagg tctgggatta cctggaatat aatatgaaaa acatttttta ggctgggtgt 2100
ggggctcaca cctgtaatcc cagcactttg ggaggctgag gcgggcggat cacctgaggt 2160
tgggagctcg agaccagcct gaccaatgtg aagaaacccc gtcttacta aaaaaaatc 2220
aaaattagcc tgggtgtgtg gcgcatgcct gtaatcccgg ctactcagga ggcggaggca 2280
ggagaatcac ttgaacccgg gaggcaggtt gcggtgagac gagatcacgc cattgcactc 2340
cagcctgggc aaaaagagct aaaaactcca tctcaaaaaa aaaaagaaaa aaagaaaaac 2400
gttttttaat tcaacccaaa gtctgaattt cttgaggaag aggtcaatgt cttttcatct 2460
ctgtattgtt cccacctct taacacagt gctggcacat actaggtgct caataaatgt 2520
ctgttgaaat gaattattaa taatttaaga gtaggaaaaa ggagtgtcag agagagactc 2580
aaagatgagt gagagagcat ggaagatggc tgtttcatgt gagacaggaa tggaaagaag 2640
tatgaatagg gatcagatgg gggagtgaga atccctgtaa gtaaaatgtg aggaaaagaa 2700
caagctgaga acaagattct cctgtcaaaa aggcgtctta ctgtaaatag tccaaggtga 2760
catttgtagt ttttgaatac tgcttttggg tttctttttt tcatttttat attttaaaat 2820
ttttatcaaa gaacaaagac ttatgtagtt ttcttttatt ctacacaatc caaattaaac 2880
agcttttggg gatgcaattg tacactttct taagaatata tctaatacca catttttagca 2940
gaaccaagca atgactgaca ttattcattg ggatctggcc actaaataaa attcttttat 3000
gcataaaaaa aaaaaaaaaa aataaaaaaa aaa 3033

```

&lt;210&gt; 66

&lt;211&gt; 178

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 66

```

Met Asn Met Ala Ala Phe Leu Ala Leu Val Val Ala Val Leu Arg Phe
  1                      5                      10                     15

Ile Gln Leu Lys Pro Lys Val Leu Asn Pro Trp Leu Asn Ile Ser Gly
      20                      25                     30

Leu Val Ala Leu Cys Leu Ala Ser Phe Gly Met Thr Leu Leu Gly Asn
      35                      40                     45

Phe Gln Leu Thr Asn Asp Glu Glu Ile His Asn Val Gly Thr Ser Leu
      50                      55                     60

Thr Phe Gly Phe Gly Thr Leu Thr Cys Trp Ile Gln Ala Ala Leu Thr
      65                      70                     75                     80

Leu Lys Val Asn Ile Lys Asn Glu Gly Arg Arg Val Gly Ile Pro Arg
      85                      90                     95

Val Ile Leu Ser Ala Ser Ile Thr Leu Cys Val Val Leu Tyr Phe Ile
      100                     105                    110

Leu Met Ala Gln Ser Ile His Met Tyr Ala Ala Arg Val Gln Trp Gly
      115                     120                    125

```

Leu Val Met Cys Phe Leu Ser Tyr Phe Gly Thr Phe Ala Val Glu Phe  
 130 135 140

Arg His Tyr Arg Tyr Glu Ile Val Cys Ser Glu Tyr Gln Glu Asn Phe  
 145 150 155 160

Leu Ser Phe Ser Glu Ser Leu Ser Glu Ala Ser Glu Tyr Gln Thr Asp  
 165 170 175

Gln Val

<210> 67

<211> 504

<212> DNA

<213> Homo sapiens

<400> 67

aaaatccctg ctcccagctg ctttactaaa gagcaagttc ctgggcatct ctgtgtttct 60  
 ctttatgggg ttcaaaacct ttcaaggacc tctctccatg ccactgggtc cttggacct 120  
 atcactgggc tgctctctga gcccctcagt cctaccacag tctactgact tttcccatc 180  
 agctgtgagc attcaacct gtcccctgga ccttgacacc tggctcccca accctgtccc 240  
 aggaaaccca gattccacca gacacttct tcttccccc gaggtatct ggcctgagac 300  
 aacaaatgct gcctcccacc ctgagtctgg cactgggact ttcagaactc ctccttccct 360  
 gactctttgc cccagaccg tcattcaatg gctagctttt tccatgggaa aaacacgagc 420  
 accccaacc acaacggcca gttctctgat tccctaaatc cgcaccctt tcaaaacctc 480  
 aaaaaaaaa aaaaaaaaa aaaa 504

<210> 68

<211> 130

<212> PRT

<213> Homo sapiens

<400> 68

Met Gly Phe Lys Thr Phe Gln Gly Pro Leu Ser Met Pro Leu Val Pro  
 1 5 10 15  
 Trp Thr Leu Ser Leu Gly Cys Leu Leu Ser Pro Ser Val Leu Pro Gln  
 20 25 30  
 Ser Thr Asp Phe Ser His Ser Ala Val Ser Ile Gln Pro Cys Pro Leu  
 35 40 45  
 Asp Leu Asp Thr Trp Leu Pro Asn Pro Val Pro Gly Asn Pro Asp Ser  
 50 55 60  
 Thr Arg His Phe Leu Leu Pro Pro Glu Ala Ile Trp Pro Glu Thr Thr  
 65 70 75 80  
 Asn Ala Ala Ser His Pro Glu Ser Gly Thr Gly Thr Phe Arg Thr Pro  
 85 90 95  
 Pro Ser Leu Thr Leu Cys Pro Arg Pro Val Ile Gln Trp Leu Ala Phe  
 100 105 110  
 Ser Met Gly Lys Thr Arg Ala Pro Pro Thr Thr Thr Ala Ser Ser Leu  
 115 120 125

Ile Pro  
130

<210> 69  
<211> 3103  
<212> DNA  
<213> Homo sapiens

<400> 69

```

gatccgctgc tcttgtgacg ttgtggagat ggggagcgtc ctggggctgt gctccatggc 60
gagctggata ccatgtttgt gtggaagtgc cccgtgtttg ctatgccgat gctgtccatg 120
tggaacaac tccactgtaa ctagattgat ctatgcactt ttcttgcttg ttggagtagt 180
tgtagcttgt gtaatgttga taccaggaat ggaagaacaa ctgaataaga ttcctggatt 240
ttgtgagaat gagaaggtg ttgtcccttg taacattttg gttggctata aagctgtata 300
tcgtttgtgc ttgggtttgg ctatgttcta tcttcttctc tctttactaa tgatcaaatg 360
gaagagtagc agtgatccta gagctgcagt gcacaatgga ttttggttct ttaaatttgc 420
tgcagcaatt gcaattatta ttggggcatt cttcattcca gaaggaactt ttacaactgt 480
gtggttttat gtaggcatgg caggtgcctt ttgtttcacc ctcatacaac tagtcttact 540
tattgatatt gcacattcat ggaatgaatc gtgggttgaa aaaatggaag aaggggaactc 600
gagatgtttg tatgcagcct tgttatcagc tacagctctg aattatctgc tgtctttagt 660
tgctatcgtc ctgttctttg tctactacac tcatccagcc agttgttcag aaaacaagc 720
gttcatcagt gtcaacatgc tctctgcgt ttgtgcttct gtaatgtcta tactgccaaa 780
aatccaagaa tcacaaccaa gatctgggtt gttacagctc tcagtaatta cagtctacac 840
aatgtatttg acatggctcag ctatgaccaa tgaaccagaa acaaattgca acccaagtct 900
actaagcata attggctaca atacaacaag cactgtccca aaggaagggc agtcagtcca 960
gtgggtggcat gctcaaggaa ttataggact aattctcttt ttgttggtgt tattttatc 1020
cagcatccgt acttcaaaca atagtcaggt taataaactg actctaaca gtgatgaatc 1080
tacattaata gaagatggtg gagctagaag tgatggatca ctggaggatg gggacgatgt 1140
tcaccgagct gtagataatg aaagggatgg tgtcacttac agttattcct tctttcactt 1200
catgcttttc ctggcttcac ttatatcat gatgacctt accaactggg acaggtatga 1260
accctctcgt gagatgaaaa gtcagtggac agctgtctgg gtgaaaatct cttccagttg 1320
gattggcacc gtgctgtatg ttggacact cgtggcacca cttgttctta caaatcgtga 1380
ttttgactga gtgagacttc tagcatgaaa gtcccacttt gattattgct tatttgaaaa 1440
cagtattccc aacttttga aagttgtgta tgtttttgct tcccatgtaa cttctccagt 1500
gttctggcat gaattagatt ttactgcttg tcatattgtt attttcttac caagtgcatt 1560
gatattgtgaa gtagaatgaa ttgcagagga aagttttatg aatatggtga tgagttagta 1620
aaagtggcca ctattgggct tattctctgc tctatagttg tgaaatgaag agtgaaaaa 1680
aatttggttg actattttaa aattatatta gaccttaagc tgttttagca agcattaaag 1740
caaatgtatg gctgcctttt gaaatatttg atgtgttgcc tggcaggata ctgcaaagaa 1800
catgggttat tttaaaattt ataaacaagt cacttaaatg ccagtgtgct gaaaaatctt 1860
ataaggtttt acccttgata cggatttac acaggtaggg agtgtttagt ggacaatagt 1920
gtaggttatg gatggagggt tcggtactaa attgaataac gagtaataa tcttacttgg 1980
gtagagatgg cctttgcaa caaagtgaac tgttttggtt gttttaaact catgaagtat 2040
gggttcagtg gaaatgtttg gaactctgaa ggatttagac aaggttttga aaaggataat 2100
catgggttag aaggaagtgt ttgaaagtca ctttgaaggt tagttttggg ccagcacggt 2160
agctcaccct tgtaatccca gcactttggg aggtgaggt gggtagatta cttgagccca 2220
ggaattcgaag accagcctgg gcaacatggt gaaaccctgt ttctataaaa aataatctgg 2280
gctttgtagc atatgcctgt ggtcccagct actgaggagg ctgaggtggg aggattgctt 2340
gagcccagga ggcagagggt gcagttagcc aaggtcacgt cactgcactc tagcctgggc 2400
aacagagtaa gacaaaaaaa tatatatata ttgaaatcaa aggaggcaaa attttgacag 2460
ggaaggaagt aactgcaaaa cactaggctt tagtaggtac ttatataaaa tctagtccag 2520
ttctctcatt taaaaaaatg aagacactga agtacagact taaatagctc agatagctaa 2580
ttaggaaatt tcaagtgggc caataatagc attctctctg acatttataaa ataatttcta 2640
ttcaaaatc atgcataatt gattttacac ctcattactg gtggataatt tatgtgatgt 2700
ggattgctgg tgtccagcat gaccataaaa caggtcagaa gaatgatgga atgttttaga 2760
ataaactcct gcttatagta tactacacag ttcaaaagat gtttaaaatg cttttgtatt 2820
tactgccatg taattgaaat atatagatta ttgtaacctt tcaacctgaa aatcaagcag 2880
tatgagagtt tagttatttg tatgtgtcac tagtgtctaa tgaagctttt aaaaactaca 2940
atttcttctt taaaaatatt tattaatgtg aatggaatat aacaattcag ctttaattccc 3000

```

caaccttatt ctgtgtgttag acattgtatt ccacaatttt gaatggctgt gttttacctc 3060  
 taaataaatg aattcagaga aaaaaaaaaa aaaaaaaaaa aaa 3103

<210> 70  
 <211> 453  
 <212> PRT  
 <213> Homo sapiens

<400> 70  
 Met Gly Ser Val Leu Gly Leu Cys Ser Met Ala Ser Trp Ile Pro Cys  
 1 5 10 15  
 Leu Cys Gly Ser Ala Pro Cys Leu Leu Cys Arg Cys Cys Pro Ser Gly  
 20 25 30  
 Asn Asn Ser Thr Val Thr Arg Leu Ile Tyr Ala Leu Phe Leu Leu Val  
 35 40 45  
 Gly Val Cys Val Ala Cys Val Met Leu Ile Pro Gly Met Glu Glu Gln  
 50 55 60  
 Leu Asn Lys Ile Pro Gly Phe Cys Glu Asn Glu Lys Gly Val Val Pro  
 65 70 75 80  
 Cys Asn Ile Leu Val Gly Tyr Lys Ala Val Tyr Arg Leu Cys Phe Gly  
 85 90 95  
 Leu Ala Met Phe Tyr Leu Leu Leu Ser Leu Leu Met Ile Lys Val Lys  
 100 105 110  
 Ser Ser Ser Asp Pro Arg Ala Ala Val His Asn Gly Phe Trp Phe Phe  
 115 120 125  
 Lys Phe Ala Ala Ala Ile Ala Ile Ile Ile Gly Ala Phe Phe Ile Pro  
 130 135 140  
 Glu Gly Thr Phe Thr Thr Val Trp Phe Tyr Val Gly Met Ala Gly Ala  
 145 150 155 160  
 Phe Cys Phe Ile Leu Ile Gln Leu Val Leu Leu Ile Asp Phe Ala His  
 165 170 175  
 Ser Trp Asn Glu Ser Trp Val Glu Lys Met Glu Glu Gly Asn Ser Arg  
 180 185 190  
 Cys Trp Tyr Ala Ala Leu Leu Ser Ala Thr Ala Leu Asn Tyr Leu Leu  
 195 200 205  
 Ser Leu Val Ala Ile Val Leu Phe Phe Val Tyr Tyr Thr His Pro Ala  
 210 215 220  
 Ser Cys Ser Glu Asn Lys Ala Phe Ile Ser Val Asn Met Leu Leu Cys  
 225 230 235 240  
 Val Gly Ala Ser Val Met Ser Ile Leu Pro Lys Ile Gln Glu Ser Gln  
 245 250 255  
 Pro Arg Ser Gly Leu Leu Gln Ser Ser Val Ile Thr Val Tyr Thr Met  
 260 265 270

Tyr Leu Thr Trp Ser Ala Met Thr Asn Glu Pro Glu Thr Asn Cys Asn  
 275 280 285  
 Pro Ser Leu Leu Ser Ile Ile Gly Tyr Asn Thr Thr Ser Thr Val Pro  
 290 295 300  
 Lys Glu Gly Gln Ser Val Gln Trp Trp His Ala Gln Gly Ile Ile Gly  
 305 310 315 320  
 Leu Ile Leu Phe Leu Leu Cys Val Phe Tyr Ser Ser Ile Arg Thr Ser  
 325 330 335  
 Asn Asn Ser Gln Val Asn Lys Leu Thr Leu Thr Ser Asp Glu Ser Thr  
 340 345 350  
 Leu Ile Glu Asp Gly Gly Ala Arg Ser Asp Gly Ser Leu Glu Asp Gly  
 355 360 365  
 Asp Asp Val His Arg Ala Val Asp Asn Glu Arg Asp Gly Val Thr Tyr  
 370 375 380  
 Ser Tyr Ser Phe Phe His Phe Met Leu Phe Leu Ala Ser Leu Tyr Ile  
 385 390 395 400  
 Met Met Thr Leu Thr Asn Trp Tyr Arg Tyr Glu Pro Ser Arg Glu Met  
 405 410 415  
 Lys Ser Gln Trp Thr Ala Val Trp Val Lys Ile Ser Ser Ser Trp Ile  
 420 425 430  
 Gly Ile Val Leu Tyr Val Trp Thr Leu Val Ala Pro Leu Val Leu Thr  
 435 440 445  
 Asn Arg Asp Phe Asp  
 450  
 <210> 71  
 <211> 1981  
 <212> DNA  
 <213> Homo sapiens  
 <400> 71  
 gagatccaag ttgggagcag ctctgcgtgc ggggcctcag agaatgagggc cggcggttcgc 60  
 cctgtgcctc ctctggcagg cgctctggcc cgggcggggc ggcggcggaac accccactgc 120  
 cgaccgtgct ggctgctcgg cctcggggggc ctgctacagc ctgcaccacg ctaccatgaa 180  
 gcggcaggcg gccgaggagg cctgcatcct gcgaggtggg gcgctcagca ccgtgcgtgc 240  
 gggcgccgag ctgcgcgctg tgctcgcgct cctgcgggca ggcccagggc ccggaggggg 300  
 ctccaaagac ctgctgttct gggtcgcact ggagcgcagg cgttcccact gcaccctgga 360  
 gaacgagcct ttgcgggggt tctcctggct gtccctccgac cccggcggtc tcgaaagcga 420  
 cagctgcag tgggtggagg agccccaacg etcctgcacc gcgcggagat gcgcggtact 480  
 ccaggccacc ggtgggggtcg agcccgagg ctggaaggag atgcgatgcc acctgcgcgc 540  
 caacggctac ctgtgcaagt accagtttga ggtcttgtgt cctgcgcgcg gccccggggc 600  
 cgcctctaac ttgagctatc gcgcgcctt ccagctgcac agcgcgcgtc tggacttcag 660  
 tccacctggg accgaggtga gtgcgctctg ccggggacag ctcccgatct cagttacttg 720  
 catcgcggaac gaaatcggcg ctgcgtggga caaactctcg ggcgatgtgt tgtgtccctg 780  
 ccccgaggag tacctccgtg ctggcaaatg cgcagagctc cctaactgcc tagacgactt 840  
 gggaggcttt gcctgcgaat gtgctacggg cttcgagctg gggaaggacg gccgctcttg 900  
 tgtgaccagt ggggaaggac agccgaccct tggggggacc ggggtgcccc ccaggcgccc 960  
 gccggccact gcaaccagcc ccgtgccgca gagaacatgg ccaatcaggg tcgacgagaa 1020



```

gctgggagag acaccacttg tccctgaaca agacaattca gtaacatcta ttcctgagat 1080
tcctcgatgg ggatcacaga gcacgatgtc tacccttcaa atgtcccttc aagccgagtc 1140
aaaggccact atcaccccat cagggagcgt gatttccaag tttaattcta cgacttcctc 1200
tgccactcct caggctttcg actcctcctc tgccgtggtc ttcataatttg tgagcacagc 1260
agtagtagtg ttggtgatct tgaccatgac agtactgggg cttgtcaagc tctgctttca 1320
cgaaagcccc tcttcccagc caaggaagga gtctatgggc ccgcccgggc tggagagtga 1380
tcctgagccc gctgctttgg gctccagttc tgcacattgc acaacaatg gggtgaaagt 1440
cggggactgt gatctgcggg acagagcaga ggggtgccttg ctggcggagt cccctcttgg 1500
ctctagtgat gcatagggaa acaggggaca tgggcactcc tgtgaacagt ttttcacttt 1560
tgatgaaacg gggaaccaag aggaacttac ttgtgtaact gacaatttct gcagaaatcc 1620
cccttcctct aaattccctt tactccactg aggagctaaa tcagaactgc acactccttc 1680
cctgatgata gaggaagtgg aagtgccttt aggatgggtga tactggggga ccgggtagtg 1740
ctggggagag atattttctt atgtttattc ggagaatttg gagaagtgat tgaacttttc 1800
aagacattgg aaacaaatag aacacaatat aatttacatt aaaaaataat ttctaccaa 1860
atggaaaagga aatgttctat gttgttcagg ctaggagtat attggttcga aatcccaggg 1920
aaaaaaaaaa aaataaaaaa ttaaaggatt gttgataaaa aaaaaaaaaa aaaaaaaaaa 1980
a

```

1981

&lt;210&gt; 72

&lt;211&gt; 490

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 72

```

Met Arg Pro Ala Phe Ala Leu Cys Leu Leu Trp Gln Ala Leu Trp Pro
 1           5           10           15

Gly Pro Gly Gly Gly Glu His Pro Thr Ala Asp Arg Ala Gly Cys Ser
      20           25           30

Ala Ser Gly Ala Cys Tyr Ser Leu His His Ala Thr Met Lys Arg Gln
      35           40           45

Ala Ala Glu Glu Ala Cys Ile Leu Arg Gly Gly Ala Leu Ser Thr Val
      50           55           60

Arg Ala Gly Ala Glu Leu Arg Ala Val Leu Ala Leu Leu Arg Ala Gly
      65           70           75           80

Pro Gly Pro Gly Gly Gly Ser Lys Asp Leu Leu Phe Trp Val Ala Leu
      85           90           95

Glu Arg Arg Arg Ser His Cys Thr Leu Glu Asn Glu Pro Leu Arg Gly
      100          105          110

Phe Ser Trp Leu Ser Ser Asp Pro Gly Gly Leu Glu Ser Asp Thr Leu
      115          120          125

Gln Trp Val Glu Glu Pro Gln Arg Ser Cys Thr Ala Arg Arg Cys Ala
      130          135          140

Val Leu Gln Ala Thr Gly Gly Val Glu Pro Ala Gly Trp Lys Glu Met
      145          150          155          160

Arg Cys His Leu Arg Ala Asn Gly Tyr Leu Cys Lys Tyr Gln Phe Glu
      165          170          175

Val Leu Cys Pro Ala Pro Arg Pro Gly Ala Ala Ser Asn Leu Ser Tyr
      180          185          190

```

Arg Ala Pro Phe Gln Leu His Ser Ala Ala Leu Asp Phe Ser Pro Pro  
 195 200 205  
 Gly Thr Glu Val Ser Ala Leu Cys Arg Gly Gln Leu Pro Ile Ser Val  
 210 215 220  
 Thr Cys Ile Ala Asp Glu Ile Gly Ala Arg Trp Asp Lys Leu Ser Gly  
 225 230 235 240  
 Asp Val Leu Cys Pro Cys Pro Gly Arg Tyr Leu Arg Ala Gly Lys Cys  
 245 250 255  
 Ala Glu Leu Pro Asn Cys Leu Asp Asp Leu Gly Gly Phe Ala Cys Glu  
 260 265 270  
 Cys Ala Thr Gly Phe Glu Leu Gly Lys Asp Gly Arg Ser Cys Val Thr  
 275 280 285  
 Ser Gly Glu Gly Gln Pro Thr Leu Gly Gly Thr Gly Val Pro Thr Arg  
 290 295 300  
 Arg Pro Pro Ala Thr Ala Thr Ser Pro Val Pro Gln Arg Thr Trp Pro  
 305 310 315 320  
 Ile Arg Val Asp Glu Lys Leu Gly Glu Thr Pro Leu Val Pro Glu Gln  
 325 330 335  
 Asp Asn Ser Val Thr Ser Ile Pro Glu Ile Pro Arg Trp Gly Ser Gln  
 340 345 350  
 Ser Thr Met Ser Thr Leu Gln Met Ser Leu Gln Ala Glu Ser Lys Ala  
 355 360 365  
 Thr Ile Thr Pro Ser Gly Ser Val Ile Ser Lys Phe Asn Ser Thr Thr  
 370 375 380  
 Ser Ser Ala Thr Pro Gln Ala Phe Asp Ser Ser Ser Ala Val Val Phe  
 385 390 395 400  
 Ile Phe Val Ser Thr Ala Val Val Val Leu Val Ile Leu Thr Met Thr  
 405 410 415  
 Val Leu Gly Leu Val Lys Leu Cys Phe His Glu Ser Pro Ser Ser Gln  
 420 425 430  
 Pro Arg Lys Glu Ser Met Gly Pro Pro Gly Leu Glu Ser Asp Pro Glu  
 435 440 445  
 Pro Ala Ala Leu Gly Ser Ser Ser Ala His Cys Thr Asn Asn Gly Val  
 450 455 460  
 Lys Val Gly Asp Cys Asp Leu Arg Asp Arg Ala Glu Gly Ala Leu Leu  
 465 470 475 480  
 Ala Glu Ser Pro Leu Gly Ser Ser Asp Ala  
 485 490

&lt;210&gt; 73

&lt;211&gt; 3098

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 73

```

aggaaagcca gaacaagtgc ttcaagctgc cegtgtccgt ggtcaacacc aactgcccgt 60
gcgtggccta cgtgctgctg tcaactcgtg acttgcccgg cgtgctggcg gctgcccctgc 120
agctgcccgc cggcaccaag taccagcgtt tccccgactg gctggaccac tggctacagc 180
accgcaagca gatcgggctg ctcaagctct tctgcccgcg cctgcacgcc ctctacagct 240
tctgcttgcc gctgcccgcg gccacccgct acgacctggg caacctggca gtcaagcagg 300
tcttgcccaa caagagccac ctctgggtgg aggaggaggt ctggcggatg gagatctacc 360
tctccctggg agtgctggcc ctccggcagt tgcctctgct ggccgtgacc tcaactgccg 420
ccattgcaaa ctgctcaaac tggagggagt tcagcttctg tcagtcctca ctgggctttg 480
tggccctcgt gctgagcaca ctgcacacgc tcacctagg ctggaccgcg gccttcgagg 540
agagccgcta caagtcttac ctgcctccca ccttcacgct cagctgctg gtgcccctgcg 600
tcgtcatcct ggccaaagcc ctgtttctcc tgccctgcat cagccgcaga ctgcccagga 660
tccggagagg ctgggagagg gagagcaca tcaagttcac gctgcccaca gaccacgccc 720
tggccgagaa gacgagccac gtatgaggtg cctgcccctg gctctggacc ccgggacac 780
gagggacggt gccctgagcc cgttaggttt tcttttcttg gtggtgcaaa gtggtataac 840
tgtgtgcaaa taggaggttt gaggtccaaa ttccctggac tcaaatgtat gcagtactat 900
tcagaatgat atacacacat atgtgtatat gtatttacat atattccaca tatataacag 960
gatttgcaat tatacatagc tagctaaaaa gttgggtctc tgagatttca actttagat 1020
ttaaaaaaaa gtgccgtacg ttaagagaag agcagatcat gctattgtga catttgaga 1080
gatatacaca cactttttgt acagaagagg cttgtgctgt ggtgggttcg atttatccct 1140
gcccacccca tccccacaac ttcccttttg ctacttcccc aaggctcttg cagagctagg 1200
gctctgaagg ggaggggaag caacggctct gccagagcc atccctggag catgtgagca 1260
gcggctggtc tcttccctcc acctggggca gcagcaggag gcctggggag gaggaaaatc 1320
aggcagtcgg cctggagctt gtgcctggtc ctttgcccgg tgggtggagg atggagggat 1380
tgggtgaag ctgctccacc tcactcctgc tgagtggggg agacattttc cctgaaagtc 1440
agaagtacc atagagcctg caaatggatc ctctgtgag agtgacgtca cctcctttcc 1500
agagccatta gtgagcctgg cttgggaaca agtgtaattt ccttccctcc ttaacctgg 1560
cgatgagcgt cctttaaac actgtgcctt ctcacctttt ccactttcag tttgaacgac 1620
tcccagggaag gcctagagca gaccctttag aaatcagccc aagggggaga gcaagagaaa 1680
acactctagg gagtaagct ccccgggcgt cagagttgag cctgctggg gctgaaggac 1740
tgtcttcacg aagtcagtcc tgaggaaaaa tattggggac tccaaatgtc ctctggcaga 1800
ggaccagaaa aaccacactg gctccaactt cctcctcatg gggcattaca cttcaaaaca 1860
gtggggagca acttttccac caaagctaca aacctaaaat gctgctgccc caaagcaca 1920
gaggggaagag caccgcccgg gccacaggac gtctgtctc cagtcacagg ccactcctgc 1980
tgctccctac tgactctagc ttacttcccc tgtgaagaaa cagggtgttct cggctgagcc 2040
cccaaccctc tgcagaacca ggttgatctg ccacagaaaa agcatctttg aagacaaaga 2100
gggtgaggtc ttcatgagtc tcctggggcc aaagccatct tctgatggaa ggaagagagt 2160
agggccagtg aaggctgccc agagagaatg tcacagatga ggctgcccct gcccctccc 2220
cgccaggag gtttcatgag ctcatgtcta tgacgacat aagggttctt cagtgaagg 2280
caggagaaga gccactgca aggatagctc attaggcaca tgaccgatgc aggggaaggcc 2340
atgccgggga agctcttctt gcaggatatt tccatctgct gtgccaaggc tgagcggcag 2400
aaacttgtct cataaattgg cactgatgga gcatcagctg tggccacag agagccttg 2460
tgagaagggg gcaggtaaaag cagagatttt agcatctcct tggcataaca agggcccatc 2520
gattccctac taatgagagg caggagagac atgggcaatg gagaccacc aatgatcccc 2580
aaccgccggt ggtactggct gcctgcccgt ggccagggaa tggctcctta taccaaagat 2640
gctggcacat agcagaaccc agtgacgtc ctccccctcc caccacctc tggctgaagg 2700
tgctcaagag ggaagcaatt ataaggtggg tggcaggagg gaacagggtg cacctgctgg 2760
acaatcacac gaaaggcagg cgggctgtgt actgggcccct gactgtgctg ccactgctgt 2820
cttccctacc tcaccaggct actggcagca gcatcccag agcacatcat ctccacagcc 2880
tggtaaatc catgtgcctc tgggtacaaa agtgcctcaa cgacatgtc tggaaatccc 2940
aaatgccaca gtctgaggtt gatattctaa atctatgcct tcaaaagagt ctctgttttt 3000
ttttttttaa cctggtagac ggtataaaag cagtgcacaa aaacacctaa ccttctgcaa 3060
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa

```

3098

&lt;210&gt; 74

&lt;211&gt; 132

&lt;212&gt; PRT

<213> Homo sapiens

<400> 74

```

Met Glu Ile Tyr Leu Ser Leu Gly Val Leu Ala Leu Gly Thr Leu Ser
 1             5             10             15

Leu Leu Ala Val Thr Ser Leu Pro Ser Ile Ala Asn Ser Leu Asn Trp
          20             25             30

Arg Glu Phe Ser Phe Val Gln Ser Ser Leu Gly Phe Val Ala Leu Val
      35             40             45

Leu Ser Thr Leu His Thr Leu Thr Tyr Gly Trp Thr Arg Ala Phe Glu
 50             55             60

Glu Ser Arg Tyr Lys Phe Tyr Leu Pro Pro Thr Phe Thr Leu Thr Leu
65             70             75             80

Leu Val Pro Cys Val Val Ile Leu Ala Lys Ala Leu Phe Leu Leu Pro
          85             90             95

Cys Ile Ser Arg Arg Leu Ala Arg Ile Arg Arg Gly Trp Glu Arg Glu
          100             105             110

Ser Thr Ile Lys Phe Thr Leu Pro Thr Asp His Ala Leu Ala Glu Lys
115             120             125

Thr Ser His Val
130

```

<210> 75

<211> 922

<212> DNA

<213> Homo sapiens

<400> 75

```

gtgaaatgag aaagggccag aacgtgcagg tctgcggagg ggaagtgtcc tgagtgaagg 60
aggggacccc catcctgggg gatgctggga gtgagtgagt gagtgagatg gctgagtgag 120
ggttatgggg agcctgaggt tttatgggcc tgtgtatccc cttctcccg cccagcctg 180
cctccctect gcccgcctgg ccacaggtc tccctctggt cctgtccct ctggtgggtg 240
gggatggagc ggcagcaagg ggtgtaatgg ggctgggttc tgtcttctac aggccacccc 300
gaggtcctca gtggttgctt ggggagccgg acggggctcc tgaggggtac aggttgggtg 360
ggcctccctt gaggtcttgg ggtcaggctt tggcctctgc tgcctctcag tcaccaagtc 420
acctccctct gaaaatccag tcccttcttt ggatgtcctt gtgagtcact ctgggccttg 480
ctgtcgtccc tctcagctt cttgttctg ggacaagggt caagccagga tgggcccagg 540
cctgggatcc cccaccccag gacccccagg cccctcccc tgetgctttg cggggggcag 600
ggcagaaatg gactcctttt gggtcctcga ggtgggtccc cctccagccc ctgcatectc 660
cgtgccgtag acctgtctcc cagaggaggg gccttgaccc acaggacgtg tgggtggcgc 720
tggcactcag ggacccccag ctgccccagc cctggtctct ggcgcatctc ttcctctctg 780
tcccgaagat ctgcgcctct agtgcctttt gaggggttcc catcatccct cctgatattt 840
gtattgaaaa tattatgcac actgttcatg cttctactaa tcaataaacg ctttatttaa 900
aaaaaaaaaa aaaaaaaaaa aa
922

```

<210> 76

<211> 106

<212> PRT

<213> Homo sapiens

<400> 76

Met Gly Leu Cys Ile Pro Phe Ser Arg Pro Gln Pro Ala Ser Leu Leu  
 1 5 10 15  
 Pro Ala Trp Pro Thr Gly Leu Pro Leu Val Pro Val Pro Leu Val Val  
 20 25 30  
 Gly Asp Gly Ala Ala Ala Arg Gly Val Met Gly Leu Gly Ser Val Phe  
 35 40 45  
 Tyr Arg Pro Pro Arg Gly Pro Gln Trp Leu Pro Gly Glu Pro Asp Gly  
 50 55 60  
 Ala Pro Glu Gly Tyr Arg Leu Gly Gly Pro Ser Leu Arg Val Trp Gly  
 65 70 75 80  
 Gln Ala Leu Ala Ser Ala Ala Ser Gln Ser Pro Ser His Leu Pro Leu  
 85 90 95  
 Lys Ile Gln Ser Leu Leu Trp Met Ser Leu  
 100 105

<210> 77  
 <211> 823  
 <212> DNA  
 <213> Homo sapiens

<400> 77  
 agcccccctgc agcaccgcgt ggctacctca ccttggtgcc cctgctggcc accatgctct 60  
 tcatcatggg ctacgccgtg ggctggggtc ccatcacctg gctgctcatg tctgaggtcc 120  
 tgcccctgcg tgcccgtggc gtggcctcag ggtctctcgt gctggccagc tggctcaccg 180  
 ccttcgtcct caccaagtcc ttctgccag tggtagcac ctctggcctc caggtgcctt 240  
 tctttcttct cgcgccatc tgcttggtga gcctggtgtt cacaggctgc tgtgtgcccg 300  
 agaccaaggg acggtccctg gagcagatcg agtccttctt ccgcacgggg agaaggtcct 360  
 tcttgcgcta ggtcaaggtc cccgcctgga gggggccaaa ccccagtggt ctgggcctct 420  
 gtgttggtg tgcaaggaag acccggtctt gccctcaca gtcttatggg caccacaggg 480  
 aacatcctgg acttaaaaag ccagggcagg ccgggcacag tggctcacgc ctgtaatccc 540  
 agcacttttg gaggccaaag caggtggatt acccaaggcc aggagttaa gaccagcctg 600  
 gccaacatgg tgaacccccg tctctactaa aaaatacaaa aaagctgggt gtggtggcac 660  
 acaccgtag ttccagctac ttgggaggt gaggcagcat tgcttgaacc cgggaggtgg 720  
 aggctgcaat gagctgagat catgccattg cactccagcc tgggcaacga gagtgaaact 780  
 ccgtcccccac ccctggccaa aaaaaaaaaa aaaaaaaaaa aaa 823

<210> 78  
 <211> 105  
 <212> PRT  
 <213> Homo sapiens

<400> 78  
 Met Leu Phe Ile Met Gly Tyr Ala Val Gly Trp Gly Pro Ile Thr Trp  
 1 5 10 15  
 Leu Leu Met Ser Glu Val Leu Pro Leu Arg Ala Arg Gly Val Ala Ser  
 20 25 30  
 Gly Leu Cys Val Leu Ala Ser Trp Leu Thr Ala Phe Val Leu Thr Lys  
 35 40 45  
 Ser Phe Leu Pro Val Val Ser Thr Phe Gly Leu Gln Val Pro Phe Phe  
 50 55 60

Phe Phe Ala Ala Ile Cys Leu Val Ser Leu Val Phe Thr Gly Cys Cys  
 65 70 75 80

Val Pro Glu Thr Lys Gly Arg Ser Leu Glu Gln Ile Glu Ser Phe Phe  
 85 90 95

Arg Thr Gly Arg Arg Ser Phe Leu Arg  
 100 105

<210> 79

<211> 1224

<212> DNA

<213> Homo sapiens

<400> 79

aatgaactgg gagctgctgc tgtggctgct ggtgctgtgc gcgctgctcc tgcctcttgg 60  
 gcagctgctg cgcttcctga gggctgacgg cgacctgacg ctactatggg ccgagtggca 120  
 gggacgacgc ccagaatggg agctgactga tatggtgggtg tgggtgactg gagcctcgag 180  
 tggaattggg gaggagctgg cttaccagtt gtctaaacta ggagtttctc ttgtgctgtc 240  
 agccagaaga gtgcatgagc tggaaagggt gaaaagaaga tgcctagaga atggcaattt 300  
 aaaagaaaaa gatatacttg ttttgccctc tgacctgacc gacctgggtt cccatgaagc 360  
 ggctacccaa gctgttctcc aggagtttgg tagaatcgac attctgggtca acaatgggtg 420  
 aatgtcccg cgttctctgt gcatggatac cagcttggat gtctacagaa agctaataga 480  
 gcttaactac ttagggacgg tgccttgac aaaatgtgtt ctgcctcaca tgatcgagag 540  
 gaagcaagga aagattgtta ctgtgaatag catcctgggt atcatatctg tacctcttcc 600  
 cattggatac tgtgctagca agcatgctct ccgggggtttt tttaatggcc ttcgaacaga 660  
 acttgccaca taccaggta taatagtctc taacatttgc ccaggacctg tgcaatcaaa 720  
 tattgtggag aattccctag ctggagaagt cacaaagact ataggcaata atggagacca 780  
 gtcccacaag atgacaacca gtcgttgtgt gcggctgatg ttaatcagca tggccaatga 840  
 tttgaaagaa gtttggatct cagaacaacc tttcttggtta gtaacatatt tgtggcaata 900  
 catgcccaacc tgggcctggt ggataaccaa caagatgggg aagaaaagga ttgagaactt 960  
 taagagtggg gtggatgcag actcttctta ttttaaaatc ttttaagacaa aacatgactg 1020  
 aaaagagcac ctgtactttt caagccactg gagggagaaa tggaaaacat gaaaacagca 1080  
 atcttcttat gcttctgaat aatcaaagac taatttgtga ttttactttt taatagatat 1140  
 gactttgctt ccaacatgga atgaaataaa aaataaataa taaaagattg ccatgaatct 1200  
 tgcaaaaaaa aaaaaaaaaa aaaa 1224

<210> 80

<211> 339

<212> PRT

<213> Homo sapiens

<400> 80

Met Asn Trp Glu Leu Leu Trp Leu Leu Val Leu Cys Ala Leu Leu  
 1 5 10 15

Leu Leu Leu Val Gln Leu Leu Arg Phe Leu Arg Ala Asp Gly Asp Leu  
 20 25 30

Thr Leu Leu Trp Ala Glu Trp Gln Gly Arg Arg Pro Glu Trp Glu Leu  
 35 40 45

Thr Asp Met Val Val Trp Val Thr Gly Ala Ser Ser Gly Ile Gly Glu  
 50 55 60

Glu Leu Ala Tyr Gln Leu Ser Lys Leu Gly Val Ser Leu Val Leu Ser  
 65 70 75 80

Ala Arg Arg Val His Glu Leu Glu Arg Val Lys Arg Arg Cys Leu Glu  
                     85                    90                    95  
 Asn Gly Asn Leu Lys Glu Lys Asp Ile Leu Val Leu Pro Leu Asp Leu  
                     100                    105                    110  
 Thr Asp Thr Gly Ser His Glu Ala Ala Thr Lys Ala Val Leu Gln Glu  
                     115                    120                    125  
 Phe Gly Arg Ile Asp Ile Leu Val Asn Asn Gly Gly Met Ser Gln Arg  
                     130                    135                    140  
 Ser Leu Cys Met Asp Thr Ser Leu Asp Val Tyr Arg Lys Leu Ile Glu  
                     145                    150                    155                    160  
 Leu Asn Tyr Leu Gly Thr Val Ser Leu Thr Lys Cys Val Leu Pro His  
                     165                    170                    175  
 Met Ile Glu Arg Lys Gln Gly Lys Ile Val Thr Val Asn Ser Ile Leu  
                     180                    185                    190  
 Gly Ile Ile Ser Val Pro Leu Ser Ile Gly Tyr Cys Ala Ser Lys His  
                     195                    200                    205  
 Ala Leu Arg Gly Phe Phe Asn Gly Leu Arg Thr Glu Leu Ala Thr Tyr  
                     210                    215                    220  
 Pro Gly Ile Ile Val Ser Asn Ile Cys Pro Gly Pro Val Gln Ser Asn  
                     225                    230                    235                    240  
 Ile Val Glu Asn Ser Leu Ala Gly Glu Val Thr Lys Thr Ile Gly Asn  
                     245                    250                    255  
 Asn Gly Asp Gln Ser His Lys Met Thr Thr Ser Arg Cys Val Arg Leu  
                     260                    265                    270  
 Met Leu Ile Ser Met Ala Asn Asp Leu Lys Glu Val Trp Ile Ser Glu  
                     275                    280                    285  
 Gln Pro Phe Leu Leu Val Thr Tyr Leu Trp Gln Tyr Met Pro Thr Trp  
                     290                    295                    300  
 Ala Trp Trp Ile Thr Asn Lys Met Gly Lys Lys Arg Ile Glu Asn Phe  
                     305                    310                    315                    320  
 Lys Ser Gly Val Asp Ala Asp Ser Ser Tyr Phe Lys Ile Phe Lys Thr  
                     325                    330                    335  
 Lys His Asp

&lt;210&gt; 81

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

<220>  
<223> oligonucleotide

<400> 81  
tcagtcacag aggcattcaa g

21

<210> 82  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 82  
ctgattgtgc attgtatctg g

21

<210> 83  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 83  
tcacggacac agaggaagg

19

<210> 84  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 84  
ccctagtgcc ctgtcacc

18

<210> 85  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 85  
tcagggtttc atctcggg

18

<210> 86  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 86  
ggagttcatc tatgcctccg

20



<210> 87  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 87  
ggcaaccagt agcatcagg

19

<210> 88  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 88  
tgaactcgga gtgctctgg

19

<210> 89  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 89  
ctggcttcag taccctcg

18

<210> 90  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 90  
acgccctggt ggaggacctc

20

<210> 91  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 91  
aagagttcaa ctattggggc

20

<210> 92  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 92  
tgtgaaattg tctgtagaaa g 21

<210> 93  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 93  
ctactctgga gtcagccgc 19

<210> 94  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 94  
aacattgtgg aaacagtgac c 21

<210> 95  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 95  
ggttggagag tcagactcg 20

<210> 96  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 96  
tgtgagagca cagactgcg 19

<210> 97  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 97

ctgagttgat tagaatgctg 20

<210> 98  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 98  
tgtacctcaa gcacagccc 19

<210> 99  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 99  
cacatcacia agggcacc 18

<210> 100  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 100  
agacccttgg acctggcag 19

<210> 101  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 101  
gaacctggag caaggacac 19

<210> 102  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 102  
cccgtggaag tagagagcc 19

<210> 103  
<211> 21  
<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 103

ttcaatgggt gaattgtatg g

21

<210> 104

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 104

tggaatagca tgactgggtg

20

<210> 105

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 105

tctcagaagc atcaccaact g

21

<210> 106

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 106

attctgtctt tctcttct c

21

<210> 107

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 107

cctcactgag ggtgaactgc

20

<210> 108

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 108  
tctgagggaa gaagggaaatg 20

<210> 109  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 109  
ttcaacagag ctgggcaag 19

<210> 110  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 110  
tgtcgtccgg tcaactgtg 18

<210> 111  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 111  
ctcataaagc aagagctggc 20

<210> 112  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 112  
acggaagcac cagcactag 19

<210> 113  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 113  
cccaaaagca gtattcaaaa c 21

<210> 114  
<211> 18

<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 114  
accttgacac ctggctcc 18

<210> 115  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 115  
actttcctct gcaattcatt c 21

<210> 116  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 116  
tcacttctcc aaattctccg 20

<210> 117  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 117  
tgggtctctg agatttcaac 20

<210> 118  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 118  
tcactcactc ccagcatcc 19

<210> 119  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 119  
cagacatgag cagccagg 18

<210> 120  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 120  
tgcctcacat gatcgagag 19

<210> 121  
<211> 51  
<212> PRT  
<213> Homo sapiens

<400> 121  
Met Gly Leu Lys Asn Ser Arg Phe Trp Glu Pro Ser Met Leu Ser Leu  
1 5 10 15  
Ser Pro Leu Leu Ser Thr Ser Leu Pro Asn Glu Arg Val Thr Glu Asn  
20 25 30  
Cys Phe Phe Ile Asn Arg Ser Phe Leu Ile Val Ser Gly Phe Asp Thr  
35 40 45  
Ser Val Val  
50

<210> 122  
<211> 101  
<212> PRT  
<213> Homo sapiens

<400> 122  
Met Glu Thr Val Met Thr Leu Glu Gly Gly Gln Lys Pro Val Met Glu  
1 5 10 15  
Ser Tyr Thr Leu Leu Leu Ser Pro Pro Glu Thr Gln Asp Glu Gly Trp  
20 25 30  
Thr Leu Ala Leu Ser Leu Ala Leu Thr Tyr Arg Leu Gly Pro Gly Ser  
35 40 45  
Val Leu Leu Ala Pro Arg Ser Pro Trp Arg Leu Gln Lys Pro Pro Pro  
50 55 60  
Ala Gly Gly Leu Pro His Ser Gln Leu Pro Ala Arg Gln Trp Gly Leu  
65 70 75 80  
Leu Arg Gln Ala Gly Pro Lys Thr Val Ser Pro Leu Gly Thr Tyr Ser  
85 90 95  
Pro Pro Ser Ile Cys  
100

<210> 123  
 <211> 90  
 <212> PRT  
 <213> Homo sapiens

<400> 123  
 Met Arg Leu Ser Thr Val Gly Thr Val His Ser Gly Pro Pro Leu Ser  
           1                  5                  10                  15  
 Ser Gly Leu Leu Thr Gly Ala Pro Cys His Val Val Thr Met Ala His  
                   20                  25                  30  
 Arg Pro Leu Leu Leu Leu Leu Leu Leu Pro Pro Glu Glu Leu Arg  
           35                  40                  45  
 Leu Cys Phe Gly Leu Leu Val Ser Gly Gln Phe Leu Val Val Leu Leu  
           50                  55                  60  
 Arg Leu Arg Gly Pro Asp Leu Phe Leu His His Arg Ile Val His Leu  
           65                  70                  75                  80  
 Val Phe Leu Thr Leu Arg Phe Leu Ala Cys  
                   85                  90

<210> 124  
 <211> 325  
 <212> PRT  
 <213> Homo sapiens

<400> 124  
 Met Gly Gly Ala Leu Cys Asp Val Val Leu Gln Thr Glu Arg Lys Gly  
           1                  5                  10                  15  
 Gly Ser Val Leu Ser Tyr Ile Asn Gly Val Thr Thr Ser Lys Pro Gly  
                   20                  25                  30  
 Val Ser Leu Val Tyr Ser Met Pro Ser Arg Asn Leu Ser Leu Arg Leu  
           35                  40                  45  
 Glu Gly Leu Gln Glu Lys Asp Ser Gly Pro Tyr Ser Cys Ser Val Asn  
           50                  55                  60  
 Val Gln Asp Lys Gln Gly Lys Ser Arg Gly His Ser Ile Lys Thr Leu  
           65                  70                  75                  80  
 Glu Leu Asn Val Leu Val Pro Pro Ala Pro Pro Ser Cys Arg Leu Gln  
                   85                  90                  95  
 Gly Val Pro His Val Gly Ala Asn Val Thr Leu Ser Cys Gln Ser Pro  
           100                  105                  110  
 Arg Ser Lys Pro Ala Val Gln Tyr Gln Trp Asp Arg Gln Leu Pro Ser  
           115                  120                  125  
 Phe Gln Thr Phe Phe Ala Pro Ala Leu Asp Val Ile Arg Gly Ser Leu  
           130                  135                  140



Ser Leu Thr Asn Leu Ser Ser Ser Met Ala Gly Val Tyr Val Cys Lys  
 145 150 155 160  
 Ala His Asn Glu Val Gly Thr Ala Gln Cys Asn Val Thr Leu Glu Val  
 165 170 175  
 Ser Thr Gly Pro Gly Ala Ala Val Val Ala Gly Ala Val Val Gly Thr  
 180 185 190  
 Leu Val Gly Leu Gly Leu Leu Ala Gly Leu Val Leu Leu Tyr His Arg  
 195 200 205  
 Arg Gly Lys Ala Leu Glu Glu Pro Ala Asn Asp Ile Lys Glu Asp Ala  
 210 215 220  
 Ile Ala Pro Arg Thr Leu Pro Trp Pro Lys Ser Ser Asp Thr Ile Ser  
 225 230 235 240  
 Lys Asn Gly Thr Leu Ser Ser Val Thr Ser Ala Arg Ala Leu Arg Pro  
 245 250 255  
 Pro His Gly Pro Pro Arg Pro Gly Ala Leu Thr Pro Thr Pro Ser Leu  
 260 265 270  
 Ser Ser Gln Ala Leu Pro Ser Pro Arg Leu Pro Thr Thr Asp Gly Ala  
 275 280 285  
 His Pro Gln Pro Ile Ser Pro Ile Pro Gly Gly Val Ser Ser Ser Gly  
 290 295 300  
 Leu Ser Arg Met Gly Ala Val Pro Val Met Val Pro Ala Gln Ser Gln  
 305 310 315 320  
 Ala Gly Ser Leu Val  
 325

&lt;210&gt; 125

&lt;211&gt; 127

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 125

Met Ile Ser Leu Pro Gly Pro Leu Val Thr Asn Leu Leu Arg Phe Leu  
 1 5 10 15  
 Phe Leu Gly Leu Ser Ala Leu Ala Pro Pro Ser Arg Ala Gln Leu Gln  
 20 25 30  
 Leu His Leu Pro Ala Asn Arg Leu Gln Ala Val Glu Gly Gly Glu Val  
 35 40 45  
 Val Leu Pro Ala Trp Tyr Thr Leu His Gly Glu Val Ser Ser Ser Gln  
 50 55 60  
 Pro Trp Glu Val Pro Phe Val Met Trp Phe Phe Lys Gln Lys Glu Lys  
 65 70 75 80  
 Glu Asp Gln Cys Cys Pro Thr Ser Met Gly Ser Gln Gln Ala Asn Leu  
 85 90 95

Glu Tyr Pro Trp Ser Thr Pro Cys Pro Pro Gly Thr Cys Pro Cys Gly  
 100 105 110

Trp Arg Val Ser Arg Arg Lys Thr Leu Ala Pro Thr Ala Ala Pro  
 115 120 125

<210> 126

<211> 91

<212> PRT

<213> Homo sapiens

<400> 126

Met Asn Gly Val Thr His His Val Ala Phe Cys Val Trp Phe Leu Pro  
 1 5 10 15

Leu Ser Ile Met Ile Ser Ser Ser Cys Arg Ser Met His Leu Ser Phe  
 20 25 30

Ile Pro Ser His Gly Ala Ile Ile Phe His Cys Val Tyr Lys Thr Thr  
 35 40 45

Phe Cys Gln Ser Thr His Pro Trp Ile Thr Ser Ala Phe Leu Lys Gly  
 50 55 60

Ala Phe Thr Ser Gln Thr Gln Phe Ser Val Gly Glu Gly Val Arg Gly  
 65 70 75 80

Gln Tyr Arg Ser Asp Cys Lys Arg His Lys Leu  
 85 90

<210> 127

<211> 84

<212> PRT

<213> Homo sapiens

<400> 127

Met Leu Phe Pro Ser Ser Ser Lys Pro Phe Ser Leu Leu Ser Leu  
 1 5 10 15

Thr Ile Trp Ala Arg Leu Val Gly Arg Leu Thr Asn Arg Ile Cys Pro  
 20 25 30

Val Pro Pro Gly Ser Val Ala Ser Ser Met Ser Leu Gln Ala Gly Arg  
 35 40 45

Cys Gly Asn Pro Val Val Leu Pro Gln Pro Met Pro Pro Gly Leu Leu  
 50 55 60

Cys Met Asn Glu Cys Ser Leu Val Pro Gly Leu Gly Arg Gly Gln Val  
 65 70 75 80

Asn Ser Arg Val

<210> 128

<211> 120

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 128

```

Met Arg Ala Thr His Cys Gln Ala Ala Arg Met Phe Val Leu Phe Ser
 1             5             10             15

Arg Glu Gly Asn Pro Thr Gly Phe Cys Gly Asn Asn Gly Asn Leu Gln
          20             25             30

Met Pro Ala Trp Asp Asp Glu Ala His Ser Glu Gln Met Leu Phe Phe
          35             40             45

Phe Leu Arg Gln Ser Leu Ala Leu Thr Pro Arg Leu Glu Cys Ser Gly
          50             55             60

Ala Ile Ser Ala His Cys Lys Leu Cys Leu Pro Gly Ser Ser Asp Ser
          65             70             75             80

Pro Thr Ser Ala Ser Arg Val Ala Gly Ile Thr Pro Pro Cys Pro Ala
          85             90             95

Asn Phe Cys Val Phe Ser Arg Asp Gly Val Ser Pro Cys Trp Pro Gly
          100            105            110

Trp Ser Arg Thr Pro Asp Leu Arg
          115            120

```

&lt;210&gt; 129

&lt;211&gt; 143

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 129

```

Met Ala Glu Gly Glu Lys Asn Gln Asp Phe Thr Phe Lys Lys Glu Ser
 1             5             10             15

Pro Ser Asp Ser Ala Val Val Leu Pro Ser Thr Pro Gln Ala Ser Ala
          20             25             30

Asn Pro Ser Ser Pro Tyr Thr Asn Ser Ser Arg Lys Gln Pro Met Ser
          35             40             45

Ala Thr Leu Arg Glu Arg Leu Arg Lys Thr Arg Phe Ser Phe Asn Ser
          50             55             60

Ser Tyr Asn Val Val Lys Arg Leu Lys Val Glu Ser Glu Glu Asn Asp
          65             70             75             80

Gln Thr Phe Ser Glu Lys Pro Ala Ser Ser Thr Glu Glu Asn Cys Leu
          85             90             95

Glu Phe Gln Glu Ser Phe Lys His Ile Asp Ser Glu Phe Glu Glu Asn
          100            105            110

Thr Asn Leu Lys Asn Thr Leu Thr Ala Ile Ser Met Ser Val Asn Leu
          115            120            125

Ser His Leu Ile Leu Asp His Ala Val Leu Ser Lys Met Ser Leu

```

130                      135                      140  
  
 <210> 130  
 <211> 284  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 130  
 Met Glu Asp Leu Asp Gln Ser Pro Leu Val Ser Ser Ser Asp Ser Pro  
   1                          5                          10                          15  
  
 Pro Arg Pro Gln Pro Ala Phe Lys Tyr Gln Phe Val Arg Glu Pro Glu  
                   20                          25                          30  
  
 Asp Glu Glu Glu Glu Glu Glu Glu Glu Glu Asp Glu Asp Glu Asp  
           35                          40                          45  
  
 Leu Glu Glu Leu Glu Val Leu Glu Arg Lys Pro Ala Ala Gly Leu Ser  
   50                          55                          60  
  
 Ala Ala Pro Val Pro Thr Ala Pro Ala Ala Gly Ala Pro Leu Met Asp  
   65                          70                          75                          80  
  
 Phe Gly Asn Asp Phe Val Pro Pro Ala Pro Arg Gly Pro Leu Pro Ala  
                   85                          90                          95  
  
 Ala Pro Pro Val Ala Pro Glu Arg Gln Pro Ser Trp Asp Pro Ser Pro  
                   100                          105                          110  
  
 Val Ser Ser Thr Val Pro Ala Pro Ser Pro Leu Ser Ala Ala Ala Val  
   115                          120                          125  
  
 Ser Pro Ser Lys Leu Pro Glu Asp Asp Glu Pro Pro Ala Arg Pro Pro  
   130                          135                          140  
  
 Pro Pro Pro Pro Ala Ser Val Ser Pro Gln Ala Glu Pro Val Trp Thr  
   145                          150                          155                          160  
  
 Pro Pro Ala Pro Ala Pro Ala Ala Pro Pro Ser Thr Pro Ala Ala Pro  
                   165                          170                          175  
  
 Lys Arg Arg Gly Ser Ser Gly Ser Val Asp Glu Thr Leu Phe Ala Leu  
                   180                          185                          190  
  
 Pro Ala Ala Ser Glu Pro Val Ile Arg Ser Ser Ala Glu Asn Met Asp  
   195                          200                          205  
  
 Leu Lys Glu Gln Pro Gly Asn Thr Ile Ser Ala Gly Gln Glu Asp Phe  
   210                          215                          220  
  
 Pro Ser Val Leu Leu Glu Thr Ala Ala Ser Leu Pro Ser Leu Ser Pro  
   225                          230                          235                          240  
  
 Leu Ser Ala Ala Ser Phe Lys Glu His Glu Tyr Leu Gly Asn Leu Ser  
                   245                          250                          255  
  
 Thr Val Leu Pro Thr Glu Gly Thr Leu Gln Glu Asn Val Ser Glu Ala  
   260                          265                          270

Ser Lys Glu Val Ser Glu Lys Ala Lys Leu Tyr Ser  
 275 280

<210> 131  
 <211> 175  
 <212> PRT  
 <213> Homo sapiens

<400> 131  
 Met Arg Pro Ala Phe Ala Leu Cys Leu Leu Trp Gln Ala Leu Trp Pro  
 1 5 10 15

Gly Pro Gly Gly Gly Glu His Pro Thr Ala Asp Arg Ala Gly Cys Ser  
 20 25 30

Ala Ser Gly Ala Cys Tyr Ser Leu His His Ala Thr Met Lys Arg Gln  
 35 40 45

Ala Ala Glu Glu Ala Cys Ile Leu Arg Gly Gly Ala Leu Ser Thr Val  
 50 55 60

Arg Ala Gly Ala Glu Leu Arg Ala Val Leu Ala Leu Leu Arg Ala Gly  
 65 70 75 80

Pro Gly Pro Gly Gly Gly Ser Lys Asp Leu Leu Phe Trp Val Ala Leu  
 85 90 95

Glu Arg Arg Arg Ser His Cys Thr Leu Glu Asn Glu Pro Leu Arg Gly  
 100 105 110

Phe Ser Trp Leu Ser Ser Asp Pro Gly Gly Leu Glu Ser Asp Thr Leu  
 115 120 125

Gln Trp Val Glu Glu Pro Gln Arg Ser Cys Thr Arg Ala Glu Met Arg  
 130 135 140

Gly Thr Pro Gly His Arg Trp Gly Arg Ala Arg Arg Leu Glu Gly Asp  
 145 150 155 160

Ala Met Pro Pro Ala Arg Gln Arg Leu Pro Val Gln Val Pro Val  
 165 170 175

<210> 132  
 <211> 147  
 <212> PRT  
 <213> Homo sapiens

<400> 132  
 Met Leu Phe Ile Met Gly Tyr Ala Val Gly Trp Gly Pro Ile Thr Trp  
 1 5 10 15

Leu Leu Met Ser Glu Val Leu Pro Leu Arg Ala Arg Gly Val Ala Ser  
 20 25 30

Gly Leu Cys Val Leu Ala Ser Trp Leu Thr Ala Phe Val Leu Thr Lys  
 35 40 45

Ser Phe Leu Pro Val Val Ser Thr Phe Gly Leu Gln Val Pro Phe Phe

[illegible]

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/19351

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>																										
IPC(6) : C07H 21/04; C07K 14/705; C12N 15/09; 15/63; C12Q 1/68 US CL : 536/23.1, 24.3; 435/7.2, 69.1, 320.1; 530/350, 300 According to International Patent Classification (IPC) or to both national classification and IPC																										
<b>B. FIELDS SEARCHED</b>																										
Minimum documentation searched (classification system followed by classification symbols) U.S. : 536/23.1, 24.3; 435/7.2, 69.1, 320.1; 530/350, 300																										
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched																										
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) MEDLINE, JAPIO, BIOSIS, WPIDS, CAPLUS, EMBASE, N-GENESEQ35, N-ISSUED, EMBL-EST58, PIR60, SWISS-PROT37, SPTREMBL19, EMBL58 search terms: secreted proteins, 98862, 98846, VB11																										
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>																										
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																								
X	Database EMBASE EST-58 , Accession No. AA458581, NID g2183488, HILLIER et al., 'aa12e01.rl Soares-NhHMPu-S1 Homo sapiens cDNA clone IMAGE:813048 5' similar to contains element PTR5 repetitive element; mRNA sequence, 9 June, 1997	1-11																								
0 X	Database EMBASE EST 158, Accession No. Z99943, NID g2887308, PEARCE, A., 'Human DNA sequence from PAC 313L4 on chromosome 1q24. Contains ESTs.' 12 February 1998	44-45																								
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.																										
<table border="0"> <tr> <td colspan="2">* Special categories of cited documents:</td> <td>*T</td> <td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>*A</td> <td>document defining the general state of the art which is not considered to be of particular relevance</td> <td>*X</td> <td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>*E</td> <td>earlier document published on or after the international filing date</td> <td>*Y</td> <td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>*L</td> <td>document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>*A</td> <td>document member of the same patent family</td> </tr> <tr> <td>*O</td> <td>document referring to an oral disclosure, use, exhibition or other means</td> <td></td> <td></td> </tr> <tr> <td>*P</td> <td>document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			* Special categories of cited documents:		*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A	document defining the general state of the art which is not considered to be of particular relevance	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E	earlier document published on or after the international filing date	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A	document member of the same patent family	*O	document referring to an oral disclosure, use, exhibition or other means			*P	document published prior to the international filing date but later than the priority date claimed		
* Special categories of cited documents:		*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																							
*A	document defining the general state of the art which is not considered to be of particular relevance	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																							
*E	earlier document published on or after the international filing date	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																							
*L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A	document member of the same patent family																							
*O	document referring to an oral disclosure, use, exhibition or other means																									
*P	document published prior to the international filing date but later than the priority date claimed																									
Date of the actual completion of the international search 16 NOVEMBER 1999		Date of mailing of the international search report 11 JAN 2000																								
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer NIRMAL S. BASI Telephone No. (703) 308-0196																								

**INTERNATIONAL SEARCH REPORT**

International application No.  
PCT/US99/19351

**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:  
1-11 and 44-45
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/19351

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s)1-11, drawn to polynucleotide comprising SEQ ID NO:1, fragments thereof, expression vector containing said sequence, cell transformed with said vector, polypeptide of SEQ ID NO:2, fragments of the polypeptide of SEQ ID NO:2 and process for preparing said polypeptide.

Group II, claim(s)12-13, drawn to polynucleotide comprising SEQ ID NO:3, fragments thereof, polypeptide of SEQ ID NO:4 and fragments of the polypeptide of said polypeptide.

Group III, claim(s)14-15, drawn to polynucleotide comprising SEQ ID NO:5, fragments thereof, polypeptide of SEQ ID NO:6 and fragments of the polypeptide of said polypeptide.

Group IV, claim(s)16-17, drawn to polynucleotide comprising SEQ ID NO:7, fragments thereof, polypeptide of SEQ ID NO:8 and fragments of the polypeptide of said polypeptide.

Group V, claim(s)18-19, drawn to polynucleotide comprising SEQ ID NO:9, fragments thereof, polypeptide of SEQ ID NO:10 and fragments of the polypeptide of said polypeptide.

Group VI, claim(s)20-21, drawn to polynucleotide comprising SEQ ID NO:11, fragments thereof, polypeptide of SEQ ID NO:12 and fragments of the polypeptide of said polypeptide.

Group VII, claim(s)22-23, drawn to polynucleotide comprising SEQ ID NO:13, fragments thereof, polypeptide of SEQ ID NO:14 and fragments of the polypeptide of said polypeptide.

Group VIII, claim(s)24-25, drawn to polynucleotide comprising SEQ ID NO:15, fragments thereof, polypeptide of SEQ ID NO:16 and fragments of the polypeptide of said polypeptide.

Group IX, claim(s)26-27, drawn to polynucleotide comprising SEQ ID NO:17, fragments thereof, polypeptide of SEQ ID NO:18 and fragments of the polypeptide of said polypeptide.

Group X, claim(s)28-29, drawn to polynucleotide comprising SEQ ID NO:19, fragments thereof, polypeptide of SEQ ID NO:20 and fragments of the polypeptide of said polypeptide.

Group XI, claim(s)30-31, drawn to polynucleotide comprising SEQ ID NO:21, fragments thereof, polypeptide of SEQ ID NO:22 and fragments of the polypeptide of said polypeptide.

Group XII, claim(s)32-33, drawn to polynucleotide comprising SEQ ID NO:23, fragments thereof, polypeptide of SEQ ID NO:24 and fragments of the polypeptide of said polypeptide.

Group XIII, claim(s)34-35, drawn to polynucleotide comprising SEQ ID NO:25, fragments thereof, polypeptide of SEQ ID NO:26 and fragments of the polypeptide of said polypeptide.

Group XIV, claim(s)36-37, drawn to polynucleotide comprising SEQ ID NO:27, fragments thereof, polypeptide of SEQ ID NO:28 and fragments of the polypeptide of said polypeptide.

Group XV, claim(s)38-39, drawn to polynucleotide comprising SEQ ID NO:29, fragments thereof, polypeptide of SEQ ID NO:30 and fragments of the polypeptide of said polypeptide.

Group XVI, claim(s)40-41, drawn to polynucleotide comprising SEQ ID NO:31, fragments thereof, polypeptide of SEQ ID NO:32 and fragments of the polypeptide of said polypeptide.

Group XVII, claim(s)42-43, drawn to polynucleotide comprising SEQ ID NO:33, fragments thereof, polypeptide of SEQ ID NO:34 and fragments of the polypeptide of said polypeptide.

Group XVIII, claim(s)44-45, drawn to polynucleotide comprising SEQ ID NO:35, fragments thereof, polypeptide of SEQ ID NO:36 and fragments of the polypeptide of said polypeptide.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/19351

Group XIX, claim(s)46-47, drawn to polynucleotide comprising SEQ ID NO:37, fragments thereof, polypeptide of SEQ ID NO:38 and fragments of the polypeptide of said polypeptide.

Group XX, claim(s)48-49, drawn to polynucleotide comprising SEQ ID NO:39, fragments thereof, polypeptide of SEQ ID NO:40 and fragments of the polypeptide of said polypeptide.

Group XXI, claim(s)50-51, drawn to polynucleotide comprising SEQ ID NO:41, fragments thereof, polypeptide of SEQ ID NO:42 and fragments of the polypeptide of said polypeptide.

Group XXII, claim(s)52-53, drawn to polynucleotide comprising SEQ ID NO:43, fragments thereof, polypeptide of SEQ ID NO:44 and fragments of the polypeptide of said polypeptide.

Group XXIII, claim(s)54-55, drawn to polynucleotide comprising SEQ ID NO:45, fragments thereof, polypeptide of SEQ ID NO:46 and fragments of the polypeptide of said polypeptide.

Group XXIV, claim(s)56-57, drawn to polynucleotide comprising SEQ ID NO:47, fragments thereof, polypeptide of SEQ ID NO:48 and fragments of the polypeptide of said polypeptide.

Group XXV, claim(s)58-59, drawn to polynucleotide comprising SEQ ID NO:49, fragments thereof, polypeptide of SEQ ID NO:50 and fragments of the polypeptide of said polypeptide.

Group XXVI, claim(s)60-61, drawn to polynucleotide comprising SEQ ID NO:51, fragments thereof, polypeptide of SEQ ID NO:52 and fragments of the polypeptide of said polypeptide.

Group XXVII, claim(s)62-63, drawn to polynucleotide comprising SEQ ID NO:53, fragments thereof, polypeptide of SEQ ID NO:54 and fragments of the polypeptide of said polypeptide.

Group XXVIII, claim(s)64-65, drawn to polynucleotide comprising SEQ ID NO:55, fragments thereof, polypeptide of SEQ ID NO:56 and fragments of the polypeptide of said polypeptide.

Group XXIX, claim(s)66-67, drawn to polynucleotide comprising SEQ ID NO:57, fragments thereof, polypeptide of SEQ ID NO:58 and fragments of the polypeptide of said polypeptide.

Group XXX, claim(s)68-69, drawn to polynucleotide comprising SEQ ID NO:59, fragments thereof, polypeptide of SEQ ID NO:60 and fragments of the polypeptide of said polypeptide.

Group XXXI, claim(s)70-71, drawn to polynucleotide comprising SEQ ID NO:61, fragments thereof, polypeptide of SEQ ID NO:62 and fragments of the polypeptide of said polypeptide.

Group XXXII, claim(s)72-73, drawn to polynucleotide comprising SEQ ID NO:63, fragments thereof, polypeptide of SEQ ID NO:64 and fragments of the polypeptide of said polypeptide.

Group XXXIII, claim(s)74-75, drawn to polynucleotide comprising SEQ ID NO:65, fragments thereof, polypeptide of SEQ ID NO:66 and fragments of the polypeptide of said polypeptide.

Group XXXIV, claim(s)76-77, drawn to polynucleotide comprising SEQ ID NO:67, fragments thereof, polypeptide of SEQ ID NO:68 and fragments of the polypeptide of said polypeptide.

Group XXXV, claim(s)78-79, drawn to polynucleotide comprising SEQ ID NO:69, fragments thereof, polypeptide of SEQ ID NO:70 and fragments of the polypeptide of said polypeptide.

Group XXXVI, claim(s)80-81, drawn to polynucleotide comprising SEQ ID NO:71, fragments thereof, polypeptide of SEQ ID NO:72 and fragments of the polypeptide of said polypeptide.

Group XXXVII, claim(s)82-83, drawn to polynucleotide comprising SEQ ID NO:73, fragments thereof, polypeptide of SEQ ID NO:74 and fragments of the polypeptide of said polypeptide.

Group XXXVIII, claim(s)84-85, drawn to polynucleotide comprising SEQ ID NO:75, fragments thereof, polypeptide of SEQ ID NO:76 and fragments of the polypeptide of said polypeptide.

Group XXXIX, claim(s)86-87, drawn to polynucleotide comprising SEQ ID NO:77, fragments thereof, polypeptide of

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/19351

SEQ ID NO:78 and fragments of the polypeptide of said polypeptide.

Group XL, claim(s)88-89, drawn to polynucleotide comprising SEQ ID NO:79, fragments thereof, polypeptide of SEQ ID NO:80 and fragments of the polypeptide of said polypeptide.

The inventions listed as Groups I-XL do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The main invention is Group I, which is first product, first method of making the product and first method of using the product. Pursuant to 37 CFR 1.474 (d), these claims are considered by the ISA/US to constitute the main invention. The products of Groups II-XL do not share the same or corresponding special technical feature with group I because they are drawn to products having materially different structures and functions, each defines a separate invention over the art. Therefore, the claims are not linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.